

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

NAME OF DRUG PRODUCT : Mirtazapine Orodispersible Tablets 15 mg.
Mirtazapine Orodispersible Tablets 30 mg.
Mirtazapine Orodispersible Tablets 45 mg.

(TRADE) NAME OF PRODUCT : AUROZAPINE OD 15
AUROZAPINE OD 30
AUROZAPINE OD 45

STRENGTH : 15 mg, 30 mg and 45 mg.

PHARMACEUTICAL DOSAGE FORM: Orodispersible tablet.

2. QUALITATIVE AND QUANTITATIVE COMPOSITIONS:

Mirtazapine Orodispersible Tablets 15 mg:

Each Orodispersible tablet contains Mirtazapine Ph.Eur. 15 mg.

Mirtazapine Orodispersible Tablets 30 mg:

Each Orodispersible tablet contains Mirtazapine Ph.Eur. 30 mg.

Mirtazapine Orodispersible Tablets 45 mg:

Each Orodispersible tablet contains Mirtazapine Ph.Eur. 45 mg.

Excipient (s) of known effect (s): Mannitol

For full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM:

Mirtazapine Orodispersible Tablets 15 mg: White round tablets debossed with '36' on one side and 'A' on the other side with an embossed circular edge.

Mirtazapine Orodispersible Tablets 30 mg: White round tablets debossed with '37' on one side and 'A' on the other side with an embossed circular edge.

Mirtazapine Orodispersible Tablets 45 mg: White round tablets debossed with '38' on one side and 'A' on the other side with an embossed circular edge.

4. CLINICAL PARTICULARS:

4.1. Therapeutic indications

Mirtazapine is indicated in adults for the treatment of episodes of major depression.

4.2.Posology and method of administration

Posology

Adults

The effective daily dose is usually between 15 and 45 mg; the starting dose is 15 or 30 mg. Mirtazapine begins to exert its effect in general after 1-2 weeks of treatment. Treatment with an adequate dose should result in a positive response within 2-4 weeks. With an insufficient response, the dose can be increased up to the maximum dose. If there is no response within a further 2-4 weeks, then treatment should be stopped.

Patients with depression should be treated for a sufficient period of at least 6 months to ensure that they are free from symptoms.

It is recommended to discontinue treatment with mirtazapine gradually to avoid withdrawal symptoms (see section Special warnings and precautions for use).

Elderly

The recommended dose is the same as that for adults. In elderly patients an increase in dosing should be done under close supervision to elicit a satisfactory and safe response.

Renal impairment

The clearance of mirtazapine may be decreased in patients with moderate to severe renal impairment (creatinine clearance <40 ml/min). This should be taken into account when prescribing Mirtazapine to this category of patients (see section Special warnings and precautions for use).

Hepatic impairment

The clearance of mirtazapine may be decreased in patients with hepatic impairment. This should be taken into account when prescribing Mirtazapine Orodispersible to this category of patients, particularly with severe hepatic impairment, as patients with severe hepatic impairment have not been investigated (see section Special warnings and precautions for use).

Paediatric population

Mirtazapine should not be used in children and adolescents under the age of 18 years as efficacy was not demonstrated in two short-term clinical trials and because of safety concerns.

Method of administration

Mirtazapine has an elimination half-life of 20-40 hours and therefore Mirtazapine is suitable for once daily administration. It should be taken preferably as a single night-time dose before going to bed. Mirtazapine Orodispersible Tablets may also be given in two divided doses (once in the morning and once at night-time, the higher dose should be taken at night).

The tablets should be taken orally. The tablet will rapidly disintegrate and can be swallowed without water.

4.3.Contraindications

Hypersensitivity to active substance or any of the other ingredients of Mirtazapine Orodispersible Tablets.

Concomitant use of mirtazapine with monoamine oxidase (MAO) inhibitors.

4.4.Special warnings and precautions for use

Paediatric population

Mirtazapine Orodispersible Tablets should not be used in the treatment of children and adolescents under the age of 18 years. Suicide-related behaviours (suicide attempt and suicidal thoughts), and hostility (predominantly aggression, oppositional behaviour and anger) may occur more frequently observed in clinical trials among children and adolescents treated with antidepressants compared to those treated with placebo. If, based on clinical need, a decision to treat is nevertheless taken, the patient should be carefully monitored for the appearance of suicidal symptoms. In addition, long-term safety data in children and adolescents concerning growth, maturation and cognitive and behavioural development are lacking.

Suicide/suicidal thoughts or clinical worsening

Depression is associated with an increased risk of suicidal thoughts, self-harm and suicide (suicide-related events). This risk persists until significant remission occurs. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored until such improvement occurs. It is general clinical experience that the risk of suicide may increase in the early stages of recovery.

Patients with a history of suicide-related events or those exhibiting a significant degree of suicidal ideation prior to commencement of treatment are known to be at greater risk of suicidal thoughts or suicide attempts, and should receive careful monitoring during treatment. A meta-analysis of placebo-controlled clinical trials of antidepressants in adult patients with psychiatric disorders showed an increased risk of suicidal behaviour with

antidepressants compared to placebo in patients less than 25 years old.

Close supervision of patients and in particular those at high risk should accompany therapy with antidepressants especially in early treatment and following dose changes. Patients (and caregivers of patients) should be alerted about the need to monitor for any clinical worsening, suicidal behaviour or thoughts and unusual changes in behaviour and to seek medical advice immediately if these symptoms present.

With regard to the chance of suicide, in particular at the beginning of treatment, only the smallest amount of Mirtazapine Orodispersible Tablets should be given to the patient consistent with good patient management, in order to reduce the risk of overdose.

Bone marrow depression

Bone marrow depression, usually presenting as granulocytopenia or agranulocytosis, has been reported during treatment with Mirtazapine Orodispersible Tablets. Reversible agranulocytosis has been reported as a rare occurrence in clinical studies with Mirtazapine Orodispersible Tablets. In the postmarketing period with Mirtazapine Orodispersible Tablets very rare cases of agranulocytosis have been reported, mostly reversible, but in some cases fatal. Fatal cases mostly concerned patients with an age above 65. The physician should be alert for symptoms like fever, sore throat, stomatitis or other signs of infection; when such symptoms occur, treatment should be stopped and blood counts taken.

Jaundice

Treatment should be discontinued if jaundice occurs.

Conditions which need supervision

Careful dosing as well as regular and close monitoring is necessary in patients with:

- epilepsy and organic brain syndrome: Although clinical experience indicates that epileptic seizures are rare during mirtazapine treatment, as with other antidepressants, Mirtazapine Orodispersible Tablets should be introduced cautiously in patients who have a history of seizures. Treatment should be discontinued in any patient who develops seizures, or where there is an increase in seizure frequency.
- hepatic impairment: Following a single 15 mg oral dose of mirtazapine, the clearance of mirtazapine was approximately 35 % decreased in mild to moderate hepatically impaired patients, compared to subjects with normal hepatic function. The average plasma concentration of mirtazapine was about 55 % increased.
- renal impairment: Following a single 15 mg oral dose of mirtazapine, in patients with moderate (creatinine clearance <40 ml/min) and severe (creatinine clearance $\leq$ 10 ml/min) renal impairment the clearance of mirtazapine was about 30 % and 50 % decreased

respectively, compared to normal subjects. The average plasma concentration of mirtazapine was about 55 % and 115 % increased respectively. No significant differences were found in patients with mild renal impairment (creatinine clearance <80 ml/min) as compared to the control group.

- cardiac diseases like conduction disturbances, angina pectoris and recent myocardial infarction, where normal precautions should be taken and concomitant medicines carefully administered.

- low blood pressure.

- diabetes mellitus: In patients with diabetes, antidepressants may alter glycaemic control. Insulin and/or oral hypoglycaemic dosage may need to be adjusted and close monitoring is recommended.

Like with other antidepressants, the following should be taken into account:

- Worsening of psychotic symptoms can occur when antidepressants are administered to patients with schizophrenia or other psychotic disturbances; paranoid thoughts can be intensified

- When the depressive phase of bipolar disorder is being treated, it can transform into the manic phase. Patients with a history of mania/hypomania should be closely monitored. Mirtazapine should be discontinued in any patient entering a manic phase.

- Although Mirtazapine Orodispersible Tablets is not addictive, post-marketing experience shows that abrupt termination of treatment after long term administration may sometimes result in withdrawal symptoms. The majority of withdrawal reactions are mild and self-limiting. Among the various reported withdrawal symptoms, dizziness, agitation, anxiety, headache and nausea are the most frequently reported. Even though they have been reported as withdrawal symptoms, it should be realized that these symptoms may be related to the underlying disease. It is recommended to discontinue treatment with mirtazapine gradually.

- Care should be taken in patients with micturition disturbances like prostate hypertrophy and in patients with acute narrow-angle glaucoma and increased intra-ocular pressure (although there is little chance of problems with Mirtazapine Orodispersible Tablets because of its very weak anticholinergic activity).

- Akathisia/psychomotor restlessness: The use of antidepressants have been associated with the development of akathisia, characterized by a subjectively unpleasant or distressing restlessness and need to move often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop

these symptoms, increasing the dose may be detrimental.

– Cases of QT prolongation, Torsade de Pointes, ventricular tachycardia, and sudden death, have been reported during the post-marketing use of mirtazapine. The majority of reports occurred in association with overdose or in patients with other risk factors for QT prolongation, including concomitant use of QTc prolonging medicines. Caution should be exercised when Mirtazapine is prescribed in patients with known cardiovascular disease or family history of QT prolongation, and in concomitant use with other medicinal products thought to prolong the QTc interval.

Severe cutaneous adverse reactions

Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), bullous dermatitis and erythema multiforme, which can be life-threatening or fatal, have been reported in association with Mirtazapine Orodispersible Tablets treatment.

If signs and symptoms suggestive of these reactions appear, Mirtazapine Orodispersible Tablets should be withdrawn immediately.

If the patient has developed one of these reactions with the use of Mirtazapine Orodispersible Tablets, treatment with Mirtazapine Orodispersible Tablets must not be restarted in this patient at any time.

Hyponatraemia

Hyponatraemia, probably due to inappropriate antidiuretic hormone secretion (SIADH), has been reported very rarely with the use of mirtazapine. Caution should be exercised in patients at risk, such as elderly patients or patients concomitantly treated with medications known to cause hyponatraemia.

Serotonin syndrome

Interaction with serotonergic active substances: serotonin syndrome may occur when selective serotonin reuptake inhibitors (SSRIs) are used concomitantly with other serotonergic active substances. Symptoms of serotonin syndrome may be hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, mental status changes that include confusion, irritability and extreme agitation progressing to delirium and coma. Caution should be advised and a closer clinical monitoring is required when these active substances are combined with mirtazapine. Treatment with mirtazapine should be discontinued if such events occur and supportive symptomatic treatment initiated. From post marketing experience it appears that serotonin syndrome occurs very rarely in patients treated with Mirtazapine Orodispersible Tablets alone.

Elderly

Elderly are often more sensitive, especially with regard to the undesirable effects of antidepressants. During clinical research with Mirtazapine Orodispersible Tablets, undesirable effects have not been reported more often in elderly patients than in other age groups.

Aspartame

Mirtazapine Orodispersible Tablets contains aspartame, a source of phenylalanine. Each 15 mg orodispersible tablet contains 3mg aspartame. Each 30 mg orodispersible tablet contains 6 mg aspartame. Each 45 mg orodispersible tablet contains 9 mg aspartame. It may be harmful for patients with phenylketonuria.

4.5.Interaction with other medicinal products and other forms of interaction:

Pharmacodynamic interactions

- Mirtazapine should not be administered concomitantly with MAO inhibitors or within two weeks after discontinuation of MAO inhibitor therapy. In the opposite way about two weeks should pass before patients treated with mirtazapine should be treated with MAO inhibitors.

In addition, as with SSRIs, co-administration with other serotonergic active substances (L-tryptophan, triptans, tramadol, linezolid, methylene blue, SSRIs, venlafaxine, lithium and St. John's Wort – *Hypericum perforatum* – preparations) may lead to an incidence of serotonin associated effects. Caution should be advised and a closer clinical monitoring is required when these active substances are combined with mirtazapine.

- Mirtazapine may increase the sedating properties of benzodiazepines and other sedatives (notably most antipsychotics, antihistamine H1 antagonists, opioids). Caution should be exercised when these medicinal products are prescribed together with mirtazapine.
- Mirtazapine may increase the CNS depressant effect of alcohol. Patients should therefore be advised to avoid alcoholic beverages while taking mirtazapine. Mirtazapine dosed at 30 mg once daily causes a small but statistically significant increase in the international normalized ratio (INR) in subjects treated with warfarin. As at a higher dose of mirtazapine a more pronounced effect cannot be excluded, it is advisable to monitor the INR in case of concomitant treatment of warfarin with mirtazapine.
- The risk of QT prolongation and/or ventricular arrhythmias (e.g. Torsade de Pointes) may be increased with concomitant use of medicines which prolong the QTc interval (e.g. some antipsychotics and antibiotics).

Pharmacokinetic interactions

- Carbamazepine and phenytoin, CYP3A4 inducers, increases mirtazapine clearance about two fold, resulting in a decrease in average plasma mirtazapine concentration of 60 % and 45 %, respectively. When carbamazepine or any other inducer of hepatic metabolism (such as rifampicin) is added to mirtazapine therapy, the mirtazapine dose may have to be increased. If treatment with such medicinal product is discontinued, it may be necessary to reduce the mirtazapine dose.
- Co-administration of the potent CYP3A4 inhibitor ketoconazole increases the peak plasma levels and the AUC of mirtazapine by approximately 40% and 50% respectively.
- When cimetidine (weak inhibitor of CYP1A2, CYP2D6 and CYP3A4) is administered with mirtazapine, the mean plasma concentration of mirtazapine may increase more than 50 %. Caution should be exercised and the dose may have to be decreased when co-administering mirtazapine with potent CYP3A4 inhibitors, HIV protease inhibitors, azole antifungals, erythromycin, cimetidine or nefazodone.
- Interaction studies did not indicate any relevant pharmacokinetic effects on concurrent treatment of mirtazapine with paroxetine, amitriptyline, risperidone or lithium.

Paediatric population

Interaction studies have only been performed in adults.

4.6.Fertility, pregnancy and lactation

Pregnancy

Limited data of the use of mirtazapine in pregnant women do not indicate an increased risk for congenital malformations. Studies in animals have not shown any teratogenic effects of clinical relevance, however developmental toxicity has been observed.

Epidemiological data have suggested that the use of SSRIs in pregnancy, particularly in late pregnancy, may increase the risk of persistent pulmonary hypertension in the newborn (PPHN). Although no studies have investigated the association of PPHN to mirtazapine treatment, this potential risk cannot be ruled out taking into account the related mechanism of action (increase in serotonin concentrations).

Caution should be exercised when prescribing to pregnant women. If Mirtazapine Orodispersible Tablets is used until, or shortly before birth, postnatal monitoring of the newborn is recommended to account for possible discontinuation effects.

Breast-feeding

Animal studies and limited human data have shown excretion of mirtazapine in breast milk only in very small amounts. A decision on whether to continue/discontinue breast-

feeding or to continue/discontinue therapy with Mirtazapine Orodispersible Tablets should be made taking into account the benefit of breast-feeding to the child and the benefit of Mirtazapine Orodispersible Tablets therapy to the woman.

Fertility

Non-clinical reproductive toxicity studies in animals did not show any effect on fertility.

4.7.Effects on ability to drive and use machines

Mirtazapine Orodispersible Tablets has minor or moderate influence on the ability to drive and use machines. Mirtazapine Orodispersible Tablets may impair concentration and alertness (particularly in the initial phase of treatment). Patients should avoid the performance of potentially dangerous tasks, which require alertness and good concentration, such as driving a motor vehicle or operating machinery, at any time when affected.

4.8.Undesirable effects

Depressed patients display a number of symptoms that are associated with the illness itself. It is therefore sometimes difficult to ascertain which symptoms are result of the illness itself and which are result of treatment with Mirtazapine.

Summary of safety profile

The most commonly reported adverse reactions, occurring in more than 5 % of patients treated with Mirtazapine Orodispersible Tablets in randomised placebo-controlled trials (see below) are somnolence, sedation, dry mouth, weight increased, increase in appetite, dizziness and fatigue.

Severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), bullous dermatitis and erythema multiforme have been reported in association with Mirtazapine Orodispersible Tablets treatment.

Tabulated list of adverse reactions

All randomised placebo-controlled trials in patients (including indications other than major depressive disorder), have been evaluated for adverse reactions of Mirtazapine Orodispersible Tablets. The meta-analysis considered 20 trials, with a planned duration of treatment up to 12 weeks, with 1,501 patients (134 person years) receiving doses of mirtazapine up to 60 mg and 850 patients (79 person years) receiving placebo. Extension

phases of these trials have been excluded to maintain comparability to placebo treatment. Table 1 shows the categorised incidence of the adverse reactions, which occurred in the clinical trials statistically significantly more frequently during treatment with Mirtazapine Orodispersible Tablets than with placebo, added with adverse reactions from spontaneous reporting. The frequencies of the adverse reactions from spontaneous reporting are based on the reporting rate of these events in the clinical trials. The frequency of adverse reactions from spontaneous reporting for which no cases in the randomised placebo-controlled patient trials were observed with mirtazapine has been classified as 'not known'.

Table 1. Adverse reactions of Mirtazapine Orodispersible Tablets

System organ class	Very common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)	Frequency not known (cannot be estimated from the available data)
Blood and the lymphatic system disorders					▪ Bone marrow depression (granulocytopenia, agranulocytosis, aplastic anaemia, thrombocytopenia) ▪ Eosinophilia
Endocrine disorders					▪ Inappropriate antidiuretic hormone secretion ▪ Hyperprolactinemia (and related symptoms: galactorrhea and gynecomastia)
Metabolism and nutrition disorders	▪ Weight increased¹ ▪ Increased appetite¹				Hyponatraemia
Psychiatric disorders		▪ Abnormal dreams - Confusion - Anxiety^{2, 5} - Insomnia 3, 5	Nightmares ² Mania Agitation ² ▪ Hallucinations ▪ Psychomotor restlessness (incl. akathisia, hyperkinesia)	▪ Aggression	Suicidal ideation ⁶ ▪ Suicidal behaviour⁶ Somnambulism
Nervous system disorders	▪ Somnolence 1, 4 - Sedation 1, 4 - Headache 2	Lethargy ¹ Dizziness Tremor Amnesia ⁷	▪ Paraesthesia² ▪ Restless legs Syncope	▪ Myoclonus	▪ Convulsions (insults) ▪ Serotonin syndrome - Oral paraesthesia - Dysarthria

Vascular disorders		▪ Orthostatic hypotension	▪ Hypotension ²		
Gastrointestinal disorders	▪ Dry mouth	Nausea ³ Diarrhea ² ▪ Vomiting ² ▪ Constipation ¹	▪ Oral hypoesthesia	▪ Pancreatitis	Mouth oedema ▪ Increased salivation
Hepatobiliary disorders				▪ Elevations in serum transaminase activities	
Skin and Subcutaneous tissue disorders		Exanthema ²			▪ Stevens-Johnson Syndrome ▪ Dermatitis bullous ▪ Erythema multiforme ▪ Toxic epidermal necrolysis ▪ Drug reaction with eosinophilia and systemic symptoms (DRESS)
Musculoskeletal and connective tissue disorders		Arthralgia Myalgia ▪ Back pain ¹			Rhabdomyolysis
Renal and urinary disorders					Urinary retention
Reproductive system and breast disorders					Priapism
General disorders and administration site conditions		▪ Peripheral oedema ¹ Fatigue			▪ Generalised oedema ▪ Localised oedema
Investigations					▪ Increased creatine kinase

¹ In clinical trials these events occurred statistically significantly more frequently during treatment with Mirtazapine Orodispersible Tablets than with placebo.

² In clinical trials these events occurred more frequently during treatment with placebo than with Mirtazapine Orodispersible Tablets, however not statistically significantly more frequently.

³ In clinical trials these events occurred statistically significantly more frequently during treatment with placebo than with Mirtazapine Orodispersible Tablets.

⁴ N.B. dose reduction generally does not lead to less somnolence/sedation but can jeopardize antidepressant efficacy.

⁵ Upon treatment with antidepressants in general, anxiety and insomnia (which may be

symptoms of depression) can develop or become aggravated. Under mirtazapine treatment, development or aggravation of anxiety and insomnia has been reported.

⁶Cases of suicidal ideation and suicidal behaviours have been reported during mirtazapine therapy or early after treatment discontinuation.

⁷In most cases patients recovered after drug withdrawal.

In laboratory evaluations in clinical trials transient increases in transaminases and gamma-glutamyltransferase have been observed (however associated adverse events have not been reported statistically significantly more frequently with Mirtazapine Orodispersible Tablets than with placebo).

Paediatric population

The following adverse events were observed commonly in clinical trials in children: weight gain, urticaria and hypertriglyceridaemia.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the National Regulatory Authority pharmacovigilance reporting channels.

4.9. Overdose

Present experience concerning overdose with Mirtazapine Orodispersible Tablets alone indicates that symptoms are usually mild. Depression of the central nervous system with disorientation and prolonged sedation have been reported, together with tachycardia and mild hyper- or hypotension. However, there is a possibility of more serious outcomes (including fatalities) at dosages much higher than the therapeutic dose, especially with mixed overdoses. In these cases QT prolongation and Torsades de Pointes have also been reported.

Cases of overdose should receive appropriate symptomatic and supportive therapy for vital functions. ECG monitoring should be undertaken. Activated charcoal or gastric lavage should also be considered.

Paediatric population

The appropriate actions as described for adults should be taken in case of an overdose in paediatrics.

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

Pharmacotherapeutic group: Other Antidepressants, ATC code: N06AX11.

Mechanism of action/pharmacodynamic effects

Mirtazapine is a centrally active presynaptic α_2 -antagonist, which increases central noradrenergic and serotonergic neurotransmission. The enhancement of serotonergic neuro-transmission is specifically mediated via 5-HT₁ receptors, because 5-HT₂ and 5-HT₃ receptors are blocked by Mirtazapine. Both enantiomers of Mirtazapine are presumed to contribute to the antidepressant activity, the S(+) enantiomer by blocking α_2 and 5-HT₂ receptors and the R(-) enantiomer by blocking 5-HT₃ receptors.

Clinical efficacy and safety

The histamine H₁-antagonistic activity of mirtazapine is associated with its sedative properties. It has practically no anticholinergic activity and, at therapeutic doses, has only limited effects (e.g. orthostatic hypotension) on the cardiovascular system.

The effect of Mirtazapine Orodispersible Tablets on QTc interval was assessed in a randomised, placebo and moxifloxacin controlled clinical trial involving 54 healthy volunteers using a regular dose of 45 mg and a supra-therapeutic dose of 75mg. Linear e-max modelling suggested that prolongation of QTc intervals remained below the threshold for clinically meaningful prolongation.

Paediatric population

Two randomised, double-blind, placebo-controlled trials in children aged between 7 and 18 years with major depressive disorder (n=259) using a flexible dose for the first 4 weeks (15-45 mg mirtazapine) followed by a fixed dose (15, 30 or 45mg mirtazapine) for another 4 weeks failed to demonstrate significant differences between mirtazapine and placebo on the primary and all secondary endpoints. Significant weight gain ($\geq 7\%$) was observed in 48.8 % of the Mirtazapine Orodispersible Tablets treated subjects compared to 5.7 % in the placebo arm. Urticaria (11.8 % vs. 6.8 %) and hypertriglyceridaemia (2.9 % vs. 0 %) were also commonly observed.

5.2. Pharmacokinetic properties

Absorption

After oral administration Mirtazapine Orodispersible Tablets, the active substance mirtazapine is rapidly and well absorbed (bioavailability $\approx 50\%$), reaching peak plasma levels after approx. two hours. Food intake has no influence on the pharmacokinetics of mirtazapine.

Distribution

Binding of mirtazapine to plasma proteins is approx. 85 %.

Biotransformation

Major pathways of biotransformation are demethylation and oxidation, followed by conjugation. In vitro data from human liver microsomes indicate that cytochrome P450 enzymes CYP2D6 and CYP1A2 are involved in the formation of the 8-hydroxy metabolite of mirtazapine, whereas CYP3A4 is considered to be responsible for the formation of the N-demethyl and N-oxide metabolites. The demethyl metabolite is pharmacologically active and appears to have the same pharmacokinetic profile as the parent compound.

Elimination

Mirtazapine is extensively metabolised and eliminated via the urine and faeces within a few days. The mean half-life of elimination is 20-40 hours; longer half-lives, up to 65 hours, have occasionally been recorded and shorter half-lives have been seen in young men. The half-life of elimination is sufficient to justify once-a-day dosing. Steady state is reached after 3-4 days, after which there is no further accumulation.

Linearity/non-linearity

Mirtazapine displays linear pharmacokinetics within the recommended dose range.

Special populations

The clearance of mirtazapine may be decreased as a result of renal or hepatic impairment.

5.3.Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential, toxicity to reproduction and development. In reproductive toxicity studies in rats and rabbits no teratogenic effects were observed. At two-fold systemic exposure compared to maximum human therapeutic exposure, there was an increase in post-implantation loss, decrease in the pup birth weights, and reduction in pup survival during the first three days of lactation in rats.

Mirtazapine was not genotoxic in a series of tests for gene mutation and chromosomal and DNA damage. Thyroid gland tumours found in a rat carcinogenicity study and hepatocellular neoplasms found in a mouse carcinogenicity study are considered to be species-specific, non-genotoxic responses associated with long-term treatment with

high doses of hepatic enzyme inducers.

6. PHARMACEUTICAL PARTICULARS

6.1.List of excipients

Crospovidone, Mannitol, Cellulose Microcrystalline, Aspartame, Strawberry guarana flavor, Peppermint flavor, Silica Colloidal Anhydrous and Magnesium Stearate.

6.2.Incompatibilities

None known.

6.3.Shelf life

36 Months.

6.4.Special precautions for storage

Do not store above 30°C.

6.5.Nature and contents of container

Blister Pack: AUROZAPINE OD are packaged in 6's Peelable PVC/ALU-ALU blister pack.

6.6.Special precautions for disposal and other handling:

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORIZATION HOLDER

Marketing Authorisation Holder:

Aurobindo Pharma Ltd.,

Plot No: 2, Maitrivihar,

Ameerpet, Hyderabad – 500 038, Telangana State, India.

Manufactured by:

Aurobindo Pharma Limited, Unit- III,

Survey No. 313 & 314,

Bachupally, Bachupally Mandal,

Medchal-Malkajgiri District,

Telangana State, India.

8. MARKETING AUTHORIZATION NUMBER (S)

AUROZAPINE OD 45: H2026/CTD11559/15091

AUROZAPINE OD 30: H2025/CTD12167/22596

AUROZAPINE OD 15:

9. DATE OF FIRST AUTHORIZATION/RENEWAL OF THE AUTHORIZATION

AUROZAPINE OD 45:

AUROZAPINE OD 30:

AUROZAPINE OD 15:

10. DATE OF REVISION OF THE TEXT

May 2022.