

SUMMARY OF PRODUCT CHARACTERISTICS

1. Name of the medicinal product

Apisopt eye drops.

2. Qualitative and quantitative composition

Active Ingredients:

Each ml contains 22.26 mg of Dorzolamide hydrochloride equivalent to 20 mg of Dorzolamide.

Excipient with known effect: Benzalkonium chloride

For the full list of excipients, see section 6.1.

3. Pharmaceutical form

Eye drops, solution.

4. Clinical particulars

4.1 Therapeutic indications

Apisopt is indicated:

- As adjunctive therapy to beta-blockers,
- As monotherapy in patients unresponsive to beta-blockers or in whom beta-blockers are contraindicated,

In the treatment of elevated intra-ocular pressure in:

- Ocular hypertension,
- Open-angle glaucoma,
- Pseudoexfoliative glaucoma.

4.2 Posology and method of administration

Posology

When used as monotherapy, the dose is one drop of Dorzolamide in the conjunctival sac of the affected eye(s), three times daily. When used as adjunctive therapy with an ophthalmic beta-blocker, the dose is one drop of Dorzolamide in the conjunctival sac of the affected eye(s), two times daily.

When substituting Dorzolamide for another ophthalmic anti-glaucoma agent, discontinue the other agent after proper dosing on one day, and start Dorzolamide on the next day. If more than one topical ophthalmic drug is being used, the drugs should be administered at least ten minutes apart. Patients

should be instructed to wash their hands before use and avoid allowing the tip of the container to come into contact with the eye or surrounding structures. Patients should also be instructed that ocular solutions, if handled improperly, can become contaminated by common bacteria known to cause ocular infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated solutions.

Paediatric population

Limited clinical data in paediatric patients with administration of Dorzolamide three times a day are available. (For information regarding paediatric dosing see section 5.1).

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Dorzolamide has not been studied in patients with severe renal impairment ($\text{CrCl} < 30$ ml/min) or with hyperchloraemic acidosis. Because Dorzolamide and its metabolites are excreted predominantly by the kidney, Dorzolamide is therefore contra-indicated in such patients.

4.4 Special warnings and precautions for use

Dorzolamide has not been studied in patients with hepatic impairment and should therefore be used with caution in such patients.

The management of patients with acute angle-closure glaucoma requires therapeutic interventions in addition to ocular hypotensive agents. Dorzolamide has not been studied in patients with acute angle-closure glaucoma.

Dorzolamide contains a sulphonamido group, which also occurs in sulphonamides and although administered topically, is absorbed systemically. Therefore the same types of adverse reactions that are attributable to sulphonamides may occur with topical administration, including severe reactions such as Stevens-Johnson syndrome and toxic epidermal necrolysis. If signs of serious reactions or hypersensitivity occur, discontinue the use of this preparation.

Therapy with oral carbonic anhydrase inhibitors has been associated with urolithiasis as a result of acid-base disturbances, especially in patients with a prior history of renal calculi.

Although no acid-base disturbances have been observed with Dorzolamide, urolithiasis has been reported infrequently. Because Dorzolamide is a topical carbonic anhydrase inhibitor that is absorbed systemically, patients with a prior history of renal calculi may be at increased risk of urolithiasis while using Dorzolamide.

If allergic reactions (e.g., conjunctivitis and eye-lid reactions) are observed, discontinuation of treatment should be considered.

There is a potential for an additive effect on the known systemic effects of carbonic anhydrase inhibition in patients receiving an oral carbonic anhydrase inhibitor and Dorzolamide. The concomitant administration of Dorzolamide and oral carbonic anhydrase inhibitors is not recommended.

Corneal oedemas and irreversible corneal decompensations have been reported in patients with pre-existing chronic corneal defects and/or a history of intra-ocular surgery while using

Apisopt. Topical Dorzolamide should be used with caution in such patients.

Choroidal detachment concomitant with ocular hypotony have been reported after filtration procedures with administration of aqueous suppressant therapies.

Apisopt contains the preservative benzalkonium chloride, which may cause eye irritation.

Contact lenses should be removed prior to application and wait at least 15 minutes before reinsertion. Benzalkonium chloride is known to discolor soft contact lenses.

Paediatric population

Dorzolamide has not been studied in patients less than 36 weeks gestational age and less than

1 week of age. Patients with significant renal tubular immaturity should only receive

Dorzolamide after careful consideration of the risk benefit balance because of the possible risk of metabolic acidosis.

4.5 Interaction with other medicinal products and other forms of interaction

Specific drug interaction studies have not been performed with Dorzolamide. Dorzolamide was used concomitantly with the following medications without evidence of adverse interactions: timolol ophthalmic solution, betaxolol ophthalmic solution and systemic medications, including ACE-inhibitors, calcium-channel blockers, diuretics, non-steroidal anti-inflammatory drugs including aspirin, and hormones (e.g. oestrogen, insulin, and thyroxine). Association between Dorzolamide and miotics and adrenergic agonists has not been fully evaluated during glaucoma therapy.

4.6 Fertility, Pregnancy and lactation

Pregnancy

Pregnancy

Dorzolamide should not be used during pregnancy. No adequate clinical data in exposed pregnancies are available. In rabbits, Dorzolamide produced teratogenic effects at maternotoxic doses (see Section 5.3)

Breast-feeding

It is not known whether Dorzolamide is excreted in human milk. In lactating rats, decreases in the body weight gain of offspring were observed. If treatment with Dorzolamide is required, then lactation is not recommended.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

Possible side effects such as dizziness and visual disturbances may affect the ability to drive and use machines.

4.8 Undesirable effects

In long-term studies as monotherapy or as adjunctive therapy with an ophthalmic betablocker, the most frequent cause of discontinuation (approximately 3%) from treatment with Apisopt was drug-related ocular adverse reactions, primarily conjunctivitis and lid reactions.

The following adverse reactions have been reported:

[Very Common: ($\geq 1/10$), Common: ($\geq 1/100$ to $<1/10$), Uncommon: ($\geq 1/1,000$ to $<1/100$),

Rare: ($\geq 1/10,000$ to $<1/1,000$)]

Nervous system disorders:

Common: headache

Rare: dizziness, paraesthesia

Eye disorders:

Very common: burning and stinging

Common: superficial punctate keratitis, tearing, conjunctivitis, eyelid

inflammation, eye itching, eyelid irritation, blurred vision *Uncommon:*

iridocyclitis *Rare:* irritation including redness, pain, eyelid crusting, transient

myopia (which resolved upon discontinuation of therapy), corneal oedema,

ocular hypotony, choroidal detachment following filtration surgery

Respiratory, thoracic, and mediastinal disorders:

Rare: epistaxis

Gastrointestinal disorders:

Common: nausea, bitter taste

Rare: throat irritation, dry mouth

Skin and subcutaneous tissue disorders:

Rare: contact dermatitis, Stevens-Johnson syndrome, toxic epidermal necrolysis

Renal and urinary disorders:

Rare: urolithiasis

General disorders and administration site conditions:

Common: asthenia/fatigue

Rare: Hypersensitivity: signs and symptoms of local reactions (palpebral reactions) and systemic allergic reactions, including angioedema, urticaria and pruritus, rash, shortness of breath, rarely bronchospasm

Investigations:

Dorzolamide was not associated with clinically meaningful electrolyte disturbances.

Paediatric population

See section 5.1.

Reporting suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after Authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org/>

4.9 Overdose

Only limited information is available with regard to human overdose by accidental or deliberate ingestion of Dorzolamide hydrochloride.

Symptoms

The following have been reported with oral ingestion: somnolence; topical application: nausea, dizziness, headache, fatigue, abnormal dreams, and dysphagia.

Treatment

Treatment should be symptomatic and supportive. Electrolyte imbalance, development of an acidotic state, and possible central nervous system effects may occur. Serum electrolyte levels (particularly potassium) and blood pH levels should be monitored.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiglaucoma preparations and miotics, Carbonic Anhydrase

Inhibitors, Dorzolamide, ATC code: S01EC03

Mechanism of Action

Carbonic anhydrase (CA) is an enzyme found in many tissues of the body including the eye.

In humans, carbonic anhydrase exists as a number of isoenzymes, the most active being carbonic anhydrase II (CA-II) found primarily in red blood cells (RBCs) but also in other tissues. Inhibition of carbonic anhydrase in the ciliary processes of the eye decreases aqueous

humour secretion. The result is a reduction in intra-ocular pressure (IOP).

Apisopt contains Dorzolamide hydrochloride, a potent inhibitor of human carbonic anhydrase

II. Following topical ocular administration, Dorzolamide reduces elevated intra-ocular pressure, whether or not associated with glaucoma. Elevated intra-ocular pressure is a major risk factor in the pathogenesis of optic nerve damage and visual-field loss. Dorzolamide does not cause pupillary constriction and reduces intra-ocular pressure without side effects such as night blindness, accommodative spasm. Dorzolamide has minimal or no effect on pulse rate or blood pressure.

Topically applied beta-adrenergic blocking agents also reduce IOP by decreasing aqueous

humour secretion but by a different mechanism of action. Studies have shown that when

Dorzolamide is added to a topical beta-blocker, additional reduction in IOP is observed; this finding is consistent with the reported additive effects of beta-blockers and oral carbonic anhydrase inhibitors.

Pharmacodynamic effects

Clinical effects:

Adult patients

In patients with glaucoma or ocular hypertension, the efficacy of Dorzolamide given t.i.d. as monotherapy (baseline IOP ≥ 23 mmHg) or given b.i.d. as adjunctive therapy while receiving ophthalmic beta-blockers (baseline IOP ≥ 22 mmHg) was demonstrated in large-scale clinical studies of up to one-year duration. The IOP-lowering effect of Dorzolamide as monotherapy and as adjunctive therapy was demonstrated throughout the day and this effect was maintained during long-term administration. Efficacy during long-term monotherapy was similar to betaxolol and slightly less than timolol. When used as adjunctive therapy to ophthalmic beta-blockers, Dorzolamide demonstrated additional IOP lowering similar to pilocarpine 2% q.i.d.

Paediatric population

A 3-month, double-masked, active-treatment controlled, multicentre study was undertaken in

184 (122 for Dorzolamide) paediatric patients from 1 week of age to <6 years of age with glaucoma or elevated intraocular pressure (baseline IOP ≥ 22 mmHg) to assess the safety of

Apisopt when administered topically t.i.d. (three times a day). Approximately half the patients in both treatment groups were diagnosed with congenital glaucoma; other common etiologies were Sturge Weber syndrome, iridocorneal mesenchymal dysgenesis, aphakic patients. The

distribution by age and treatments Timolol

in the monotherapy phase was as

follows: Dorzolamide 2%

Age cohort < 2 years	N=56	Timolol GS 0.25%
	Age range: 1 to 23 months	N=27
		Age range: 0.25 to 22 months
Age cohort ≥ 2 - < 6 years	N=66	Timolol 0.50% N=35
	Age range: 2 to 6 years	Age range: 2 to 6 years

Overall, this study did not reveal additional safety concerns in paediatric patients: approximately 26% (20% in Dorzolamide monotherapy) of paediatric

patients were observed to experience drug related adverse effects, the majority of which were local, non-serious ocular effects such as ocular burning and stinging, injection and eye pain. A small percentage < 4% was observed to have corneal oedema or haze. Local reactions appeared similar in frequency to comparator. In post marketing data, metabolic acidosis in the very young particularly with renal immaturity/impairment has been reported.

Efficacy results in paediatric patients suggest that the mean IOP decrease observed in the

Dorzolamide group was comparable to the mean IOP decrease observed in the timolol group even if a slight numeric advantage was observed for timolol.

Longer-term efficacy studies (>12 weeks) are not available.

5.2 Pharmacokinetic properties

Unlike oral carbonic anhydrase inhibitors, topical administration of Dorzolamide hydrochloride allows for the active substance to exert its effects directly in the eye at substantially lower doses and therefore with less systemic exposure. This resulted in a reduction in IOP without the acid-base disturbances or alterations in electrolytes characteristic of oral carbonic anhydrase inhibitors.

When topically applied, Dorzolamide reaches the systemic circulation. To assess the potential for systemic carbonic anhydrase inhibition following topical administration, active substance and metabolite concentrations in red blood cells (RBCs) and plasma and carbonic anhydrase inhibition in RBCs were measured. Dorzolamide accumulates in RBCs during chronic dosing as a result of selective binding to CA-II while extremely low concentrations of free active substance in plasma are maintained. The parent active substance forms a single N-desethyl metabolite that inhibits CA-II less potently than the parent active substance but also inhibits a less active isoenzyme (CA-I). The metabolite also accumulates in RBCs where it binds primarily to CA-I. Dorzolamide binds moderately to plasma proteins (approximately 33%).

Dorzolamide is primarily excreted unchanged in the urine; the metabolite is also excreted in urine. After dosing ends, Dorzolamide washes out of RBCs non linearly, resulting in a rapid decline of active substance concentration initially, followed by a slower elimination phase with a half-life of about four months. When Dorzolamide was given orally to simulate the maximum systemic exposure after long term topical ocular administration, steady state was reached within 13 weeks. At steady state, there was virtually no free active substance or metabolite in plasma; CA inhibition in RBCs was less than that anticipated to be necessary for a pharmacological effect on renal function or respiration. Similar pharmacokinetic results were observed after chronic, topical administration of Dorzolamide. However, some elderly patients with renal impairment (estimated CrCl 30-60 ml/min) had higher metabolite concentrations in RBCs, but no meaningful differences in carbonic anhydrase inhibition, and no clinically significant systemic side effects were directly attributable to this finding.

5.3 Preclinical safety data

Not applicable.

6. Pharmaceutical particulars

6.1 List of excipients

Mannitol

Sodium citrate

Hydroxypropyl methylcellulose (Hypermellose 2910)

Sodium hydroxide

Benzalkonium chloride

Highly purified water

6.2 Incompatibilities

Not Applicable.

6.3 Shelf life

2 years.

After first opening the container, Apisopt should be used no longer than 1 month.

6.4 Special precautions for storage

Don't store above 30° C.

6.5 Nature and contents of container

LDPE bottle and HDPE cap.

6.6 Special precautions for disposal and other handling

No special requirements.

7. Marketing authorization holder

Amman Pharmaceutical Industries (API).

Jordan / Amman, Sahab,

Second King Abdullah II Industrial City.

8. Marketing authorization number(s)

H2025/CTD6890/13288

9. Date of first authorization/renewal of the authorization

17/02/2025

10. Date of revision of the text

22/08/2026