

SUMMARY OF PRODUCT CHARECTERISTICS (SPC)

1.NAME OF THE MEDICINAL PRODUCT

Bectoxime 400 Tablet

2. Qualitative and Quantitative Composition

Each film coated tablet contains:

Cefixime Trihydrate USP

Equivalent to Cefixime (Anhydrous).....400 mg

Excipient of know effect:

Lactose

For full list of excipients, see section 6.1

3. Pharmaceutical Form

Tablet, Oral dosage form

White coloured oblong shaped, film coated tablet break line on one side and plain on another

4.Clinical particulars

4.1 Therapeutic indications

Bectoxime 400 is an orally active cephalosporin antibiotic which has marked in vitro bactericidal activity against a wide variety of Gram-positive and Gram-negative organisms.

It is indicated for the treatment of the following acute infections when caused by susceptible micro-organisms:

Upper Respiratory Tract Infections (URTI): e.g. otitis media; and other URTI where the causative organism is known or suspected to be resistant to other commonly used antibiotics, or where treatment failure may carry significant risk.

Lower Respiratory Tract Infection: e.g. bronchitis.

Urinary Tract Infections: e.g. cystitis, cystourethritis, uncomplicated pyelonephritis.

Clinical efficacy has been demonstrated in infections caused by commonly occurring pathogens including *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Escherichia coli*, *Proteus mirabilis*, *Kliebsiella* species, *Haemophilus influenzae* (beta-lactamase positive and negative), *Branhamella catarrhalis* (beta-lactamase positive and negative) and *Enterobacter* species. Bectoxime 400 is highly stable in the presence of beta-lactamase enzymes.

Most strains of enterococci (*Streptococcus faecalis*, group D *Streptococci*) and *Staphylococci* (including coagulase positive and negative strains and meticillin-resistant strains) are resistant to Bectoxime 400. In addition, most strains of *Pseudomonas*, *Bacteriodes fragalis*, *Listeria monocytogenes* and *Clostridia* are resistant to Bectoxime 400.

4.2 Posology and method of administration

The usual course of treatment is 7 days. This may be continued for up to 14 days if required. Posology

Adults and Children over 10 Years or weighing more than 50 kg:

The recommended adult dosage is 400 mg daily according to the severity of infection, given either as a single dose or in two divided doses.

Elderly:

Elderly patients may be given the same dose as recommended for adults. Renal function should be assessed and dosage should be adjusted in severe renal impairment (See "Dosage in Renal Impairment").

Children under 10 Years:

Bectoxime 400 are not recommended for use in children under 10 years old. The safety and efficacy of cefixime has not been established in children less than 6 months.

Renal Impairment:

Bectoxime 400 may be administered in the presence of impaired renal function. Normal dose and schedule may be given in patients with creatinine clearances of 20 ml/min or greater. In patients whose creatinine clearance is less than 20 ml/min, it is recommended that a dose of once daily should not be exceeded. The dose and regimen for patients who are maintained on chronic ambulatory peritoneal dialysis or haemodialysis should follow the same recommendation as that for patients with creatinine clearances of less than 20 ml/min.

Method for administration

For oral administration.

Absorption of Bectoxime 400 is not significantly modified by the presence of food.

4.3 Contraindications

Hypersensitivity to cephalosporin antibiotics or to any of the excipients

4.4 Special warnings and precautions for use

Encephalopathy

Beta-lactams, including cefixime, predispose the patient to encephalopathy risk (which may include convulsions, confusion, impairment of consciousness, movement disorders), particularly in case of overdose or renal impairment.

Severe cutaneous adverse reactions

Severe cutaneous adverse reactions such as toxic epidermal necrolysis, Stevens- Johnson syndrome and drug rash with eosinophilia and systemic symptoms (DRESS) have been reported in some patients on cefixime. When severe cutaneous adverse reactions occur, cefixime should be discontinued and appropriate therapy and/or measures should be taken.

Bectoxime 400 should be given with caution to patients who have shown hypersensitivity to other drugs.

Hypersensitivity to penicillins

As with other cephalosporins, cefixime should be given with caution to patients

with a history of hypersensitivity to penicillin, as there is some evidence of partial cross-allergenicity between the penicillins and cephalosporins.

Patients have had severe reactions (including anaphylaxis) to both classes of drugs. If an allergic effect occurs with Bectoxime 400, the drug should be discontinued and the patient treated with appropriate agents if necessary.

Haemolytic anaemia

Drug-induced haemolytic anaemia, including severe cases with a fatal outcome, has been described for cephalosporins (as a class). The recurrence of haemolytic anaemia after re-administration of cephalosporins in a patient with a history of cephalosporin (including cefixime) –associated haemolytic anaemia has also been reported.

Acute renal failure

As with other cephalosporins, cefixime may cause acute renal failure including tubulointerstitial nephritis as an underlying pathological condition. When acute renal failure occurs, cefixime should be discontinued and appropriate therapy and/or measures should be taken.

Renal impairment

Bectoxime 400 should be administered with caution in patients with markedly impaired renal function

Paediatric use

Safety of cefixime in premature or newborn infant has not been established.

Antibiotic-associated colitis

Treatment with broad spectrum antibiotics alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is a primary cause of antibiotic-associated diarrhoea.

Pseudomembranous colitis is associated with the use of broad-spectrum antibiotics (including macrolides, semi-synthetic penicillins, lincosamides and cephalosporins); it is therefore important to consider its diagnosis in patients who develop diarrhoea in association with the use of antibiotics. Symptoms of pseudomembranous colitis may occur during or after antibiotic treatment.

Management of pseudomembranous colitis should include sigmoidoscopy, appropriate bacteriologic studies, fluids, electrolytes and protein supplementation. If the colitis does not improve after the drug has been discontinued, or if the symptoms are severe, oral vancomycin is the drug of choice for antibiotic-associated pseudomembranous colitis produced by *C. difficile*. Other causes of colitis should be excluded.

4.5 Interaction with other medicinal products and other forms of interaction

Anticoagulants

In common with other cephalosporins, increases in prothrombin times have been noted in a few patients. Care should therefore be taken in patients receiving anticoagulation therapy.

Cefixime should be administered with caution to patients receiving coumarin-type anticoagulants, e.g. warfarin potassium. Since cefixime may enhance effects of the anticoagulants, prolonged prothrombin time with or without

bleeding may occur.

Other forms of interaction

A false positive reaction for glucose in the urine may occur with Benedict's or Fehling's solutions or with copper sulphate test tablets, but not with tests based on enzymatic glucose oxidase reactions.

A false positive direct Coombs test has been reported during treatment with cephalosporin antibiotics; therefore it should be recognised that a positive Coombs test may be due to the drug.

4.6 Fertility, pregnancy and lactation

Reproduction studies have been performed in mice and rats at doses up to 400 times the human dose and have revealed no evidence of impaired fertility or harm to the foetus due to cefixime. In the rabbit, at doses up to 4 times the human dose, there was no evidence of a teratogenic effect; there was a high incidence of abortion and maternal death which is an expected consequence of the known sensitivity of rabbits to antibiotic-induced changes in the population of the microflora of the intestine. There are no adequate and well-controlled studies in pregnant women. Bectoxime 400 should therefore not be used in pregnancy or in nursing mothers unless considered essential by the physician.

4.7 Effects on ability to drive and use machines

In the case of side effects such as encephalopathy (which may include convulsion, confusion, impairment of consciousness, movement disorders), the patient should not operate machines or drive a vehicle.

4.8 Undesirable effects

Bectoxime 400 is generally well tolerated. The majority of adverse reactions observed in clinical trials were mild and self-limiting in nature.

The following adverse reaction (Preferred term# or equivalent) will be considered listed:

Blood and lymphatic system disorders:	Eosinophilia Hypereosinophilia Agranulocytosis Leucopenia Neutropenia Granulocytopenia Haemolytic anaemia Thrombocytopenia Thrombocytosis
Gastrointestinal disorders:	Abdominal pain Diarrhoea* Dyspepsia Nausea Vomiting Flatulence
Hepatobiliary disorders:	Jaundice
Infections and infestations:	Pseudomembranous colitis

Investigations:	Aspartate aminotransferase increased Alanine aminotransferase increased Blood bilirubin increased Blood urea increased Blood creatinine increased
Nervous system disorders:	Dizziness Headache Cases of convulsions have been reported with cephalosporins including cefixime (frequency not known) ** Beta-lactams, including cefixime, predispose the patient to encephalopathy risk (which may include convulsions, confusion, impairment of consciousness, movement disorders), particularly in case of overdose or renal impairment (frequency not known) **
Respiratory, thoracic and mediastinal disorders:	Dyspnoea
Renal and urinary disorders:	Renal failure acute including tubulointerstitial nephritis as an underlying pathological condition
Immune system disorders, administrative site conditions, skin and subcutaneous tissue disorders:	Anaphylactic reaction Serum sickness-like reaction Drug rash with eosinophilia and systemic symptoms (DRESS) Pruritus Rash Drug Fever Arthralgia Erythema multiforme Stevens-Johnson syndrome Toxic epidermal necrolysis Angio-oedema Urticaria Pyrexia Face oedema Genital pruritus Vaginitis

The above mentioned listed adverse reactions have been observed during clinical studies and/or during marketed use.

*Diarrhoea has been more commonly associated with higher doses. Some cases of moderate to severe diarrhoea have been reported; this has occasionally warranted cessation of therapy. Bectoxime 400 should be discontinued if marked diarrhoea occurs

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are requested to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org/>

4.9 Overdose

There is a risk of encephalopathy in cases of administration of beta-lactam antibiotics, including cefixime, particularly in case of overdose or renal impairment.

Adverse reactions seen at dose levels up to 2 g Bectoxime 400 in normal subjects did not differ from the profile seen in patients treated at the recommended doses.

Cefixime is not removed from the circulation in significant quantities by dialysis. No specific antidote exists. General supportive measures are recommended.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: third generation

cephalosporin ATC code: J01DD08

Cefixime is an oral third generation cephalosporin which has marked in vitro bactericidal activity against a wide variety of Gram-positive and Gram-negative organisms.

Clinical efficacy has been demonstrated in infections caused by commonly occurring pathogens including *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Escherichia coli*, *Proteus mirabilis*, *Klebsiella* species, *Haemophilus influenzae* (beta-lactamase positive and negative), *Branhamella catarrhalis* (beta-lactamase positive and negative) and *Enterobacter* species. It is highly stable in the presence of beta-lactamase enzymes.

Most strains of enterococci (*Streptococcus faecalis*, group D Streptococci) and Staphylococci (including coagulase positive and negative strains and meticillin-resistant strains) are resistant to cefixime. In addition, most strains of *Pseudomonas*, *Bacteroides fragilis*, *Listeria monocytogenes* and Clostridia are resistant to cefixime.

5.2 Pharmacokinetic properties

The absolute oral bioavailability of cefixime is in the range of 22-54%. Absorption is not significantly modified by the presence of food. Cefixime may therefore be given without regard to meals.

From in vitro studies, serum or urine concentrations of 1 mcg/mL or greater were considered to be adequate for most common pathogens against which cefixime is active. Typically, the peak serum levels following the recommended adult or paediatric doses are between 1.5 and 3 mcg/mL. Little or no accumulation of cefixime occurs following multiple dosing.

The pharmacokinetics of cefixime in healthy elderly (age > 64 years) and young volunteers (11-35) compared the administration of 400 mg doses once daily for 5 days. Mean C_{max} and AUC values were slightly greater in the elderly. Elderly patients may be given the same dose as the general population.

Cefixime is predominantly eliminated as unchanged drug in the urine. Glomerular filtration is considered the predominant mechanism. Metabolites of cefixime have not been isolated from human serum or urine.

Serum protein binding is well characterised for human and animal sera;

cefixime is almost exclusively bound to the albumin fraction, the mean free fraction being approximately 30%. Protein binding of cefixime is only concentration dependent in human serum at very high concentrations which are not seen following clinical dosing.

Transfer of ¹⁴C-labelled cefixime from lactating rats to their nursing offspring through breast milk was quantitatively small (approximately 1.5% of the mothers' body content of cefixime in the pup). No data are available on secretion of cefixime in human breast milk. Placental transfer of cefixime was small in pregnant rats dosed with labelled cefixime.

5.3 Preclinical safety data

There are no pre-clinical data of relevance to the prescriber which are additional to that already included in other sections of the Summary of Product Characteristics.

6. Pharmaceutical Particulars

6.1. List of Excipients

Maize Starch

Lactose

Microcrystalline cellulose

Sodium benzoate

Purified Talc

Magnesium Stearate

Cross Carmellose Sodium

Colloidal Silicon Dioxide

Isopropyl alcohol

Methylene Dichloride

Titanium Dioxide BP.

6.2. Incompatibilities

None

6.3. Shelf life

36Months.

6.4 Special precautions for storage

Store below above 30 °C.

Protect from light & moisture.

6.5. Nature and contents of container

Aluminum - Aluminum Blister Pack

10 Tablet are blister packed with Aluminum - Aluminum foil; such 1 blister is packed in one carton pack.

Pack size: 1 x 10 Tablets (i.e. 10 Tablets) in one carton box along with packing leaflet.

6.6. Special precautions for disposal and other handling

There are no special storage precautions. Any unused product or waste material

should be disposed of in accordance with local requirements.

7.0. MARKETING AUTHORIZATION HOLDER AND MANUFACTURING SITE ADDRESSES

MAH:

Eastleigh Pharmaceuticals Co. Ltd
P.O Box 167-00610 Nairobi, Kenya

Manufacturer

MARS REMEDIES PVT LTD

Address: 635, GIDC Estate, Waghodia-391760, Vadodara, Gujarat India

8.0. MARKETING AUTHORIZATION NUMBER

H2022/CTD9056/20433

9.0. DATE OF FIRST REGISTRATION /RENEWAL OF REGISTRATION

2022 September 14

10.0 DATE OF REVISION OF THE TEXT

Last revised on 10-July-2020