

SUMMARY OF PRODUCT CHARACTERISTICS

Caberlin 0.5 (Cabergoline 0.5 mg Tablet)

1. NAME OF THE FINISHED PHARMACEUTICAL PRODUCT

Caberlin 0.5

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Caberlin 0.5

Each film-coated tablet contains:

Cabergoline..... 0.5 mg.

For excipients, see **section 6.1**

3. PHARMACEUTICAL FORM

4. CLINICAL PARTICULARS

4.1 *Therapeutic indications*

Inhibition/suppression of physiological lactation

Cabergoline is indicated for the inhibition of physiological lactation soon after delivery and for suppression of already established lactation:

- After parturition, when the mother elects not to breast feed the infant or when breast feeding is contraindicated due to medical reasons related to the mother or the new-born.
- After stillbirth or abortion.

Cabergoline prevents/suppresses physiological lactation by inhibiting prolactin secretion.

In the reported controlled clinical studies, cabergoline given as a single 1 mg administration during the first day post-partum, was effective in inhibiting milk secretion, as well as breast engorgement and pain in 70 - 90% of the women. Less than 5% of women experienced rebound breast symptomatology during the third post-partum week (which was usually mild in severity).

Suppression of milk secretion and relief of breast engorgement and pain are obtained in approximately 85% of nursing women treated with a total dose of 1 mg cabergoline given

in four divided doses over two days. Rebound breast symptomatology after day 10 is uncommon (approximately 2% of cases).

Treatment of hyperprolactinaemic disorders

Cabergoline is indicated for the treatment of dysfunctions associated with hyperprolactinemia, including amenorrhoea, oligomenorrhoea, anovulation and galactorrhoea. Cabergoline is indicated in patients with prolactin-secreting pituitary adenomas (micro- and macroprolactinomas), idiopathic hyperprolactinemia, or empty sella syndrome with associated hyperprolactinemia, which represent the basic underlying pathologies contributing to the above clinical manifestations.

On chronic therapy, cabergoline at doses ranging between 1 and 2 mg per week, was effective in normalising serum prolactin levels in approximately 84% of hyperprolactinaemic patients. Regular cycles were resumed in 83% of previously amenorrhoeic women. Restoration of ovulation was documented in 89% of women with progesterone levels monitored during the luteal phase. Galactorrhoea disappeared in 90% of cases showing this symptom before therapy. Reduction in tumour size was obtained in 50 - 90% of female and male patients with micro- or macroprolactinoma.

4.2 Posology and method of administration

Posology

Inhibition/suppression of physiological lactation

For inhibition of lactation cabergoline should be administered during the first day post-partum. The recommended therapeutic dose is 1 mg (two 0.5 mg tablets) given as a single dose.

For suppression of established lactation, the recommended therapeutic dosage regimen is 0.25 mg (one-half 0.5 mg tablet) every 12 hours for two days (1 mg total dose). This dosage regimen has been demonstrated to be better tolerated than the single dose regimen in women electing to suppress lactation having a lower incidence of adverse events, in particular of hypotensive symptoms.

Treatment of hyperprolactinemic disorders

The recommended initial dosage of cabergoline is 0.5 mg per week given in one or two (one-half of one 0.5 mg tablet) doses (e.g. on Monday and Thursday) per week. The weekly dose should be increased gradually, preferably by adding 0.5 mg per week at monthly intervals until an optimal therapeutic response is achieved. The therapeutic dosage is usually 1 mg per week and ranges from 0.25 mg to 2 mg per week. Doses of cabergoline

up to 4.5 mg per week have been used in hyperprolactinaemic patients.

The maximum dose should not exceed 3 mg per day.

The weekly dose may be given as a single administration or divided into two or more doses per week according to patient tolerability. Division of the weekly dose into multiple administrations is advised when doses higher than 1 mg per week are to be given since the tolerability of doses greater than 1 mg taken as a single weekly dose has been evaluated only in a few patients.

Patients should be evaluated during dose escalation to determine the lowest dosage that produces the therapeutic response. Monitoring of serum prolactin levels at monthly intervals is advised since, once the effective therapeutic dosage regimen has been reached, serum prolactin normalisation is usually observed within two to four weeks.

After cabergoline withdrawal, recurrence of hyperprolactinaemia is usually observed. However, persistent suppression of prolactin levels has been observed for several months in some patients. Of the group of women followed up, majority of women had ovulatory cycles which continued for greater than 6 months after cabergoline discontinuation.

Paediatrics

The safety and efficacy of cabergoline has not been established in subjects less than 16 years of age.

Elderly

As a consequence of the indications for which cabergoline is presently proposed, the experience in elderly is very limited. Available data do not indicate a special risk.

Method of administration

Cabergoline is to be administered by the oral route. Since in the reported clinical studies cabergoline has been mainly administered with food and since the tolerability of this class of compounds is improved with food, it is recommended that cabergoline be preferably taken with meals for all the therapeutic indications.

4.3 Contraindications

- Hypersensitivity to cabergoline, any of the excipients or any ergot alkaloid.
- History of pulmonary, pericardial and retroperitoneal fibrotic disorders.

- Cabergoline is contraindicated in patients with hepatic insufficiency and with toxemia of pregnancy. Cabergoline should not be co-administered with anti-psychotic medications or administered to women with a history of puerperal psychosis.
- For long-term treatment: evidence of cardiac valvulopathy as determined by pre-treatment echocardiography (See **section 4.4**).

4.4 *Special warnings and precautions for use*

General

The safety and efficacy of cabergoline have not yet been established in patients with renal and hepatic disease. As with other ergot derivatives, cabergoline should be given with caution to patients with severe cardiovascular disease, Raynaud's syndrome, renal insufficiency, peptic ulcer or gastrointestinal bleeding, or with a history of serious, particularly psychotic, mental disorders. Particular care should be taken when patients are taking concomitant psychoactive medication.

Symptomatic hypotension can occur with cabergoline administration for any indication. Care should be exercised when administering cabergoline concomitantly with other drugs known to lower blood pressure.

The effects of alcohol on overall tolerability of cabergoline are currently unknown. Before cabergoline administration, pregnancy should be excluded and after treatment pregnancy should be prevented for at least one month.

Hepatic Insufficiency

Lower doses should be considered in patients with severe hepatic insufficiency who receive prolonged treatment with cabergoline. Compared to normal volunteers and those with lesser degrees of hepatic insufficiency, an increase in AUC has been reported in patients with severe hepatic insufficiency (Child-Pugh Class C) who received a single 1 mg dose.

Postural Hypotension

Postural hypotension can occur following administration of cabergoline. Care should be exercised when administering cabergoline concomitantly with other drugs known to lower blood pressure.

Somnolence/Sudden Sleep Onset

Cabergoline has been associated with somnolence. Dopamine agonists can be associated with sudden sleep onset episodes in patients with Parkinson's disease. Sudden onset of sleep during daily activities, in some cases without awareness or warning signs, has been reported uncommonly. Patients must be informed of this and advised to exercise caution while driving

or operating machines during treatment with cabergoline. Patients who have experienced somnolence and/or an episode of sudden sleep onset must refrain from driving or operating machines. Furthermore, a reduction in dosage or termination of therapy may be considered (See **section 4.4**)

Impulse control disorders

Patients should be regularly monitored for the development of impulse control disorders. Patients and carers should be made aware that behavioral symptoms of impulse control disorders including pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists including cabergoline. Dose reduction/tapered discontinuation should be considered if such symptoms develop.

Inhibition/suppression of physiological lactation

As with other ergot derivatives, cabergoline should not be used in women with pregnancy-induced hypertension, for example, preeclampsia or post-partum hypertension, unless the potential benefit is judged to outweigh the possible risk.

Serious adverse events including hypertension, myocardial infarction, seizures, stroke or psychiatric disorders have been reported in postpartum women treated with cabergoline for inhibition of lactation. In some patients the development of seizures or stroke was preceded by severe headache and/or transient visual disturbances. Blood pressure should be carefully monitored after the treatment. If hypertension, suggestive chest pain, severe, progressive, or unremitting headache (with or without visual disturbances), or evidence of central nervous system toxicity develop, cabergoline should be discontinued and the patient should be evaluated promptly.

In the reported post-partum studies with cabergoline, blood pressure decreases were mostly asymptomatic and were frequently observed on a single occasion 2 to 4 days after treatment. Since decreases in blood pressure are frequently noted during the puerperium, independently of drug therapy, it is likely that many of the observed decreases in blood pressure after cabergoline administration were not drug-induced. However, periodic monitoring of blood pressure, particularly during the first few days after cabergoline administration, is advised.

A single dose of 0.25 mg of cabergoline should not be exceeded in nursing women treated for suppression of established lactation to avoid potential postural hypotension (see **section 4.2**). A reported clinical study explored the efficacy and tolerability of 0.5 mg of cabergoline given as a single dose for suppression of lactation had shown that the risk of side effects is approximately doubled in this indication if the drug is administered as a single dose of 0.5 mg.

Treatment of hyperprolactinaemic disorders

Because hyperprolactinemia accompanied with amenorrhoea/galactorrhoea and infertility may be associated with pituitary tumour, a complete evaluation of the pituitary is indicated before treatment with cabergoline is initiated.

Cabergoline restores ovulation and fertility in women with hyperprolactinaemic hypogonadism.

Because pregnancy might occur prior to reinitiation of menses, a pregnancy test is recommended at least every four weeks during the amenorrhoeic period and, once menses are reinitiated, every time a menstrual period is delayed by more than three days. Women who wish to avoid pregnancy should be advised to use mechanical contraception during treatment with cabergoline and after discontinuation of cabergoline until recurrence of anovulation. As a precautionary measure, women who become pregnant should be monitored to detect signs of pituitary enlargement since expansion of pre-existing pituitary tumours may occur during gestation.

Before administration of cabergoline, pregnancy should be excluded. Because clinical experience is still limited and the product has a long half-life, as a precautionary measure it is recommended that once regular ovulatory cycles have been achieved women seeking pregnancy discontinue cabergoline one month before intended conception. Should pregnancy occur during treatment, cabergoline is to be discontinued. As a precautionary measure, women who become pregnant should be monitored to detect signs of pituitary enlargement since expansion of pre-existing pituitary tumours may occur during gestation.

Regular gynaecological assessment, including cervical and endometrial cytology, is recommended for patients taking cabergoline for extensive periods.

Fibrosis and cardiac valvulopathy and possibly related clinical phenomena

Fibrotic and serosal inflammatory disorders such as pleuritis, pleural effusion, pleural fibrosis, pulmonary fibrosis, pericarditis, pericardial effusion, cardiac valvulopathy involving one or more valves (aortic, mitral and tricuspid) or retroperitoneal fibrosis have occurred after prolonged usage of ergot derivatives with agonist activity at the serotonin 5HT_{2B} receptor, such as cabergoline. In some cases, symptoms or manifestations of cardiac valvulopathy improved after discontinuation of cabergoline.

Erythrocyte sedimentation rate (ESR) has been found to be abnormally increased in association with pleural effusion/fibrosis. Chest x-ray examination is recommended in cases of unexplained ESR increases to abnormal values.

Valvulopathy has been associated with cumulative doses, therefore, patients should be treated with the lowest effective dose. At each visit, the risk benefit profile of cabergoline treatment for the patient should be reassessed to determine the suitability of continued treatment with cabergoline.

Before initiating long-term treatment

All patients must undergo a cardiovascular evaluation, including echocardiogram to assess the potential presence of asymptomatic valvular disease. It is also appropriate to perform baseline investigations of erythrocyte sedimentation rate or other inflammatory markers, lung function/chest X-ray and renal function prior to initiation of therapy. In patients with valvular regurgitation, it is not known whether cabergoline treatment might worsen the underlying disease. If fibrotic valvular disease is detected, the patient should not be treated with cabergoline (see **section 4.3**).

During long-term treatment

Fibrotic disorders can have an insidious onset and patients should be regularly monitored for possible manifestations of progressive fibrosis. Therefore, during treatment, attention should be paid to the signs and symptoms of:

- Pleuro-pulmonary disease such as Dyspnoea, shortness of breath, persistent cough or chest pain.
- Renal insufficiency or ureteral/abdominal vascular obstruction that may occur with pain in the loin/flank and lower limb oedema as well as any possible abdominal masses or tenderness that may indicate retroperitoneal fibrosis.
- Cardiac failure: cases of valvular and pericardial fibrosis have often manifested as cardiac failure. Therefore, valvular fibrosis (and constrictive pericarditis) should be excluded if such symptoms occur.

Clinical diagnostic monitoring for development of fibrotic disorders, as appropriate, is essential. Following treatment initiation, the first echocardiogram must occur within 3-6 months, thereafter, the frequency of echocardiographic monitoring should be determined by appropriate individual clinical assessment with particular emphasis on the above-mentioned signs and symptoms, but must occur at least every 6 to 12 months.

Cabergoline should be discontinued if an echocardiogram reveals new or worsened valvular regurgitation, valvular restriction or valve leaflet thickening (see **section 4.3**). The need for other clinical monitoring (e.g. physical examination including, cardiac auscultation, X-ray, CT scan) should be determined on an individual basis.

Additional appropriate investigations such as erythrocyte sedimentation rate, and serum creatinine measurements should be performed if necessary to support a diagnosis of a fibrotic disorder.

4.5 *Interaction with other medicinal products and other forms of interaction*

The concomitant use of other drugs during early puerperium, particularly of ergot alkaloids, was not associated with detectable interactions modifying the efficacy and safety of cabergoline.

No information is available about the interaction between cabergoline and other ergot alkaloids; therefore, the concomitant use of these medications during long-term treatment with cabergoline is not recommended.

Since cabergoline exerts its therapeutic effect by direct stimulation of dopamine receptors, it should not be concurrently administered with drugs which have dopamine-antagonist activity (such as phenothiazines, butyrophenones, thioxanthenes, metoclopramide) since these might reduce the prolactin-lowering effect of cabergoline.

As with other ergot derivatives, cabergoline should not be used with macrolide antibiotics (e.g. erythromycin) due to increased systemic bioavailability of cabergoline.

4.6 *Fertility, pregnancy and lactation*

There are no adequate and well-controlled studies from the use of cabergoline in pregnant women. The reported animal studies have not demonstrated teratogenic effects, but reduced fertility and embryo-toxicity were observed in association with pharmacodynamic activity (see **section 5.3**).

In a reported twelve-year observational study on pregnancy outcomes following cabergoline therapy, information is available on 256 pregnancies. Seventeen of these 256 pregnancies (6.6%) eventuated in major congenital malformations or abortion. Information is available on 23/258 infants who had a total of 27 neonatal abnormalities, both major and minor. Musculoskeletal malformations were the most common neonatal abnormality (10), followed by cardio-pulmonary abnormalities (5). There is no information on perinatal disorders or long-term development of infants exposed to intra-uterine cabergoline. Based on recent published literature, the prevalence of major congenital malformations in the general population has been reported to be 6.9% or greater. Rates of congenital abnormality vary between different populations. It is not possible to accurately determine if there is an increased risk as no control group was included.

Cabergoline should only be used during pregnancy if clearly indicated and after an accurate benefit/risk evaluation (See **section 4.4**).

Due to the long half-life of the drug and limited data on in utero exposure, women planning to become pregnant should discontinue cabergoline one month before intended conception. If conception occurs during therapy, treatment should be discontinued as soon as pregnancy is confirmed to limit foetal exposure to the drug.

In rats, cabergoline and/or its metabolites are excreted in milk. No information is available on the excretion in breast milk in humans; however, mothers should be advised not to breast-feed in case of failed lactation inhibition/suppression by cabergoline. Since it prevents lactation, cabergoline should not be administered to mothers with hyperprolactinemic disorders who wish to breast-feed their infants.

4.7 *Effects on ability to drive and use machines*

Patients should be careful when performing actions which require fast and accurate reaction during treatment initiation.

During the first days of cabergoline administration, patients should be cautioned about re-engaging in activities requiring rapid and precise responses such as driving an automobile or operating machinery.

Patients being treated with cabergoline and presenting with somnolence must be informed to refrain from driving or engaging in activities where impaired alertness may put themselves and others at risk of serious injury or death (e.g. operating machines) until such episodes and somnolence have resolved. (See **section 4.4**).

4.8 *Undesirable effects*

Adverse events are generally dose-related. In patients known to be intolerant to dopaminergic drugs, the likelihood of adverse events may be lessened by starting therapy with cabergoline at reduced doses, e.g. 0.25 mg once a week, with subsequent gradual increase until the therapeutic dosage is reached. If persistent or severe adverse events occur, temporary reduction of dosage followed by a more gradual increase, e.g. increments of 0.25 mg/week every two weeks, may increase tolerability.

The following undesirable effects have been reported during treatment with cabergoline with the following frequencies: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$);

uncommon ($\geq 1/1,000$ to $\leq 1/100$); rare ($\geq 1/10,000$ to $\leq 1/1,000$); very rare ($\leq 1/10,000$), not known (cannot be estimated from the available data).

MedDRA System Organ Class	Frequency	Undesirable Effects
Cardiac disorders	Very Common	Valvulopathy (including regurgitation) and related disorders (pericarditis and pericardial effusion)
	Uncommon	Palpitations
	Not Known	Angina pectoris
Respiratory, thoracic and mediastinal disorders	Uncommon	Dyspnea, pleural effusion, fibrosis, (including pulmonary fibrosis), epistaxis
	Very rare	Pleural fibrosis
	Not Known	Respiratory disorder, respiratory failure, pleuritis, chest pain
Immune system disorders	Uncommon	Hypersensitivity reaction
Nervous system disorders	Very common	Headache [*] , dizziness/vertigo [*]
	Common	somnolence
	Uncommon	Transient hemianopsia, syncope, paresthesia
	Not Known	Sudden sleep onset, tremor
Eye disorders	Not Known	Visual impairment
Psychiatric disorders	Common	Depression
	Uncommon	Increased libido
	Not Known	Aggression, delusions, hypersexuality, pathological gambling, psychotic disorder, hallucinations
Vascular disorders	Common	Cabergoline generally exerts a hypotensive effect in patients on long-term treatment; Postural hypotension, hot flushes ^{**}
	Uncommon	Digital vasospasm, fainting
Gastrointestinal disorders	Very common	Nausea [*] , dyspepsia, gastritis, abdominal pain [*]
	Common	Constipation, vomiting ^{**}
	Rare	Epigastric pain
General disorders and administration site conditions	Very Common	Asthenia ^{***} , fatigue
	Uncommon	Oedema, peripheral oedema
Hepato-biliary disorders	Not Known	Hepatic function abnormal
Skin and subcutaneous tissue disorders	Uncommon	Rash, alopecia
Musculoskeletal and connective tissue disorders	Uncommon	Leg cramps
Reproductive system and breast disorders	Common	Breast pain

Investigations	Common	Asymptomatic decreases in blood pressure (≥ 20 mmHg systolic and ≥ 10 mmHg diastolic)
	Uncommon	A decrease in haemoglobin values have been observed in amenorrhoeic women during the first few months after menses.
	Not Known	Blood creatinine phosphokinase increased, liver function tests abnormal

*Very common in patients treated for hyperprolactinemia disorders; Common in patients treated for inhibition/suppression of lactation

** Common in patients treated for hyperprolactinemia disorders; Uncommon in patients treated for inhibition/suppression of lactation

*** Very common in patients treated for hyperprolactinemia disorders; Uncommon in patients treated for inhibition/suppression of lactation

Impulse control disorders

Pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists including cabergoline (see **section 4.4**).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after Authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Pharmacovigilance

Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org/>

4.9 Overdose

Symptoms of overdose would likely be those of over-stimulation of dopamine receptors e.g. nausea, vomiting, gastric complaints, postural hypotension, confusion/psychosis or hallucinations.

Supportive measures should be taken to remove any unabsorbed drug and maintain blood pressure, if necessary. In addition, the administration of dopamine antagonist drugs may be advisable.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Prolactin inhibitors, ATC code: G02CB03.

Cabergoline is a dopaminergic ergoline derivative endowed with a potent and long-lasting PRL-lowering activity. It acts by direct stimulation of the D₂-dopamine receptors on pituitary lactotrophs, thus inhibiting PRL secretion. In rats the compound decreases PRL

secretion at oral doses of 3-25 mcg/kg, and *in-vitro* at a concentration of 45 pg/ml. In addition, cabergoline exerts a central dopaminergic effect via D₂ receptor stimulation at oral doses higher than those effective in lowering serum PRL levels. The long lasting PRL-lowering effect of cabergoline is probably due to its long persistence in the target organ as

suggested by the slow elimination of total radioactivity from the pituitary after single oral dose in rats ($t_{1/2}$ of approximately 60 hours).

The pharmacodynamic effects of cabergoline have been studied in healthy volunteers, puerperal women and hyperprolactinaemic patients. After a single oral administration of cabergoline (0.3 - 1.5 mg), a significant decrease in serum PRL levels was reported in each of the populations studied. The effect is prompt (within 3 hours from administration) and persistent (up to 7 - 28 days in healthy volunteers and hyperprolactinaemic patients, and up to 14 - 21 days in puerperal women). The PRL-lowering effect is dose-related both in terms of degree of effect and duration of action.

With regard to the endocrine effects of cabergoline not related to the antiprolactinemic effect, available data from humans confirm the experimental findings in animals indicating that the test compound is endowed with a very selective action with no effect on basal secretion of other pituitary hormones or cortisol. The pharmacodynamic actions of cabergoline not correlated with the therapeutic effect only relate to blood pressure decrease. The maximal hypotensive effect of cabergoline as single dose usually occurs during the first 6 hours after drug intake and is dose-dependent both in terms of maximal decrease and frequency.

5.2 Pharmacokinetic properties

The pharmacokinetic and metabolic profiles of cabergoline have been studied in healthy volunteers of both sexes and in female hyperprolactinaemic patients.

After oral administration of the labelled compound, radioactivity was rapidly absorbed from the gastrointestinal tract as the peak of radioactivity in plasma was between 0.5 and 4 hours.

Ten days after administration about 18% and 72% of the radioactive dose was recovered in urine and faeces, respectively. Unchanged drug in urine accounted for 2-3% of the dose.

In urine, the main metabolite identified was 6-allyl-8 β -carboxy-ergoline, which accounted for 4-6% of the dose. Three additional metabolites were identified in urine, which accounted overall for less than 3% of the dose. The metabolites have been found to be much less potent than cabergoline in inhibiting prolactin secretion *in vitro*. Cabergoline biotransformation was also reportedly studied in plasma of healthy male volunteers treated with [14 C]-cabergoline: a rapid and extensive biotransformation of cabergoline was shown.

The low urinary excretion of unchanged cabergoline has been confirmed also in the reported studies with non-radioactive product. The elimination half-life of cabergoline,

estimated from urinary excretion rates, is long (63-68 hours in healthy volunteers (using a radio-immuno assay), 79-115 hours in hyperprolactinaemic patients (using a HPLC method).

On the basis of the elimination half-life, steady state conditions should be achieved after 4 weeks, as confirmed by the mean peak plasma levels of cabergoline obtained after a single dose (37 ± 8 pg/ml) and after a 4-week multiple regimen (101 ± 43 pg/ml).

In vitro experiments showed that the drug at concentrations of 0.1-10 ng/ml is 41-42% bound to plasma proteins. Food does not appear to affect absorption and disposition of cabergoline.

5.2 Preclinical safety data

There were maternotoxic effects but no teratogenic effects in mice given cabergoline at doses up to 8 mg/kg/day (approximately 55 times the maximum recommended human dose) during the period of organogenesis.

A dose of 0.012 mg/kg/day (approximately 1/7 the maximum recommended human dose) during the period of organogenesis in rats caused an increase in post-implantation embryo foetal losses. These losses could be due to the prolactin inhibitory properties of cabergoline in rats. At daily doses of 0.5 mg/kg/day (approximately 19 times the maximum recommended human dose) during the period of organogenesis in the rabbit, cabergoline caused materno-toxicity characterized by a loss of body weight and decreased food consumption. Doses of 4 mg/kg/day (approximately 150 times the maximum recommended human dose) during the period of organogenesis in the rabbit caused an increased occurrence of various malformations. However, in another reported study in rabbits, no treatment-related malformations or embryofetotoxicity were observed at doses up to 8 mg/kg/day (approximately 300 times the maximum recommended human dose).

6. PHARMACEUTICAL PARTICULARS

6.1 *List of excipients:*

Microcrystalline cellulose, L Leucine

6.2 *Incompatibilities*

Not Applicable

6.3 *Shelf life*

24 Months

6.4 *Special precautions for storage*

Store up to 30 °C and protected from light.

6.5 *Nature and contents of container*

CABERLIN 0.5 is available in 2 by 2; 1 by 4 and 1 by 10 pack

6.6 *Special precautions for disposal and other handling*

No Special requirement.

7. *MARKETING AUTHORISATION HOLDER AND MANUFACTURING SITE ADDRESS*

Sun Pharmaceutical Industries Limited

8. *MARKETING AUTHORISATION NUMBER*

H2009/19598/168

9. *DATE OF FIRST REGISTRATION / RENEWAL OF THE REGISTRATION*

-

10. *DATE OF REVISION OF THE TEXT*

February 2024

REFERENCES

1. Summary of Product Characteristics of Dostinex Tablets 0.5 mg, Pfizer Limited, UK, October 2020.

Information last compiled in February 2024

Dostinex is the registered trademark of Pfizer Limited, UK and is not the trademarks of Sun Pharmaceutical Industries Limited. The maker of this brand is not affiliated with and does not endorse Sun Pharmaceutical Industries Limited or its products.