

Summary of Product Characteristics for Pharmaceutical Products

1. Name of the Medicinal Product

Cazithro 500

2. Quality and Quantitative Composition

Each Film coated tablet contains

Azithromycin Dihydrate USP

Eq.to. Azithromycin anhydrous 500 mg

Excipients q.s.

For the full list of excipients, see section 6.1.

3. Pharmaceutical Form

Solid Dosage form (Film coated tablets)

4. Clinical particulars:

4.1 Therapeutic indications:

Azithromycin is indicated for the following bacterial infections induced by micro-organisms susceptible to azithromycin:

- Acute bacterial sinusitis (adequately diagnosed)
- Acute bacterial otitis media (adequately diagnosed)
- Pharyngitis, tonsillitis
- Acute exacerbation of chronic bronchitis (adequately diagnosed)
- Mild to moderately severe community acquired pneumonia
- Infections of the skin and soft tissues of mild to moderate severity e.g. folliculitis, cellulitis, erysipelas.
- Uncomplicated Chlamydia trachomatis urethritis and cervicitis

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration:

Azithromycin should be given as a single daily dose. Duration of the treatment for the different infection diseases is given below.

The tablets can be taken with or without food. The tablets should be taken with ½ glass of water.

Children and adolescents with a body weight above 45 kg, adults and the elderly:

The total dose is 1500 mg, administered as 500 mg once daily for 3 days. Alternatively, the same total dose (1500 mg) can be administered in a period of 5 days, 500 mg on the first day and 250 mg on day 2 to 5.

In the case of uncomplicated *Chlamydia trachomatis* urethritis and cervicitis, the dosage is 1000 mg as a single oral dose.

Children and adolescents with a body weight below 45 kg:

Azithromycin tablets are not suitable for patients under 45 kg body weight. Other dosage forms are available for this group of patients.

Elderly patients

For elderly patients the same dose as for adults can be applied. Since elderly patients can be patients with ongoing proarrhythmic conditions a particular caution is recommended due to the risk of developing cardiac arrhythmia and torsades de pointes.

Patients with renal impairment:

Dose adjustment is not required in patients with mild to moderate renal impairment (GFR 10-80 ml/min).

Patients with hepatic impairment:

Dose adjustment is not required for patients with mild to moderate hepatic dysfunction.

4.2 Contraindications

The use of this product is contraindicated in patients with hypersensitivity to azithromycin, erythromycin, any macrolide or ketolide antibiotic, or to any of the excipients.

4.3 Special warnings and precautions for use

As with erythromycin and other macrolides, rare serious allergic reactions, including angioedema and anaphylaxis (rarely fatal), have been reported.

Some of these reactions with azithromycin have resulted in recurrent symptoms and required a longer period of observation and treatment. Since liver is the principal route of elimination for azithromycin, the use of azithromycin should be undertaken with caution in patients with significant hepatic disease. Cases of fulminant hepatitis potentially leading to life-threatening liver failure have been reported with azithromycin. Some patients may have had pre-existing hepatic disease or may have been taking other hepatotoxic medicinal products.

In case of signs and symptoms of liver dysfunction, such as rapid developing asthenia associated with jaundice, dark urine, bleeding tendency or hepatic encephalopathy, liver function tests / investigations should be performed immediately. Azithromycin administration should be stopped if liver dysfunction has emerged.

In patients receiving ergot derivatives, ergotism has been precipitated by co-administration of some macrolide antibiotics. There are no data concerning the possibility of an interaction between ergot and azithromycin. However, because of the theoretical possibility of ergotism, azithromycin and ergot derivatives should not be coadministered.

As with any antibiotic preparation, observation for signs of superinfection with non-susceptible organisms, including fungi is recommended.

Clostridium difficile associated diarrhoea (CDAD) has been reported with the use of nearly all antibacterial agents, including azithromycin, and may range in severity from mild diarrhoea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

4.4 Interaction with other medicinal products and other forms of interaction

Antacids

In a pharmacokinetic study investigating the effects of simultaneous administration of antacid with azithromycin, no effect on overall

bioavailability was seen although peak serum concentrations were reduced by approximately 25%.

Cetirizine

In healthy volunteers, coadministration of a 5-day regimen of azithromycin with cetirizine 20 mg at steady-state resulted in no pharmacokinetic interaction and no significant changes in the QT interval.

Didanosine (Dideoxyinosine)

Coadministration of 1200 mg/day azithromycin with 400 mg/day didanosine in 6 HIV-positive subjects did not appear to affect the steady-state pharmacokinetics of didanosine as compared with placebo.

Digoxin (P-gp substrates)

Concomitant administration of macrolide antibiotics, including azithromycin, with P-glycoprotein substrates such as digoxin, has been reported to result in increased serum levels of the P-glycoprotein substrate. Therefore, if azithromycin and P-gp substrates such as digoxin are administered concomitantly, the possibility of elevated serum concentrations of the substrate should be considered.

Zidovudine

Single 1000 mg doses and multiple 1200 mg or 600 mg doses of azithromycin had little effect on the plasma pharmacokinetics or urinary excretion of zidovudine or its glucuronide metabolite. However, administration of azithromycin increased the concentrations of phosphorylated zidovudine, the clinically active metabolite, in peripheral blood mononuclear cells. The clinical significance of this finding is unclear, but it may be of benefit to patients.

Azithromycin does not interact significantly with the hepatic cytochrome P450 system. It is not believed to undergo the pharmacokinetic drug interactions as seen with erythromycin and other macrolides. Hepatic cytochrome P450 induction or inactivation via cytochrome-metabolite complex does not occur with azithromycin.

Ergot

Due to the theoretical possibility of ergotism, the concurrent use of azithromycin with ergot derivatives is not recommended.

Pharmacokinetic studies have been conducted between azithromycin and the following drugs known to undergo significant cytochrome P450 mediated metabolism.

Ergotamine derivatives: Due to the theoretical possibility of ergotism, the concurrent use of azithromycin with ergot derivatives is not recommended.

Astemizole, alfentanil

There are no known data on interactions with astemizole or alfentanil. Caution is advised in the co-administration of these medicines with Azithromycin because of the known enhancing effect of these medicines when used concurrently with the macrolid antibiotic erythromycin.

Atorvastatin

Coadministration of atorvastatin (10 mg daily) and azithromycin (500 mg daily) did not alter the plasma concentrations of atorvastatin (based on a HMG CoA-reductase inhibition assay). However, post-marketing cases of rhabdomyolysis in patients receiving azithromycin with statins have been reported.

Carbamazepine

In a pharmacokinetic interaction study in healthy volunteers, no significant effect was observed on the plasma levels of carbamazepine or its active metabolite in patients receiving concomitant azithromycin.

Cisapride

Cisapride is metabolized in the liver by the enzyme CYP 3A4. Because macrolides inhibit this enzyme, concomitant administration of cisapride may cause the increase of QT interval prolongation, ventricular arrhythmias and torsades de pointes.

Cimetidine

In a pharmacokinetic study investigating the effects of a single dose of cimetidine, given 2 hours before azithromycin, on the pharmacokinetics of

azithromycin, no alteration of azithromycin pharmacokinetics was seen.

Coumarin-Type Oral Anticoagulants

In a pharmacokinetic interaction study, azithromycin did not alter the anticoagulant effect of a single 15 mg dose of warfarin administered to healthy volunteers. There have been reports received in the post-marketing period of potentiated anticoagulation subsequent to coadministration of azithromycin and coumarin-type oral anticoagulants. Although a causal relationship has not been established, consideration should be given to the frequency of monitoring prothrombin time when azithromycin is used in patients receiving coumarin-type oral anticoagulants.

Cyclosporin

In a pharmacokinetic study with healthy volunteers that were administered a 500 mg/day oral dose of azithromycin for 3 days and were then administered a single 10 mg/kg oral dose of cyclosporin, the resulting cyclosporin C_{max} and AUC₀₋₅ were found to be significantly elevated. Consequently, caution should be exercised before considering concurrent administration of these drugs. If coadministration of these drugs is necessary, cyclosporin levels should be monitored and the dose adjusted accordingly.

Efavirenz

Coadministration of a 600 mg single dose of azithromycin and 400 mg efavirenz daily for 7 days did not result in any clinically significant pharmacokinetic interactions.

Fluconazole

Coadministration of a single dose of 1200 mg azithromycin did not alter the pharmacokinetics of a single dose of 800 mg fluconazole. Total exposure and half-life of azithromycin were unchanged by the coadministration of fluconazole, however, a clinically insignificant decrease in C_{max} (18%) of azithromycin was observed.

Indinavir

Coadministration of a single dose of 1200 mg azithromycin had no statistically significant effect on the pharmacokinetics of indinavir administered as 800 mg three times daily for 5 days.

Methylprednisolone

In a pharmacokinetic interaction study in healthy volunteers, azithromycin had no significant effect on the pharmacokinetics of methylprednisolone.

Midazolam

In healthy volunteers, coadministration of azithromycin 500 mg/day for 3 did not cause clinically significant changes in the pharmacokinetics and pharmacodynamics of a single 15 mg dose of midazolam.

Nelfinavir

Coadministration of azithromycin (1200 mg) and nelfinavir at steady state (750 mg three times daily) resulted in increased azithromycin concentrations. No clinically significant adverse effects were observed and no dose adjustment is required.

Rifabutin

Coadministration of azithromycin and rifabutin did not affect the serum concentrations of either drug.

Neutropenia was observed in subjects receiving concomitant treatment of azithromycin and rifabutin. Although neutropenia has been associated with the use of rifabutin, a causal relationship to combination with azithromycin has not been established.

Sildenafil

In normal healthy male volunteers, there was no evidence of an effect of azithromycin (500 mg daily for 3 days) on the AUC and C_{max} of sildenafil or its major circulating metabolite. Terfenadine

Pharmacokinetic studies have reported no evidence of an interaction between azithromycin and terfenadine. There have been rare cases reported where the possibility of such an interaction could not be entirely

excluded; however there was no specific evidence that such an interaction had occurred.

Theophylline

There is no evidence of a clinically significant pharmacokinetic interaction when azithromycin and theophylline are co-administered to healthy volunteers.

Triazolam

In 14 healthy volunteers, coadministration of azithromycin 500 mg on Day 1 and 250 mg on Day 2 with 0.125 mg triazolam on Day 2 had no significant effect on any of the pharmacokinetic variables for triazolam compared to triazolam and placebo.

Trimethoprim/sulfamethoxazole

Coadministration of trimethoprim/sulfamethoxazole (160 mg/800 mg) for 7 days with azithromycin 1200 mg on Day 7 had no significant effect on peak concentrations, total exposure or urinary excretion of either trimethoprim or sulfamethoxazole. Azithromycin serum concentrations were similar to those seen in other studies.

Substances that prolong the QT interval

Azithromycin should not be used concurrently with other active substances that prolong the QT interval.

4.5 Pregnancy and lactation

Pregnancy

There are no adequate data from the use of Azithromycin in pregnant women. In reproduction toxicity studies in animals azithromycin was shown to pass the placenta, but no teratogenic effects were observed. The safety of azithromycin has not been confirmed with regard to the use of the active substance during pregnancy. Therefore Azithromycin should only be used during pregnancy if the benefit outweighs the risk.

Lactation

Azithromycin has been reported to be secreted into human breast milk,

but there are no adequate and well-controlled clinical studies in nursing women that have characterized the pharmacokinetics of azithromycin excretion into human breast milk.

Fertility

In fertility studies conducted in rat, reduced pregnancy rates were noted following administration of azithromycin. The relevance of this finding to humans is unknown.

4.6 Effects on ability to drive and use machines

Not known

4.7 Undesirable effects

The frequency grouping is defined using the following convention: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); and not known (cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Adverse reactions possibly or probably related to azithromycin based on clinical trial experience and post-marketing surveillance:

Infections and Infestations:

Uncommon ($\geq 1/1000$ to $< 1/100$): Candidiasis, Vaginal infection, Pneumonia, Fungal infection, Bacterial infection, Pharyngitis, Gastroenteritis, Respiratory disorder, Rhinitis, Oral candidiasis.

Frequency Not Known: Pseudomembranous colitis

Blood and Lymphatic System Disorders:

Uncommon ($\geq 1/1000$ to $< 1/100$): Leukopenia, Neutropenia, Eosinophilia.

Frequency Not Known: Thrombocytopenia, Haemolytic anaemia.

Immune System Disorders:

Uncommon ($\geq 1/1000$ to $< 1/100$): Angioedema, Hypersensitivity. Frequency

Not Known: Anaphylactic reaction.

Metabolism and Nutrition Disorders:

Uncommon ($\geq 1/1000$ to $< 1/100$): Anorexia

Psychiatric Disorders:

Uncommon ($\geq 1/1000$ to $< 1/100$): Nervousness,

Insomnia. Rare ($\geq 1/10,000$ to $< 1/1,000$): Agitation

Frequency Not Known: Aggression, Anxiety, Delirium, Hallucination.

Nervous System Disorders:

Common ($\geq 1/100$ to $< 1/10$): Headache

Uncommon ($\geq 1/1000$ to $< 1/100$): Dizziness, Somnolence, Dysgeusia, Paraesthesia. Frequency Not Known: Syncope, convulsion, Hypoesthesia, Psychomotor hyperactivity, Anosmia, Ageusia, Parosmia, Myasthenia gravis.

4.8 Overdose

Adverse events experienced in higher than recommended doses were similar to those seen at normal doses. In the event of overdosage, general symptomatic and supportive measures are indicated as required

Reporting of suspected adverse reactions

Healthcare professionals are requested to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org>

5. Pharmacological properties

5.1 Pharmacodynamic properties

General properties

Pharmacotherapeutic group: antibacterial for systemic use; macrolides

Antibiotics; azithromycin, ATC code: J01FA10

Mode of action

Azithromycin is an azalide, a sub-class of the macrolide antibiotics. By binding to the 50S-ribosomal sub-unit, azithromycin avoids the translocation of peptide chains from one side of the ribosome to the other. As a consequence of this, RNA-dependent protein synthesis is sensitive

organisms is prevented.

PK/PD relationship

For azithromycin the AUC/MIC is the major PK/PD parameter correlating best with the efficacy of azithromycin.

Mechanism of resistance: Resistance to azithromycin may be inherent or acquired. There are three main mechanisms of resistance in bacteria: target site alteration, alteration in antibiotic transport and modification of the antibiotic.

Complete cross resistance exists among *Streptococcus pneumoniae*, *betahaemolytic streptococcus* of group A, *Enterococcus faecalis* and *Staphylococcus aureus*, including methicillin resistant *S. aureus* (MRSA) to erythromycin, azithromycin, other macrolides and lincosamides.

5.2 Pharmacokinetic properties

Absorption

The biological availability of azithromycin after oral administration is approximately 37%. Peak plasma levels are achieved 2-3 hours after taking the medicinal product.

Distribution

After oral administration, azithromycin is distributed throughout the entire body. Pharmacokinetic studies have shown clearly higher azithromycin levels in the tissues than in the plasma (up to 50 times the maximum observed concentration in plasma). This indicates that the substance is bound in the tissues in considerable quantities.

Concentrations in the infected tissues, such as lungs, tonsil and prostate are higher than the MIC₉₀ of the most frequently occurring pathogens after a single dose of 500 mg.

The protein binding of azithromycin in serum is variable and varies, depending on the serum concentration, from 52% at 0.05 mg/l to 12% at 0.5 mg/l. The steady state distribution volume is 31.1 l/kg.

Elimination

The terminal plasma-elimination half-life closely follows the tissue depletion half-life from 2 to 4 days.

Approximately 12% of an intravenously administered dose of azithromycin is, over a period of 3 days, excreted unchanged in the urine. High concentrations of unchanged azithromycin were found in human bile. In this, ten metabolites were also detected (formed by N- and O-desmethylation, by hydroxylation of the desosamin and aglycon rings and by splitting the cladinose conjugate). A comparison of fluid chromatography and microbiological assessment methods shows that the metabolites are microbiologically inactive.

Pharmacokinetics in special populations

Renal insufficiency

Following a single oral dose of azithromycin 1 g, mean C_{max} and AUC₀₋₁₂₀ increased by 5.1% and 4.2% respectively, in subjects with mild to moderate renal impairment (glomerular filtration rate of 10-80 ml/min) compared with normal renal function (GFR > 80 ml/min). In subjects with severe renal impairment, the mean C_{max} and AUC₀₋₁₂₀ increased 61% and 33% respectively compared to normal.

Hepatic insufficiency

In patients with mild to moderate hepatic impairment, there is no evidence of a marked change in serum pharmacokinetics of azithromycin compared to normal hepatic function. In these patients, urinary recovery of azithromycin appears to increase perhaps to compensate for reduced hepatic clearance.

Elderly

The pharmacokinetics of azithromycin in elderly men was similar to that of young adults; however, in elderly women, although higher peak concentrations (increased by 30-50%) were observed, no significant accumulation occurred.

Infants, toddlers, children and adolescents

Pharmacokinetics have been studied in children aged 4 months – 15 years taking capsules, granules or suspension. At 10 mg/kg on day 1 followed by 5 mg/kg on days 2-5, the C_{max} achieved is slightly lower than adults with 224 ug/l in children aged 0.6-5 years and after 3 days dosing and 383 ug/l in those aged 6-15 years. The t_{1/2} of 36 h in the older children was within the expected range for adults.

5.3 Preclinical safety data

There are no pre-clinical data of relevance

6. Pharmaceutical particulars

6.1 List of excipients:

S. No.	Name of Material	Specification
1.	Dibasic Calcium Phosphate	BP
2.	Croscarmellose Sodium	USP NF
3.	Sodium lauryl Sulphate	BP
4.	Starch	BP
5.	Starch (for Paste)	BP
6.	Purified talc	BP
7.	Magnesium Stearate	BP
8.	Opadry White (85G68918)	IH
9.	Purified Water	BP

6.2 Incompatibilities

Not Applicable

6.3 Shelf life

24 months

6.4 Special precautions for storage

Store in a dry place, below 25°C. Protect from light. Keep out of reach of

6.5 Nature and contents of container

3 tablets packed in ALU /PVC blister and such 1 blister packed in a printed monocarton with pack insert.

6.6 Special precautions for disposal and other handling

Not Applicable

7. Marketing authorization Holder

Cachet Pharmaceuticals Pvt. Ltd

415, Shah Nahar Industrial Estate,

Dr. E. Moses Road, Worli, Mumbai-400 018,

Maharashtra, India

8. Marketing authorization Number

H2022/CTD9078/19818

9. Date of first authorisation/renewal of the authorisation

2023 May 03

10. Date of revision of the text

2023 May 03