

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE FINISHED PHARMACEUTICAL PRODUCT

CLAXY 1.2G

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each combipack contains:

a) One vial of Co-Amoxiclav for Injection BP 1.2 g

Each vial contains:

Amoxicillin Sodium (Sterile) BP

Equivalent to Amoxicillin 1000 mg

Potassium Clavulanate (Sterile) BP

Equivalent to Clavulanic Acid 200 mg

b) One Ampoule of Sterilised water for Injections BP 20ml

3. PHARMACEUTICAL FORM

Dry Powder for Injection

White to off white powder filled in transparent glass vial.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Co-Amoxiclav for Injection BP 1.2g is indicated for the treatment of the following infections in adults and children:

- Severe infections of the ear, nose and throat (such as mastoiditis, peritonsillar infections, epiglottitis, and sinusitis when accompanied by severe systemic signs and symptoms)
- Acute exacerbations of chronic bronchitis (adequately diagnosed)
- Community acquired pneumonia
- Cystitis
- Pyelonephritis
- Skin and soft tissue infections in particular cellulitis, animal bites, severe dental abscess with spreading cellulitis
- Bone and joint infections, in particular osteomyelitis
- Intra-abdominal infections
- Female genital infections.

Prophylaxis against infections associated with major surgical procedures in adults, such as those involving the:

- Gastrointestinal tract
- Pelvic cavity
- Head and neck

- Biliary tract surgery.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

Doses are expressed throughout in terms of amoxicillin/clavulanic acid content except when doses are stated in terms of an individual component.

-Posology

The dose of Co-Amoxiclav for Injection BP 1.2g that is selected to treat an individual infection should take into account:

- The expected pathogens and their likely susceptibility to antibacterial agents
- The severity and the site of the infection
- The age, weight and renal function of the patient as shown below.

The use of alternative presentations of Co-Amoxiclav for Injection BP 1.2g (e.g. those that provide higher doses of amoxicillin and/or different ratios of amoxicillin to clavulanic acid) should be considered as necessary.

This amoxicillin/clavulanic acid powder for solution for injection or infusion provides a total daily dose of 3000 mg amoxicillin and 600 mg clavulanic acid when administered as recommended below. If it is considered that a higher daily dose of amoxicillin is required, it is recommended that an alternative intravenous formulation of Co-Amoxiclav for Injection BP 1.2g is selected in order to avoid administration of unnecessarily high daily doses of clavulanic acid.

The duration of therapy should be determined by the response of the patient. Some infections (e.g. osteomyelitis) require longer periods of treatment. Treatment should not be extended beyond 14 days without review.

Consideration should be given to local guidelines on appropriate dosing frequencies for amoxicillin/clavulanic acid.

Adults and children $\geq$ 40 kg

For treatment of infections as indicated in section 4.1: 1000 mg/ 200 mg every 8 hours

For surgical prophylaxis:

For procedures less than 1 hour in duration, the recommended dose is 1000 mg/200 mg to 2000 mg/200 mg given at induction of anaesthesia (Doses of 2000 mg/200 mg can be achieved by using an alternative intravenous formulation of Co-amoxiclav).

For procedures greater than 1 hour in duration, the recommended dose is 1000 mg/200 mg to 2000 mg/200 mg given at induction of anaesthesia, with up to 3 doses of 1000 mg/200 mg in 24 hours.

Clear clinical signs of infection at operation will require a normal course of intravenous or oral therapy post-operatively.

Paediatric population

Children < 40 kg

Recommended doses:

- *Children aged 3 months and over:* 25 mg/5 mg per kg every 8 hours
- *Children aged less than 3 months or weighing less than 4 kg:* 25 mg/5 mg per kg every 12 hours.

Elderly

No dose adjustment is considered necessary.

Renal impairment

Dose adjustments are based on the maximum recommended level of amoxicillin.

No dose adjustment is required in patients with creatinine clearance (CrCl) greater than 30 ml/min.

Adults and children ≥ 40 kg

CrCl: 10-30 ml/min: Initial dose of 1000 mg/200 mg and then 500 mg/100 mg given twice daily.

CrCl < 10 ml /min: Initial dose of 1000 mg/200 mg and then 500 mg/100 mg given every 24 hours.

Haemodialysis: Initial dose of 1000 mg/200 mg and then followed by 500 mg/100 mg every 24 hours, plus a dose of 500 mg/100 mg at the end of dialysis (as serum concentrations of both amoxicillin and clavulanic acid are decreased).

Children < 40 kg

CrCl: 10-30 ml/min: 25 mg/5 mg per kg given every 12 hours.

CrCl < 10 ml /min: 25 mg/5 mg per kg given every 24 hours.

Haemodialysis: 25 mg/5 mg per kg given every 24 hours, plus a dose of 12.5 mg/2.5 mg per kg at the end of dialysis (as serum concentrations of both amoxicillin and clavulanic acid are decreased).

Hepatic impairment

Dose with caution and monitor hepatic function at regular intervals.

Method of administration

For intravenous use.

Co-Amoxiclav for Injection BP 1.2g may be administered either by slow intravenous injection over a period of 3 to 4 min directly into a vein or via a drip tube or by infusion over 30 to 40 min. Co-Amoxiclav for Injection BP 1.2g is not suitable for intramuscular administration.

Children aged less than 3 months should be administered amoxicillin/clavulanic acid by infusion only.

Treatment with amoxicillin/clavulanic acid may be initiated by the use of an intravenous preparation and completed with an appropriate oral presentation as considered appropriate for the individual patient.

4.3 Contraindications

Hypersensitivity to the active substances, to any of the penicillins or to any of the excipients.

History of a severe immediate hypersensitivity reaction (e.g. anaphylaxis) to another beta-lactam agent (e.g. a cephalosporin, carbapenem or monobactam).

History of jaundice/hepatic impairment due to amoxicillin/clavulanic acid.

4.4 Special warnings and precautions for use

Before initiating therapy with amoxicillin/clavulanic acid, careful enquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, or other beta-lactam agents.

Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous reactions) have been reported in patients on penicillin therapy. Hypersensitivity reactions can also progress to Kounis syndrome, a serious allergic reaction that can result in myocardial infarction. These reactions are more likely to occur in individuals with a history of penicillin hypersensitivity and in atopic individuals. If an allergic reaction occurs, amoxicillin/clavulanic acid therapy should be discontinued and appropriate alternative therapy instituted.

Drug-induced enterocolitis syndrome (DIES) has been reported mainly in children receiving amoxicillin/clavulanate. DIES is an allergic reaction with the leading symptom of protracted vomiting (1-4 hours after drug administration) in the absence of allergic skin or respiratory symptoms. Further symptoms could comprise abdominal pain, diarrhoea, hypotension or leucocytosis with neutrophilia. There have been severe cases including progression to shock.

In the case that an infection is proven to be due to an amoxicillin-susceptible organisms(s) then consideration should be given to switching from amoxicillin/clavulanic acid to amoxicillin in accordance with official guidance.

This presentation of amoxicillin/clavulanic acid may not be suitable for use when there is a high risk that the presumptive pathogens have resistance to beta-lactam agents that is not mediated by beta-lactamases susceptible to inhibition by clavulanic acid. As no specific data for $T > MIC$ are available and the data for comparable oral presentations are borderline, this presentation (without additional amoxicillin) may not be suitable for the treatment of penicillin-resistant *S. pneumoniae*. Convulsions may occur in patients with impaired renal function or in those receiving high doses. Amoxicillin/clavulanic acid should be avoided if infectious mononucleosis is suspected since the occurrence of a morbilliform rash has been associated with this condition following the use of amoxicillin.

Concomitant use of allopurinol during treatment with amoxicillin can increase the likelihood of allergic skin reactions.

Prolonged use may occasionally result in overgrowth of non-susceptible organisms.

The occurrence at the treatment initiation of a feverish generalised erythema associated with pustula may be a symptom of acute generalised exanthemous pustulosis (AGEP). This reaction requires amoxicillin/clavulanic acid discontinuation and contra-indicates any subsequent administration of amoxicillin.

Amoxicillin/clavulanic acid should be used with caution in patients with evidence of hepatic impairment.

Hepatic events have been reported predominantly in males and elderly patients and may be associated with prolonged treatment. These events have been very rarely reported in children. In all populations, signs and symptoms usually occur during or shortly after treatment but in some cases may not become apparent until several weeks after treatment has ceased. These are usually

reversible. Hepatic events may be severe and in extremely rare circumstances, deaths have been reported. These have almost always occurred in patients with serious underlying disease or taking concomitant medications known to have the potential for hepatic effects.

Antibiotic-associated colitis has been reported with nearly all antibacterial agents including amoxicillin and may range in severity from mild to life threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during or subsequent to the administration of any antibiotics. Should antibiotic-associated colitis occur, amoxicillin/clavulanic acid should immediately be discontinued, a physician be consulted and an appropriate therapy initiated. Anti-peristaltic drugs are contra-indicated in this situation.

Periodic assessment of organ system functions, including renal, hepatic and haematopoietic function is advisable during prolonged therapy.

Prolongation of prothrombin time has been reported rarely in patients receiving amoxicillin/clavulanic acid. Appropriate monitoring should be undertaken when anticoagulants are prescribed concomitantly. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation.

In patients with renal impairment, the dose should be adjusted according to the degree of impairment.

In patients with reduced urine output crystalluria (including acute renal injury) has been observed very rarely, predominantly with parenteral therapy. During administration of high doses of amoxicillin it is advisable to maintain adequate fluid intake and urinary output in order to reduce the possibility of amoxicillin crystalluria. In patients with bladder catheters, a regular check of patency should be maintained.

During treatment with amoxicillin, enzymatic glucose oxidase methods should be used whenever testing for the presence of glucose in urine because false positive results may occur with non-enzymatic methods.

The presence of clavulanic acid in amoxicillin/clavulanic acid may cause a non-specific binding of IgG and albumin by red cell membranes leading to a false positive Coombs test.

There have been reports of positive test results using the Bio-Rad Laboratories Platelia *Aspergillus* EIA test in patients receiving amoxicillin/clavulanic acid who were subsequently found to be free of *Aspergillus* infection. Cross-reactions with non-*Aspergillus* polysaccharides and polyfuranoses with Bio-Rad Laboratories Platelia *Aspergillus* EIA test have been reported. Therefore, positive test results in patients receiving amoxicillin/clavulanic acid should be interpreted cautiously and confirmed by other diagnostic methods.

4.5 Interaction with other medicinal products and other forms of interaction

Oral anticoagulants

Oral anticoagulants and penicillin antibiotics have been widely used in practice without reports of interaction. However, in the literature there are cases of increased international normalised ratio in patients maintained on acenocoumarol or warfarin and prescribed a course of amoxicillin. If co-administration is necessary, the prothrombin time or international normalised ratio should be

carefully monitored with the addition or withdrawal of amoxicillin. Moreover, adjustments in the dose of oral anticoagulants may be necessary.

Methotrexate

Penicillins may reduce the excretion of methotrexate causing a potential increase in toxicity.

Probenecid

Concomitant use of probenecid is not recommended. Probenecid decreases the renal tubular secretion of amoxicillin. Concomitant use of probenecid may result in increased and prolonged blood levels of amoxicillin.

Mycophenolate mofetil

In patients receiving mycophenolate mofetil, reduction in pre-dose concentration of the active metabolite mycophenolic acid (MPA) of approximately 50% has been reported following commencement of oral amoxicillin plus clavulanic acid. The change in pre-dose level may not accurately represent changes in overall MPA exposure.

Therefore, a change in the dose of mycophenolate mofetil should not normally be necessary in the absence of clinical evidence of graft dysfunction. However, close clinical monitoring should be performed during the combination and shortly after antibiotic treatment.

4.6 Fertility, pregnancy and lactation

Pregnancy

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development. Limited data on the use of amoxicillin/clavulanic acid during pregnancy in humans do not indicate an increased risk of congenital malformations. In a single study in women with preterm, premature rupture of the foetal membrane it was reported that prophylactic treatment with amoxicillin/clavulanic acid may be associated with an increased risk of necrotising enterocolitis in neonates. Use should be avoided during pregnancy, unless considered essential by the physician.

Breast-feeding

Both substances are excreted into breast milk (nothing is known of the effects of clavulanic acid on the breast-fed infant). Consequently, diarrhoea and fungus infection of the mucous membranes are possible in the breast-fed infant, so that breast-feeding might have to be discontinued. The possibility of sensitisation should be taken into account. Amoxicillin/clavulanic acid should only be used during breast-feeding after benefit/risk assessment by the physician in charge.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, undesirable effects may occur (e.g. allergic reactions, dizziness, convulsions), which may influence the ability to drive and use machines.

4.8 Undesirable effects

The most commonly reported adverse drug reactions (ADRs) are diarrhoea, nausea and vomiting.

The ADRs derived from clinical studies and post-marketing surveillance with amoxicillin/clavulanic acid, sorted by MedDRA System Organ Class are listed below.

The following terminologies have been used in order to classify the occurrence of undesirable effects.

Very common ($\geq 1/10$)

Common ($\geq 1/100$ to $< 1/10$)

Uncommon ($\geq 1/1,000$ to $< 1/100$)

Rare ($\geq 1/10,000$ to $< 1/1,000$)

Very rare ($< 1/10,000$)

Not known (cannot be estimated from the available data)

Infections and infestations

Common: Mucocutaneous candidosis

Not known: Overgrowth of non-susceptible organisms

Blood and lymphatic system disorders

Rare: Reversible leucopenia (including neutropenia), Thrombocytopenia

Not known: Reversible agranulocytosis, Haemolytic anaemia, Prolongation of bleeding time and prothrombin time

Cardiac disorders

Not known: Kounis syndrome

Immune system disorders

Not known: Angioneurotic oedema, Anaphylaxis, Serum sickness-like syndrome, Hypersensitivity vasculitis

Nervous system disorders

Uncommon: Dizziness, Headache

Not known: Convulsions, Aseptic meningitis

Vascular disorders

Rare: Thrombophlebitis

Gastrointestinal disorders

Common: Diarrhoea

Uncommon: Nausea, Vomiting, Indigestion

Not known: Antibiotic-associated colitis, Drug-induced enterocolitis syndrome, Pancreatitis acute

Hepatobiliary disorders

Uncommon: Rises in AST and/or ALT

Not known: Hepatitis, Cholestatic jaundice

Skin and subcutaneous tissue disorders

Uncommon: Skin rash, Pruritus, Urticaria

Rare: Erythema multiforme

Not known: Stevens-Johnson syndrome, Toxic epidermal necrolysis, Bullous exfoliative dermatitis, Acute generalised exanthematous pustulosis (AGEP), Drug reaction with eosinophilia and systemic symptoms (DRESS), Linear IgA disease

Renal and urinary disorders

Not known: Interstitial nephritis, Crystalluria (including acute renal injury)

Reporting of suspected adverse reactions

Reporting of suspected adverse reactions: Healthcare professionals are requested to report any suspected adverse reactions via the Pharmacy and Poisons Reporting System ([PvERS](https://pv.pharmacyboardkenya.org)) <https://pv.pharmacyboardkenya.org>

4.9 Overdose

Symptoms and signs of overdose

Gastrointestinal symptoms and disturbance of the fluid and electrolyte balances may be evident. Amoxicillin crystalluria, in some cases leading to renal failure, has been observed.

Convulsions may occur in patients with impaired renal function or in those receiving high doses. Amoxicillin has been reported to precipitate in bladder catheters, predominantly after intravenous administration of large doses. A regular check of patency should be maintained.

Treatment of intoxication

Gastrointestinal symptoms may be treated symptomatically, with attention to the water/electrolyte balance.

Amoxicillin/clavulanic acid can be removed from the circulation by haemodialysis.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic Class: Combinations of penicillins, incl. beta-lactamase inhibitors.

ATC Code: J01CR02.

Mechanism of Action

Amoxicillin is a semisynthetic penicillin (beta-lactam antibiotic) that inhibits one or more enzymes (often referred to as penicillin-binding proteins, PBPs) in the biosynthetic pathway of bacterial peptidoglycan, which is an integral structural component of the bacterial cell wall. Inhibition of peptidoglycan synthesis leads to weakening of the cell wall, which is usually followed by cell lysis and death.

Amoxicillin is susceptible to degradation by beta-lactamases produced by resistant bacteria and therefore the spectrum of activity of amoxicillin alone does not include organisms which produce these enzymes.

Clavulanic acid is a beta-lactam structurally related to penicillins. It inactivates some beta-lactamase enzymes thereby preventing inactivation of amoxicillin. Clavulanic acid alone does not exert a clinically useful antibacterial effect.

PK/PD relationship

The time above the minimum inhibitory concentration ($T > MIC$) is considered to be the major determinant of efficacy for amoxicillin.

5.2 Pharmacokinetic properties

Absorption

The pharmacokinetic results for studies in which amoxicillin/clavulanic acid was administered to groups of healthy volunteers as either 500 mg/100 mg or 1000 mg/200 mg given as a bolus

intravenous injection are presented below.

Mean ($\pm$ SD) pharmacokinetic parameters <i>Bolus intravenous injection</i>					
Dose administered	Amoxicillin				
	Dose	Mean peak serum conc (μ g/ml)	T 1/2 (h)	AUC (h.mg/l)	Urinary recovery (% 0 to 6 h)
AMX/CA mg/100 mg 500	500 mg	32.2	1.07	25.5	66.5
AMX/CA mg/200 mg 1000	1000 mg	105.4	0.9	76.3	77.4
Clavulanic acid					
AMX/CA mg/100 mg 500	100 mg	10.5	1.12	9.2	46.0
AMX/CA mg/200 mg 1000	200 mg	28.5	0.9	27.9	63.8
AMX – amoxicillin, CA – clavulanic acid					

Distribution

About 25% of total plasma clavulanic acid and 18% of total plasma amoxicillin is bound to protein. The apparent volume of distribution is around 0.3-0.4 l/kg for amoxicillin and around 0.2 l/kg for clavulanic acid.

Following intravenous administration, both amoxicillin and clavulanic acid have been found in gall bladder, abdominal tissue, skin, fat, muscle tissues, synovial and peritoneal fluids, bile and pus. Amoxicillin does not adequately distribute into the cerebrospinal fluid.

From animal studies there is no evidence for significant tissue retention of drug-derived material for either component. Amoxicillin, like most penicillins, can be detected in breast milk. Trace quantities of clavulanic acid can also be detected in breast milk.

Biotransformation

Amoxicillin is partly excreted in the urine as the inactive penicilloic acid in quantities equivalent to up to 10 to 25% of the initial dose. Clavulanic acid is extensively metabolized in man, and eliminated in urine and faeces and as carbon dioxide in expired air.

Elimination

The major route of elimination for amoxicillin is via the kidney, whereas for clavulanic acid it is by both renal and non-renal mechanisms.

Amoxicillin/clavulanic acid has a mean elimination half-life of approximately one hour and a mean total clearance of approximately 25 l/h in healthy subjects. Approximately 60 to 70% of the amoxicillin and approximately 40 to 65% of the clavulanic acid are excreted unchanged in urine during the first 6 h after administration of a single 500 mg/100 mg or a single 1000 mg/200 mg bolus intravenous injection. Various studies have found the urinary excretion to be 50-85% for amoxicillin and between 27-60% for clavulanic acid over a 24-hour period. In the case of

clavulanic acid, the largest amount of drug is excreted during the first 2 hours after administration. Concomitant use of probenecid delays amoxicillin excretion but does not delay renal excretion of clavulanic acid.

Paediatric population

The elimination half-life of amoxicillin is similar for children aged around 3 months to 2 years and older children and adults. For very young children (including preterm newborns) in the first week of life the interval of administration should not exceed twice daily administration due to immaturity of the renal pathway of elimination.

Older people

Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

Renal impairment

The total serum clearance of amoxicillin/clavulanic acid decreases proportionately with decreasing renal function. The reduction in drug clearance is more pronounced for amoxicillin than for clavulanic acid, as a higher proportion of amoxicillin is excreted *via* the renal route. Doses in renal impairment must therefore prevent undue accumulation of amoxicillin while maintaining adequate levels of clavulanic acid.

Hepatic impairment

Hepatically impaired patients should be dosed with caution and hepatic function monitored at regular intervals.

5.3 Preclinical safety data

Nonclinical data reveal no special hazard for humans based on studies of safety pharmacology, genotoxicity and toxicity to reproduction.

Repeat dose toxicity studies performed in dogs with amoxicillin/clavulanic acid demonstrate gastric irritancy and vomiting, and discoloured tongue.

Carcinogenicity studies have not been conducted with amoxicillin clavulanic acid or its components.

6. PHARMACEUTICAL PARTICULARS

6.1 List of Excipients

None

6.2 Incompatibilities

Co-Amoxiclav for Injection BP 1.2g must not be mixed with amino acid solutions, lipid emulsions, blood and glucose solutions.

6.3 Shelf Life

24 Months

6.4 Special Precautions for Storage

Store below 30°C. Protect from moisture & light.

6.5 Nature and Contents of Container

Type and nature of materials of the bottle: 20 ml clear glass vial Type-1, 20mm grey bromo butyl rubber closure Type-I and 20mm Flip-off Sterile Aluminium Seal, free from dirt, foreign matter or any other contamination both inside and outside.

Pack Size: 20ml vial + 20ml Sterilised water for Injections BP Packed in a carton along with leaflet.

6.6 Special precautions for disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Keep medicines out of the sight and reach of children.

7. MARKETING AUTHORISATION HOLDER

Syner-Med Pharmaceuticals (Kenya) Limited
Gayatri House ICD Road, Off Mombasa Road, Nairobi,
Kenya.

8. MARKETING AUTHORISATION NUMBER

H2007/224

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

23rd July, 2026

10. DATE OF REVISION OF THE TEXT

23rd July, 2026