



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

About Medicine

Prescription only medicine

1. Name of the medicinal product

Glucose 5% w/v Intravenous Infusion (D5)

2. Qualitative and quantitative composition

Each ml contains 50mg of Glucose

Approximately 840 kJ/L (or 200 kcal/L)

For the full list of excipients: see section 6.1

3. Pharmaceutical form

Solution for infusion.

A clear, colourless solution free from visible particles.

Osmolarity: 278 mOsm/L (approx.)

pH: 3.7 – 6.0

4. Clinical particulars

4.1 Therapeutic indications

- D5 is indicated for the treatment of carbohydrate and fluid depletion
- D5 is also used as a vehicle and diluent for compatible medicinal products for parenteral administration.

4.2 Posology and method of administration

Posology

Adults, older people and children:

The concentration and dosage of D5 is determined by several factors including the age, weight and clinical condition of the patient. Serum-glucose concentrations may need to be carefully monitored.

Fluid balance, serum glucose, serum sodium and other electrolytes should be monitored before and during administration, especially in patients with increased non-osmotic vasopressin release



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

(Syndrome of Inappropriate Antidiuretic Hormone Secretion, SIADH) and in patients co-medicated with vasopressin agonist drugs due to the risk of hyponatraemia. Monitoring of serum sodium is particularly important for physiologically hypotonic fluids. <Product name> may become extremely hypotonic after administration due to glucose metabolism in the body (see sections 4.4, 4.5 and 4.8).

The recommended dosage for treatment of carbohydrate and fluid depletion is:

- for adults: 500 ml to 3 litres / 24h

- for babies and children;

_0-10 kg body weight :100 ml/kg/24h.

_10-20 kg body weight :1000 ml + 50 ml /kg over 10 kg / 24h.

_0 kg body weight :1500 ml + 20 ml / kg over 20 kg / 24h.

The infusion rate depends on the patient's clinical condition.

Infusion rate should not exceed the patient's glucose oxidation capacities in order to avoid hyperglycaemia. Therefore, the maximum dose ranges from 5mg/kg/min for adults to 10-18 mg/kg/min for babies and children depending on the age and the total body mass.

The recommended dosage when used as a vehicle or diluent ranges from 50 to 250 ml per dose of medicinal product to be administered.

When D5 is used as a diluent for injectable preparations of other drugs, the dosage and the infusion rate will be principally dictated by the nature and the dose regimen of the prescribed drug.

Paediatric Population

The infusion rate and volume depend on the age, weight clinical and metabolic conditions of the patient, concomitant therapy and should be determined by the consulting physician experienced in paediatric intravenous fluid therapy.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

Method of administration:

The solution is for administration by intravenous infusion (peripheral or central vein).

When the solution is used for dilution and delivery of therapeutic additives for administration by intravenous infusion, the direction for use with additive therapeutic substances will dictate the appropriate volumes for each therapy.

D5 is an isosmotic solution.

Please see section 3 for the information about the osmolarity of the solution.

Precautions to be taken before handling or administering the medicinal product

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Use only if the solution is clear, without visible particles and the container is undamaged. Administer immediately following the insertion of infusion set.

The solution should be administered with sterile equipment using aseptic technique. The equipment should be primed with the solution in order to prevent air entering the system.

Electrolyte supplementation may be indicated according to the clinical needs of the patient. Additives may be introduced before or during infusion through the injection site.

When introducing additives, the final osmolarity of solutions need to be checked. Administration of hyperosmolar solutions may cause venous irritation and phlebitis. Thorough and careful aseptic mixing of any additive is mandatory. Solutions containing additives should be used immediately and not stored.

Please see section 4.4 for the risk of air embolism.

4.3 Contraindications

The solution is contraindicated in case of uncompensated diabetes, other known glucose intolerances (such as metabolic stress situations), hyperosmolar coma, hyperglycaemia, hyperlactataemia.

Hypersensitivity to the active substance. See sections 4.4 and 4.8 for corn allergies.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

4.4 Special warnings and precautions for use

Glucose intravenous infusions are usually isotonic solutions. In the body, however, glucose containing fluids can become extremely physiologically hypotonic due to rapid glucose metabolism (see section 4.2).

Dilution and other effects on serum electrolytes

Depending on the tonicity of the solution, the volume and rate of infusion and depending on a patient's underlying clinical condition and capability to metabolize glucose, intravenous administration of glucose can cause;

- Hyperosmolality, osmotic diuresis and dehydration
- Hypoosmolality
- Electrolyte disturbances such as;
 - hypo- or hyperosmotic hyponatraemia (see below),
 - hypokalaemia,
 - hypophosphataemia,
 - hypomagnesaemia,
 - overhydration/hypervolaemia and, for example, congested states, including pulmonary congestion and oedema.

The above effects do not only result from the administration of electrolyte-free fluid but also from glucose administration.

Hyponatraemia:

Patients with non-osmotic vasopressin release (e.g. in acute illness, pain, post-operative stress, infections, burns, and CNS diseases), patients with heart-, liver- and kidney diseases and patients exposed to vasopressin agonists (see section 4.5) are at particular risk of acute hyponatraemia upon infusion of hypotonic fluids.

Acute hyponatraemia can lead to acute hyponatraemic encephalopathy (brain oedema) characterized by headache, nausea, seizures, lethargy and vomiting. Patients with brain oedema are at particular risk of severe, irreversible and life-threatening brain injury.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

Children, women in the fertile age and patients with reduced cerebral compliance (e.g. meningitis, intracranial bleeding, and cerebral contusion) are at particular risk of the severe and life-threatening brain swelling caused by acute hyponatraemia.

Clinical evaluation and periodic laboratory determinations may be necessary to monitor changes in fluid balance, electrolyte concentrations, and acid-base balance during prolonged parenteral therapy or whenever the condition of the patient or the rate of administration warrants such evaluation.

Particular caution is advised in patients at increased risk of water and electrolyte disturbances that could be aggravated by increased free water load, hyperglycaemia or possibly required insulin administration (see below).

Hyperglycaemia

- Rapid administration of glucose solutions may produce substantial hyperglycaemia and a hyperosmolar syndrome.
- If hyperglycaemia occurs, rate of infusion should be adjusted and/or insulin administered
- If necessary, provide parenteral supplements in potassium.
- Intravenous Glucose 5% should be administered with caution in patients with, for example:
 - impaired glucose tolerance (such as in diabetes mellitus, renal failure, or in the presence of sepsis, trauma, or shock),
 - severe malnutrition (risk of precipitating a refeeding syndrome – see below),
 - thiamine deficiency, e.g., in patients with chronic alcoholism (risk of severe lactic acidosis due to impaired oxidative metabolism of pyruvate),
 - patients with ischemic stroke or severe traumatic brain injury

Avoid infusion within the first 24 hours following head trauma. Monitor blood glucose closely as early hyperglycaemia has been associated with poor outcomes in patients with severe traumatic brain injury.

- Newborns



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

Effects on Insulin Secretion

Prolonged intravenous administration of glucose and associated hyperglycaemia may result in decreased rates of glucose-stimulated insulin secretion.

Hypersensitivity Reactions

- Hypersensitivity/infusion reactions, including anaphylactic/anaphylactoid reactions, have been reported with Glucose solutions (see section 4.8). Solutions containing glucose should therefore be used with caution, if at all, in patients with known allergy to corn or corn products (see section 4.3).
- The infusion must be stopped immediately if any signs or symptoms of a suspected hypersensitivity reaction develop. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Refeeding syndrome

- Refeeding severely undernourished patients may result in the refeeding syndrome that is characterized by the shift of potassium, phosphorus, and magnesium intracellularly as the patient becomes anabolic. Thiamine deficiency and fluid retention may also develop. Careful monitoring and slowly increasing nutrient intakes while avoiding overfeeding can prevent these complications.

Paediatric population

The infusion rate and volume depend on the age, weight, clinical and metabolic conditions of the patient, concomitant therapy, and should be determined by a consulting physician experienced in paediatric intravenous fluid therapy.

In order to avoid potentially fatal over infusion of intravenous fluids to the neonate, special attention needs to be paid to the method of administration. When using a syringe pump to administer intravenous fluids or medicines to neonates, a bag of fluid should not be left connected to the syringe.

When using an infusion pump all clamps on the intravenous administration set must be closed before removing the administration set from the pump or switching the pump off. This is required regardless of whether the administration set has an anti-free flow device.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

The intravenous infusion device and administration equipment must be frequently monitored.

Paediatric glycaemia-related issues

Newborns – especially those born premature and with low birth weight - are at increased risk of developing hypo- or hyperglycaemia and therefore need close monitoring during treatment with intravenous glucose solutions to ensure adequate glycaemic control in order to avoid potential long term adverse effects. Hypoglycaemia in the newborn can cause prolonged seizures, coma and cerebral injury. Hyperglycaemia has been associated with intraventricular haemorrhage, late onset bacterial and fungal infection, retinopathy of prematurity, necrotizing enterocolitis, bronchopulmonary dysplasia, prolonged length of hospital stays, and death.

Paediatric hyponatraemia-related issues

- Children (including neonates and older children) are at increased risk of developing hypoosmotic hyponatraemia as well as for developing hyponatraemic encephalopathy.
- Plasma electrolyte concentrations should be closely monitored in the paediatric population.
- Rapid correction of hypoosmotic hyponatraemia is potentially dangerous (risk of serious neurologic complications).
- Dosage, rate, and duration of administration should be determined by a physician experienced in paediatric intravenous fluid therapy.

Geriatric Use

- When selecting the type of infusion solution and the volume/rate of infusion for a geriatric patient, consider that geriatric patients are generally more likely to have cardiac, renal, hepatic, and other diseases or concomitant drug therapy.

Blood

- Glucose 5% (an aqueous, i.e., electrolyte-free glucose solution) should not be administered simultaneously with, before or after an administration of blood through the same infusion equipment, because haemolysis and pseudo agglutination can occur.

Adding other medication or using an incorrect administration technique might cause the appearance of fever reactions due to the possible introduction of pyrogens. In case of adverse reaction, infusion must be stopped immediately.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

Risk of Air Embolism

- Do not use plastic containers in series connections. Such use could result in air embolism due to residual air being drawn from the primary container before the administration of the fluid from the secondary container is completed.
- Pressurizing intravenous solutions contained in flexible plastic containers to increase flow rates can result in air embolism if the residual air in the container is not fully evacuated prior to administration.
- Use of a vented intravenous administration set with the vent in the open position could result in air embolism. Vented intravenous administration sets with the vent in the open position should not be used with flexible plastic containers.

4.5 Interaction with Other Medicinal Products and Other Forms of Interaction

Both the glycaemic effects of D5 and its effects on water and electrolyte balance should be taken into account when using it in patients treated with other substances that affect glycaemic control, or fluid and/or electrolyte balance.

Concomitant administration of catecholamines and steroids decreases the glucose up-take.

Drugs leading to an increased vasopressin effect

The below listed drugs increase the vasopressin effect, leading to reduced renal electrolyte free water excretion and increase the risk of hospital acquired hyponatraemia following inappropriately balanced treatment with I.V. fluids (see sections 4.2, 4.4 and 4.8).

- Drugs stimulating vasopressin release, e.g.: Chlorpropamide, clofibrate, carbamazepine, vincristine, selective serotonin reuptake inhibitors, 3,4-methylenedioxy-N- methamphetamine, ifosfamide, antipsychotics, narcotics
- Drugs potentiating vasopressin action, e.g.: Chlorpropamide, NSAIDs, cyclophosphamide
- Vasopressin analogues, e.g.: Desmopressin, oxytocin, terlipressin

Other medicinal products increasing the risk of hyponatraemia also include diuretics in general and antiepileptics such as oxcarbazepine.

No interaction studies have been performed.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

4.6 Fertility, Pregnancy and Lactation

When a medicinal product is added, the nature of the drug and its use during pregnancy and lactation have to be considered separately.

Intrapartum maternal intravenous glucose infusion may result in foetal insulin production, with an associated risk of foetal hyperglycaemia and metabolic acidosis as well as rebound hypoglycaemia in the neonate.

Pregnancy

Glucose solution can be used during pregnancy. However, caution should be exercised when glucose solution is used intrapartum.

D5 should be administered with special caution for pregnant women during labour particularly if administered in combination with oxytocin due to the risk of hyponatraemia (see section 4.4, 4.5 and 4.8).

Fertility

There are no adequate data of the effect of D5 on fertility. However, no effect on fertility is expected.

Lactation

There are no adequate data of using Glucose solution during lactation. However, no effect on lactation is expected. D5 can be used during lactation.

4.7 Effects on ability to drive and use machines

No studies have been conducted on the influence of D5 on the ability to operate an automobile or other heavy machinery.

4.8 Undesirable effects

Undesirable effects which occurred in patients treated with Glucose 5% from the post-marketing experience are tabulated below.

The adverse drug reactions listed in this section are given following the recommended frequency convention: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$); and not known (cannot be estimated from the available data).



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

Tabulated list of adverse reactions

System Organ Class (SOC)	Adverse reactions (Preferred Term)	Frequency
Immune system disorders	Anaphylactic reaction* Hypersensitivity*	Not known
Metabolism and nutrition disorders	Electrolyte imbalance Hypokalaemia Hypomagnesaemia Hypophosphatemia Hyperglycaemia Dehydration Hypervolaemia Hospital acquired hyponatraemia**	Not known
Nervous system disorders	Hyponatraemic encephalopathy**	Not known
Skin and subcutaneous tissue disorders	Rash	Not known
Vascular disorders	Venous thrombosis Phlebitis	Not known
Renal and urinary disorders	Polyuria	Not known
General disorders and administration site conditions	Chills* Pyrexia* Infusion site infection Infusion site irritation for example erythema Extravasation Local reaction Pain localised	Not known

*Potential manifestation in patients with allergy to corn, see section 4.4. The following adverse reactions have not been reported with this product but may occur:

* Hospital acquired hyponatraemia may cause irreversible brain injury and death due to development of acute hyponatraemic encephalopathy (see sections 4.2 and 4.4).



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

Suspected adverse reactions can be reported via email on drugsafety@abacuspharma.com or through telephone on +256 786 557 530

4.9 Overdose

Prolonged administration or rapid infusion of large volumes of Glucose 5% may cause hyperosmolarity and hyponatraemia, dehydration, hyperglycaemia, hyperglycosuria, osmotic diuresis (due to the hyperglycaemia) and water intoxication and oedema. Severe hyperglycaemia and hyponatraemia may be fatal (see sections 4.4 and 4.8).

In case of suspected overdose, treatment with D5 must be stopped immediately. Management of overdose is symptomatic and supportive, with appropriate monitoring.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic Group – Other IV Solution Additives

ATC Code: B05BA03

The pharmacodynamic properties of this solution are those of glucose, which forms the principal source of energy in cellular metabolism. D5 is given as a source of carbohydrate in parenteral nutrition. D5 solution provides a caloric intake of 200 kcal/l. Furthermore, this glucose solution for intravenous infusion allows hydric supplementation without ionic supplementation.

D5 is an isosmotic solution, with an approximate osmolarity of 278 mOsm/l.

The pharmacodynamics of the additive will depend on the nature of the drug used.

5.2 Pharmacokinetic properties

Glucose is metabolized via pyruvic or lactic acid to carbon dioxide and water with the release of energy.

The pharmacokinetics of the additive will depend on the nature of the drug used.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

5.3 Preclinical safety data

The safety of glucose in animals is not relevant in view of its presence as a normal component in animal and human plasma.

The safety of the additive should be considered separately.

6. Pharmaceutical particulars

6.1 List of excipients

Water For Injections

6.2 Incompatibilities

As with all parenteral solutions compatibility of the additives with the solution must be assessed before addition.

It is the responsibility of the physician to judge the incompatibility of an additive medication with D5 solution by checking for eventual colour change and/or eventual precipitate, insoluble complexes or crystals apparition. The Instructions for use of the medication to be added must be consulted.

Before adding a drug, verify it is soluble and stable in water at the pH of D5.

When a compatible medication is added to the D5, the solution must be administered immediately.

Those additives known to be incompatible should not be used

6.3 Shelf life

36 months.

Use immediately after first opening.

6.4 Special precautions for storage

Do not store above 30°C and do not freeze.

6.5 Nature and contents of container

Pack sizes: 250mL and 500 mL.

The bottles are made from Low Density Polyethylene plastic; the bottles are then flowwrapped with a protective plastic pouch.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

6.6 Special precautions for disposal and other handling

For I.V. Injection

Use as directed by the physician.

Keep out of reach of children.

Do not use unless solution is clear and the container is undamaged.

Any contents of the product remaining after use should be discarded.

Do not store solution containing additives.

When introducing additives to D5 solution, aseptic techniques must be used.

Mix the solution thoroughly when additives have been introduced.

a) Opening

Remove the LDPE container from the over pouch just before use.

Check for minute leaks by squeezing inner bag firmly. If leaks are found, discard solution, as sterility may be impaired.

Check the solution for limpidity and absence of foreign matters. If solution is not clear or contains foreign matters, discard the solution.

b) Preparation for administration

Use sterile material for preparation and administration.

Suspend container from eyelet support.

Remove plastic protector from outlet port at bottom of container:

- grip the small wing on the neck of the port with one hand,
- grip the large wing on the cap with the other hand and twist,
- the cap will pop off.

Use an aseptic method to set up the infusion

Attach administration set. Refer to complete directions accompanying set for connection, priming of the set and administration of the solution.

c) Techniques for injection of additive medications

Warning: Additives may be incompatible.

To add medication before administration

- Disinfect medication site.



SUMMARY OF PRODUCT CHARACTERISTICS

GLUCOSE 5.0% w/v INTRAVENOUS INFUSION

- Using syringe with an appropriate needle, puncture resealable medication port and inject.
- Mix solution and medication thoroughly. For high-density medication such as potassium chloride, tap the ports gently while ports are upright and mix.

Caution: Do not store bags containing added medications.

To add medication during administration

- Close clamp on the set.
- Disinfect medication site.
- Using syringe with an appropriate needle, puncture resealable medication port and inject.
- Remove container from IV pole and/or turn to an upright position.
- Evacuate both ports by tapping gently while the container is in an upright position.
- Mix solution and medication thoroughly.
- Return container to in use position, re-open the clamp and continue administration.

7. Marketing authorisation holder

Abacus Parenteral Drugs Limited
Block 191, Plot No.114, Kinga Mukono
P.O. Box 31376, Kampala, Uganda.
Email: apdl@abacuspharma.com
Website: www.abacuspenteral.com

8. Marketing authorisation number

N/A

9. Date of first authorisation/renewal of the authorisation

N/A

10. Date of revision of the text

June 2025