

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

DONET-M 400SR

Doxofylline (Sustained Release) 400 mg and Montelukast 10 mg Tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each uncoated bilayered tablet contains:

Doxofylline400 mg
(As sustained release form) Montelukast Sodium BP
Equivalent Montelukast10 mg
Excipients Q.S.
Color Quinoline Yellow

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORMS

A White and Yellow Colored, Bilayered, Caplet Shape, Uncoated, Biconvex Tablet, Break Line On Yellow Side and Plain On Other Side.

4. CLINICAL PARTICULARS

4.1. Therapeutic indications

Bronchial asthma.

Pulmonary disease associated with bronchospasm.

4.2. Posology and method of administration

400 mg tablets – adults: 1 tablet two/three times daily

100 mg ampoules – adults: 2 ampoules by slow intravenous injection to patients in supine position (15-20 minutes), preferably diluted, during the acute phase.

Administration can be repeated at 12-hour intervals, at the physician's discretion.

200 mg sachets - children aged 6-12 years

1-3 sachets per day (12-18 mg/kg) dissolved in plenty of water.

2% syrup: - adults: one 20 ml measures two/three times a day (one 20 ml measure corresponds to 400 mg of doxofylline)

At the recommended posology, the plasma levels of Doxofylline do not generally exceed 20 µg/ml, so it is not essential to check these levels periodically.

If the dosage is increased, the blood levels of the drug must be measured (the therapeutic value is about 10 µg/ml, the value bordering on toxicity is 20 µg/ml).

4.3. Contraindications

DONET 400SR is contraindicated in individuals with known hypersensitivity to the drug or other xanthenes' derivatives. It is also contraindicated in patients with acute myocardial infarction, hypotension and during lactation.

4.4. Special warnings and precautions for use

Numerous factors may reduce the hepatic clearance of xanthine derivatives with increased plasma levels of the drug. These factors include age, congestive cardiac decompensation, chronic obstructive pulmonary disease, severe liver disease, concomitant infections, the concurrent administration of several drugs such as: erythromycin, TAO, lincomycin, clindamycin, allopurinol, cimetidine, influenza vaccine and propranolol. In these cases, it may prove necessary to reduce the dosage of the drug.

Phenytoin, other anticonvulsants and cigarette smoking may increase the clearance of xanthine derivatives with a reduction of plasmatic half-life. In these cases, it may prove necessary to increase the dosage of the drug.

In case of factors that may influence the clearance of xanthine derivatives, monitoring of the concentration of the blood levels of the drug is recommended for the control of the therapeutic range.

Caution should be observed in administering the product to patients with cardiac disease, hypertension, in the elderly, in patients with severe hypoxemia, hyperthyroidism, chronic cor pulmonale, congestive heart failure, liver disease, peptic ulcer and in those with renal impairment. In particular, it is to be used with caution in patients with congestive heart failure, since the clearance of the drug is considerably slower in these patients in which high blood levels may persist for long periods even after discontinuation of the treatment.

There is no risk of addiction or any other form of dependence.

Patients should be advised never to use oral montelukast to treat acute asthma attacks and to keep their usual appropriate rescue medication for this purpose readily available. If an acute attack occurs, a shortacting inhaled β -agonist should be used. Patients should seek their doctor's advice as soon as possible if they need more inhalations of short-acting β -agonists than usual. Montelukast should not be substituted abruptly for inhaled or oral corticosteroids. There are no data demonstrating that oral corticosteroids can be reduced when montelukast is given concomitantly. In rare cases, patients on therapy with anti-asthma agents including montelukast may present with systemic eosinophilia, sometimes presenting with clinical features of vasculitis consistent with Churg-Strauss syndrome, a condition which is often treated with systemic corticosteroid therapy. These cases have been sometimes associated with the reduction or withdrawal of oral corticosteroid therapy. Although a causal relationship with leukotriene receptor antagonism has not been established, physicians should be alert to eosinophilia, vasculitic rash, worsening pulmonary symptoms, cardiac complications, and/or neuropathy presenting in their patients. Patients who develop these symptoms should be reassessed and their treatment regimens evaluated. Treatment with montelukast does not alter the need for patients with aspirin-sensitive asthma to avoid taking aspirin and other non-steroidal anti-inflammatory drugs. This medicinal product contains lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucosegalactose malabsorption should not take this medicine.

Neuropsychiatric events have been reported in adults, adolescents, and children taking Montelukast 10 mg film-coated tablets (see section 4.8). Patients and physicians should be alert for neuropsychiatric events. Patients and/or caregivers should be instructed to notify their physician if these changes occur. Prescribers should carefully evaluate the risks and benefits of continuing treatment with Montelukast 10 mg film-coated tablets if such events occur.

KEEP OUT OF THE REACH AND SIGHT OF CHILDREN

4.5. Interaction with other medicinal products and other forms of interaction

DONET 400SR should not be administered with other xanthine preparations. It is recommended to limit consumption of beverages and food containing caffeine.

Caution should be exercised in administering DONET 400SR together with ephedrine or other sympathomimetic drugs.

The concurrent administration of many drugs such as erythromycin, TAO, lincomycin, clindamycin, allopurinol, cimetidine, influenza vaccine and propranolol may reduce the hepatic clearance of xanthine derivatives with an increase in the plasmatic levels of the drug.

Phenytoin, other anticonvulsants and cigarette smoking may increase the clearance of xanthine derivatives with a reduction of plasmatic half-life. In these cases, it may prove necessary to increase the dosage of the drug.

Montelukast may be administered with other therapies routinely used in the prophylaxis and chronic treatment of asthma. In drug-interactions studies, the recommended clinical dose of montelukast did not have clinically important effects on the pharmacokinetics of the following medicinal products: theophylline, prednisone, prednisolone, oral contraceptives (ethinyl estradiol/norethindrone 35/1), terfenadine, digoxin and warfarin. The area under the plasma concentration curve (AUC) for montelukast was decreased approximately 40% in subjects with co-administration of phenobarbital. Since montelukast is metabolised by CYP 3A4, 2C8, and 2C9, caution should be exercised, particularly in children, when montelukast is co-administered with inducers of CYP 3A4, 2C8, and 2C9, such as phenytoin, phenobarbital and rifampicin. In vitro studies have shown that montelukast is a potent inhibitor of CYP 2C8. However, data from a clinical drug-drug interaction study involving montelukast and rosiglitazone (a probe substrate representative of medicinal products primarily metabolized by CYP 2C8) demonstrated that montelukast does not inhibit CYP 2C8 in vivo. Therefore, montelukast is not anticipated to markedly alter the metabolism of medicinal products metabolised by this enzyme (e.g., paclitaxel, rosiglitazone, and repaglinide.) In vitro studies have shown that montelukast is a substrate of CYP 2C8, and to a less significant extent, of 2C9, and 3A4. In a clinical drug-drug interaction study involving montelukast and gemfibrozil (an inhibitor of both CYP 2C8 and 2C9) gemfibrozil increased the systemic exposure of montelukast by 4.4-fold. No routine dosage adjustment of montelukast is required upon coadministration with gemfibrozil or other potent inhibitors of CYP 2C8, but the physician should be aware of the potential for an increase in adverse reactions. Based on in vitro data, clinically important drug interactions with less potent inhibitors of CYP 2C8 (e.g., trimethoprim) are not anticipated. Co-administration of montelukast with itraconazole, a strong inhibitor of CYP 3A4, resulted in no significant increase in the systemic exposure of montelukast.

4.6. Pregnancy and lactation

Animal tests have shown that the active ingredient of DONET 400SR does not interfere with pre- and postnatal growth.

However, as there is not sufficient clinical evidence about the effects of the drug during pregnancy, use of the drug during pregnancy should be evaluated carefully case by case on the basis of the risk- benefit ratio. The drug is contraindicated during lactation.

Available data from published prospective and retrospective cohort studies with montelukast use in pregnant women evaluating major birth defects have not established a drug-associated risk. Available studies have methodologic limitations, including small sample size, in some cases retrospective data collection, and inconsistent comparator groups. Montelukast 10 mg film-coated tablets may be used during pregnancy only if it is considered to be clearly essential.

Breastfeeding It is unknown whether montelukast is excreted in human milk. Studies in rats have shown that montelukast is excreted in milk (see section 5.3). Montelukast 10 mg film-coated tablets may be used in breast-feeding only if it is considered to be clearly essential

4.7. Effects on ability to drive and use machines

The product does not affect the patient's alertness and therefore does not interfere with his/her ability to drive and use machines.

Montelukast has no or negligible influence on the ability to drive and use machines. However, individuals have reported drowsiness or dizziness.

4.8. Undesirable effects

Patients treated with xanthine derivatives may suffer nausea, vomiting, epigastric pain, headache, irritability, insomnia, tachycardia, extrasystoles, tachypnea, and in rare cases, hyperglycemia or albuminuria. In case of overdose severe cardiac arrhythmias and tonic-clonic seizure may occur. These effects may represent the first signs of intoxication.

The appearance of side effects may require discontinuation of the treatment which, if necessary, at the physician's discretion, may be resumed at lower doses after all signs and symptoms of toxicity have subsided.

Montelukast has been evaluated in clinical studies as follows:

- 10 mg film-coated tablets in approximately 4000 adult and adolescent asthmatic patients 15 years of age and older.

- 10 mg film-coated tablets in approximately 400 adult and adolescent asthmatic patients with seasonal allergic rhinitis 15 years of age and older.
- 5 mg chewable tablets in approximately 1750 pediatric asthmatic patients 6 to 14 years of age.

The following drug-related adverse reactions in clinical studies were reported commonly (1/100 to <1/10) in asthmatic patients treated with montelukast and at a greater incidence than in patients treated with placebo.

Body system Class	Adult and Adolescent Patients 15 years and older (two 12-week studies; n=795)	Paediatric Patients 6 to 14 years old (one 8-week study; n=201) (two 56-week studies; n=615)
Nervous system disorders	headache	headache
Gastrointestinal disorders	abdominal pain	

With prolonged treatment in clinical trials with a limited number of patients for up to 2 years for adults, and up to 12 months for paediatric patients 6 to 14 years of age, the safety profile did not change.

Tabulated list of Adverse Reactions

Adverse reactions reported in post-marketing use are listed, by System Organ Class and specific Adverse Experience Term, in the table below. Frequency Categories were estimated based on relevant clinical trials.

System Class	Organ	Adverse Reactions	Frequency Category*
Infections and infestations	an	upper respiratory infection [†]	Very Common
Blood disorders	an	increased bleeding tendency	Rare
Immunological disorders	an	thrombocytopenia	Very Rare
Immune disorder	system	hypersensitivity reactions including anaphylaxis	Uncommon
		hepatic eosinophilic infiltration	Very Rare
Psychiatric disorders		dream abnormalities including nightmares, insomnia, somnambulism, anxiety, agitation including aggressive behaviour or hostility, depression, psychomotor hyperactivity (including irritability, restlessness, tremor [§])	Uncommon

		disturbance in attention, memory impairment , tic	Rare
		hallucinations, disorientation, suicidal thinking and behaviour (suicidality), obsessive- compulsive symptoms, dysphemia	Very Rare
Nervous system disorder		dizziness, drowsiness paraesthesia/hypoesthesia, seizure	Uncommon
Cardiac disorders		palpitations	Rare
Respiratory, thoracic and mediastinal disorders		epistaxis	Uncommon
		Churg-Strauss Syndrome (CSS) (see section 4.4), pulmonary eosinophilia	Very Rare
Gastrointestinal disorders		diarrhoea [‡] , nausea [‡] , vomiting [‡]	Common
		dry mouth, dyspepsia	Uncommon
Hepatobiliary disorders		elevated levels of serum transaminases (ALT, AST)	Common
		hepatitis (including cholestatic, hepatocellular, and mixed-pattern liver injury).	Very Rare
Skin and subcutaneous tissue disorders		rash [‡]	Common
		bruising, urticaria, pruritus	Uncommon
		angiooedema	Rare
		erythema nodosum, erythema multiforme	Very Rare
Musculoskeletal and connective tissue disorders		arthralgia, myalgia including muscle cramps	Uncommon
Renal and urinary disorders		enuresis in children	Uncommon
General disorders and administration site conditions		pyrexia [‡]	Common
		asthenia/fatigue, malaise, oedema,	Uncommon

*Frequency Category: Defined for each Adverse Reactions by the incidence reported in the clinical trials data base: Very Common ($\geq 1/10$), Common ($\geq 1/100$ to $< 1/10$), Uncommon ($\geq 1/1000$ to $< 1/100$), Rare ($\geq 1/10,000$ to $< 1/1000$), Very Rare ($< 1/10,000$).

[†]This adverse experience, reported as Very Common in the patients who received montelukast, was also reported as Very Common in the patients who received placebo in clinical trials.

[‡]This adverse experience, reported as Common in the patients who received montelukast, was also reported as Common in the patients who received placebo in clinical trials.

§ Frequency Category: Rare

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Reporting of suspected adverse reactions: Healthcare professionals are requested to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org>

4.9. Overdose

As there is no specific antidote, in case of overdose a symptomatic treatment of cardiovascular collapse should be instituted.

In chronic asthma studies, montelukast has been administered at doses up to 200 mg/day to adult patients for 22 weeks and in short term studies, up to 900 mg/day to patients for approximately one week without clinically important adverse experiences. There have been reports of acute overdose in post-marketing experience and clinical studies with montelukast. These include reports in adults and children with a dose as high as 1000 mg (approximately 61 mg/kg in a 42-month-old child). The clinical and laboratory findings observed were consistent with the safety profile in adults and pediatric patients.

Symptoms of overdose

There were no adverse experiences in the majority of overdose reports. The most frequently occurring adverse experiences were consistent with the safety profile of montelukast and included abdominal pain, somnolence, thirst, headache, vomiting and psychomotor hyperactivity.

Management of overdose

No specific information is available on the treatment of overdose with montelukast. It is not known whether montelukast is dialysable by peritoneal- or haemo-dialysis.

5. PHARMACOLOGICAL PROPERTIES

5.1. Pharmacodynamic properties

Doxofylline directly relaxes the smooth muscles of the bronchi and pulmonary vessels. In this way, it acts mainly as a bronchodilator, pulmonary vasodilator and as a relaxant of the bronchial muscle. The action of Doxofylline may be mediated, at least in part, by inhibition of the phosphodiesterase leading to an increase in intracellular cyclic AMP which results in smooth-muscle relaxation. At higher concentrations, Doxofylline may inhibit the release of histamine by the cells. Prolonged use of the drug does not lead to addiction.

Pharmacotherapeutic group: Leukotriene receptor antagonist

ATC code: R03D03

Mechanism of action

The cysteinyl leukotrienes (LTC₄, LTD₄, and LTE₄) are potent inflammatory eicosanoids released from various cells including mast cells and eosinophils. These important pro-asthmatic mediators bind to cysteinyl leukotriene (CysLT) receptors. The CysLT type-1 (CysLT₁) receptor is found in the human airway (including airway smooth muscle cells and airway macrophages) and on other pro-inflammatory cells (including eosinophils and certain myeloid stem cells). CysLTs have been correlated with the pathophysiology of asthma and allergic rhinitis. In asthma, leukotriene-mediated effects include bronchoconstriction, mucous secretion, vascular permeability, and eosinophil recruitment. In allergic rhinitis, CysLTs are released from the nasal mucosa after allergen exposure during both early- and late-phase reactions and are associated with symptoms of allergic rhinitis. Intranasal challenge with CysLTs has been shown to increase nasal airway resistance and symptoms of nasal obstruction.

Pharmacodynamic effects

Montelukast is an orally active compound which binds with high affinity and selectivity to the CysLT₁ receptor. In clinical studies, montelukast inhibits bronchoconstriction due to inhaled LTD₄ at doses as low as 5 mg. Bronchodilation was observed within 2 hours of oral administration. The bronchodilation effect caused by a β agonist was additive to that caused by montelukast. Treatment with montelukast inhibited both early- and late phase bronchoconstriction due to antigen challenge. Montelukast, compared with placebo, decreased peripheral blood eosinophils in adult and paediatric patients. In a separate study, treatment with montelukast significantly decreased eosinophils in the airways (as measured in sputum) and in peripheral blood while improving clinical asthma control.

Clinical efficacy and safety

In studies in adults, montelukast, 10 mg once daily, compared with placebo, demonstrated significant improvements in morning FEV₁ (10.4% vs 2.7% change from baseline), AM peak expiratory flow rate (PEFR) (24.5 L/min vs 3.3 L/min change from baseline), and significant decrease in total β agonist use (-26.1% vs -4.6% change from baseline). Improvement in patient-reported daytime and nighttime asthma symptoms scores was significantly better than placebo.

Studies in adults demonstrated the ability of montelukast to add to the clinical

effect of inhaled corticosteroid (% change from baseline for inhaled beclometasone plus montelukast vs beclometasone, respectively for FEV1: 5.43% vs 1.04%; β -agonist use: -8.70% vs 2.64%). Compared with inhaled beclometasone (200 μ g twice daily with a spacer device), montelukast demonstrated a more rapid initial response, although over the 12-week study, beclometasone provided a greater average treatment effect (% change from baseline for montelukast vs beclometasone, respectively for FEV1: 7.49% vs 13.3%; β agonist use: -28.28% vs -43.89%). However, compared with beclometasone, a high percentage of patients treated with montelukast achieved similar clinical responses (e.g., 50% of patients treated with beclometasone achieved an improvement in FEV1 of approximately 11% or more over baseline while approximately 42% of patients treated with montelukast achieved the same response).

A clinical study was conducted to evaluate montelukast for the symptomatic treatment of seasonal allergic rhinitis in adult and adolescent asthmatic patients 15 years of age and older with concomitant seasonal allergic rhinitis. In this study, montelukast 10 mg tablets administered once daily demonstrated a statistically significant improvement in the Daily Rhinitis Symptoms score, compared with placebo. The Daily Rhinitis Symptoms score is the average of the Daytime Nasal Symptoms score (mean of nasal congestion, rhinorrhea, sneezing, nasal itching) and the Nighttime Symptoms score (mean of nasal congestion upon awakening, difficulty going to sleep, and nighttime awakenings scores). Global evaluations of allergic rhinitis by patients and physicians were significantly improved, compared with placebo. The evaluation of asthma efficacy was not a primary objective in this study.

In an 8-week study in paediatric patients 6 to 14 years of age, montelukast 5 mg once daily, compared with placebo, significantly improved respiratory function (FEV1 8.71% vs 4.16% change from baseline; AM PEF 27.9 L/min vs 17.8 L/min change from baseline) and decreased "as-needed" β agonist use (-11.7% vs +8.2% change from baseline).

Significant reduction of exercise-induced bronchoconstriction (EIB) was demonstrated in a 12-week study in adults (maximal fall in FEV1 22.33% for montelukast vs 32.40% for placebo; time to recovery to within 5% of baseline FEV1 44.22 min vs 60.64 min). This effect was consistent throughout the 12-week study period. Reduction in EIB was also demonstrated in a short-term study in paediatric patients (maximal fall in FEV1 18.27% vs 26.11%; time to recovery to within 5% of baseline FEV1 17.76 min vs 27.98 min). The effect in both studies was demonstrated at the end of the once-daily dosing interval.

In aspirin-sensitive asthmatic patients receiving concomitant inhaled and/or oral corticosteroids, treatment with montelukast, compared with placebo, resulted in significant improvement in asthma control (FEV1 8.55% vs -1.74% change from baseline and decrease in total β -agonist use -27.78% vs 2.09% change from baseline).

5.2. Pharmacokinetic properties

Doxofylline half-life is more than 6 hours, so constant effective plasma levels may be maintained with three administrations a day.

The kinetics after a single i.v. and oral administration have been studied in man to define the distribution and absorption of the drug.

After intravenous administration of 100 mg of Doxofylline to 5 volunteers, the distribution of the unchanged substance in the serum follows a bi-compartmental model.

The area under the curve of the concentration of the drug in the serum during the distribution phase represents a small fraction of the total area.

Plasmatic clearance is high, with values ranging from 444 to 806 ml/min, whereas

distribution volume is about 1 l/kg.

The mean half-life after intravenous administration was calculated to be 65 minutes (from 40 to 96).

After tablet administration, peak plasma levels are reached after 60 minutes, while with the syrup, due to its water and alcohol vehicle, the drug is absorbed more quickly, peak plasma concentration occurring within 30 minutes.

Absolute oral bioavailability is about 62.6%; at pH 7.4 the percentage of product bonded to the plasma proteins is about 48%.

Less than 4% of the oral dose is eliminated unchanged in the urine.

Absorption

Montelukast is rapidly absorbed following oral administration. For the 10 mg film-coated tablet, the mean peak plasma concentration (C_{max}) is achieved 3 hours (T_{max}) after administration in adults in the fasted state. The mean oral bioavailability is 64%. The oral bioavailability and C_{max} are not influenced by a standard meal. Safety and efficacy were demonstrated in clinical trials where the 10 mg film-coated tablet was administered without regard to the timing of food ingestion. For the 5 mg chewable tablet, the C_{max} is achieved in 2 hours after administration in adults in the fasted state. The mean oral bioavailability is 73% and is decreased to 63% by a standard meal.

Distribution

Montelukast is more than 99% bound to plasma proteins. The steady-state volume of distribution of montelukast averages 8-11 litres. Studies in rats with radiolabelled montelukast indicate minimal distribution across the blood-brain barrier. In addition, concentrations of radiolabelled material at 24 hours post-dose were minimal in all other tissues.

Biotransformation

Montelukast is extensively metabolised. In studies with therapeutic doses, plasma concentrations of metabolites of montelukast are undetectable at steady state in adults and children.

Cytochrome P450 2C8 is the major enzyme in the metabolism of montelukast. Additionally CYP 3A4 and 2C9 may have a minor contribution, although itraconazole, an inhibitor of CYP 3A4, was shown not to change pharmacokinetic variables of montelukast in healthy subjects that received 10 mg montelukast daily. Based on in vitro results in human liver microsomes, therapeutic plasma concentrations of montelukast do not inhibit cytochromes P450 3A4, 2C9, 1A2, 2A6, 2C19, or 2D6. The contribution of metabolites to the therapeutic effect of montelukast is minimal.

Elimination

The plasma clearance of montelukast averages 45 ml/min in healthy adults. Following an oral dose of radiolabelled montelukast, 86% of the radioactivity was recovered in 5-day faecal collections and < 0.2% was recovered in urine. Coupled with estimates of montelukast oral bioavailability, this indicates that montelukast and its metabolites are excreted almost exclusively via the bile.

Characteristics in patients

No dosage adjustment is necessary for the elderly or mild to moderate hepatic insufficiency. Studies in patients with renal impairment have not been undertaken. Because montelukast and its metabolites are eliminated by the biliary route, no dose adjustment is anticipated to be necessary in patients with renal impairment. There are no data on the pharmacokinetics of montelukast in patients with severe hepatic insufficiency (Child-Pugh score >9). With high doses of montelukast (20- and 60-fold the recommended adult dose), decrease in plasma theophylline concentration was observed. This effect was not seen at the recommended dose of 10 mg once daily.

Preclinical safety data

In animal toxicity studies, minor serum biochemical alterations in ALT, glucose, phosphorus and triglycerides were observed which were transient in nature. The signs of toxicity in animals were increased excretion of saliva, gastro- intestinal symptoms, loose stools and ion imbalance. These occurred at dosages which provided >17-fold the systemic exposure seen at the clinical dosage. In monkeys, the adverse effects appeared at doses from 150 mg/kg/day (>232-fold the systemic exposure seen at the clinical dose). In animal studies, montelukast did not affect fertility or reproductive performance at systemic exposure exceeding the clinical systemic exposure by greater than 24-fold. A slight decrease in pup body weight was noted in the female fertility study in rats at 200 mg/kg/day (>69fold the clinical systemic exposure). In studies in rabbits, a higher incidence of incomplete ossification, compared with concurrent control animals, was seen at systemic exposure >24-fold the clinical systemic exposure seen at the clinical dose. No abnormalities were seen in rats. Montelukast has been shown to cross the placental barrier and is excreted in breast milk of animals.

No deaths occurred following a single oral administration of montelukast sodium at doses up to 5000 mg/kg in mice and rats (15,000 mg/m² and 30,000 mg/m² in mice and rats, respectively), the maximum dose tested. This dose is equivalent to 25,000 times the recommended daily adult human dose (based on an adult patient weight of 50 kg).

Montelukast was determined not to be phototoxic in mice for UVA, UVB or visible light spectra at doses up to 500 mg/kg/day (approximately >200-fold based on systemic exposure).

Montelukast was neither mutagenic in in vitro and in vivo tests nor tumorigenic in rodent species.

6) PHARMACEUTICAL PARTICULARS

6.1. List of Excipients

Microcrystalline Cellulose

Hydroxypropyl methylcellulose

Polyvinyl pyrrolidone K30

Purified water PEG 6000

Purified Talc

Magnesium Stearate

Croscarmellose sodium

Crospovidone XL 10

Lake Quinoline Yellow

Colloidal silicon Dioxide

Magnesium Stearate

H.P.M.C. K 100 M

Lactose

6.2. Incompatibilities

No incompatibility with other substances has been reported for any of the available pharmaceutical forms.

6.3. Shelf Life

24 Months

6.4 Special precautions for storage

Store in the original blister in order to protect from moisture.

6.5 Nature and contents of container

Alu-Alu Blister

6.6 Special precautions for disposal

Special precautions for storage

The preparation must be stored "at ordinary environmental conditions" as laid down by. DONET 400SR

7.MARKETING AUTHORIZATION HOLDER

4Care life Science (P) Limited
SurveyNo.23/3P&24, Opp. Jeans
Factory, Daduram Vistar, Village-
Bagdol, Tal-Kathlal, Dist- Kheda -
387630, Gujarat, India.

8. MARKETING AUTHORIZATION NUMBER(S)

H2022/CTD9859/22531

9. DATE OF FIRST AUTHORIZATION

2023 August 04

10. DATE OF REVISION OF THE TEXT

2023 August 04