

## **SUMMARY OF PRODUCT CHARACTERISTICS**

### **1. NAME OF THE MEDICINAL PRODUCT**

Docetaxel 120 mg/3mL, Concentrate and Solvent for Solution for Infusion.

### **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

#### For Concentrate :

Each single-dose vial of Docetaxel 120mg/3mL concentrate contains docetaxel (as trihydrate) corresponding to 120 mg of Docetaxel (anhydrous). The viscous solution contains 40 mg/mL Docetaxel (anhydrous).

#### Excipients with known effect:

Citric Acid

Anhydrous

Polysorbate 80 For

#### Solvent :

Each single-dose vial of solvent contains 13% (w/v) ethanol 95% v/v (Absolute Alcohol content USP 15.25% v/v) in water for injection q.s. to 9mL.

#### Excipients with known effect:

Absolute Alcohol

Water for Injection

For the full list of excipients, see section 6.1.

### **3. PHARMACEUTICAL FORM**

Concentrate and Solvent for solution for infusion.

The concentrate is a pale yellow or brownish yellow, viscous solution.

The solvent is a colorless solution.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

#### Breast Cancer

Docetaxel in combination with doxorubicin and cyclophosphamide is indicated for the adjuvant treatment of patients with:

- operable node-positive breast cancer
- operable node-negative breast cancer

For patients with operable node-negative breast cancer, adjuvant treatment should be restricted to patients eligible to receive chemotherapy according to internationally established criteria for primary therapy of early breast cancer (see section 5.1).

Docetaxel in combination with doxorubicin is indicated for the treatment of patients with locally advanced or metastatic breast cancer who have not previously received cytotoxic therapy for this condition.

Docetaxel monotherapy is indicated for the treatment of patients with locally advanced or metastatic breast cancer after failure of cytotoxic therapy. Previous chemotherapy should have included an anthracycline or an alkylating agent.

Docetaxel in combination with trastuzumab is indicated for the treatment of patients with metastatic breast cancer whose tumours over express HER2 and who previously have not received chemotherapy for metastatic disease.

Docetaxel in combination with capecitabine is indicated for the treatment of patients with locally advanced or metastatic breast cancer after failure of cytotoxic chemotherapy. Previous therapy should have included an anthracycline.

#### Non-small cell lung cancer

Docetaxel is indicated for the treatment of patients with locally advanced or metastatic non-small cell lung cancer after failure of prior chemotherapy.

Docetaxel in combination with cisplatin is indicated for the treatment of patients with unresectable, locally advanced or metastatic non-small cell lung cancer, in patients who have not previously received chemotherapy for this condition.

#### Prostate cancer

Docetaxel in combination with prednisone or prednisolone is indicated for the treatment of patients with hormone refractory metastatic prostate cancer.

Docetaxel in combination with androgen-deprivation therapy (ADT), with or without prednisone or prednisolone, is indicated for the treatment of patients with metastatic hormone-sensitive prostate cancer.

#### Gastric adenocarcinoma

Docetaxel in combination with cisplatin and 5-fluorouracil is indicated for the treatment of patients with metastatic gastric adenocarcinoma, including adenocarcinoma of the gastro esophageal junction, who have not received prior chemotherapy for metastatic disease.

#### Head and neck cancer

Docetaxel in combination with cisplatin and 5-fluorouracil is indicated for the treatment of patients with locally advanced squamous cell carcinoma of the head and neck.

### **4.2 Posology and method of administration**

The use of docetaxel should be confined to units specialised in the administration of cytotoxic chemotherapy and it should be administered under the supervision of a physician qualified in the use of anticancer chemotherapy (see section 6.6).

#### Posology :

For breast, non-small cell lung, gastric, and head and neck cancers, premedication consisting of an oral corticosteroid, such as dexamethasone 16 mg per day (e.g. 8mg BID) for 3 days starting 1 day prior to docetaxel administration, unless contraindicated, can be used (see section 4.4).

For metastatic castration-resistant prostate cancer, given the concurrent use of prednisone or prednisolone the recommended premedication regimen is oral dexamethasone 8 mg, 12 hours, 3 hours and 1 hour before the docetaxel infusion (see section 4.4).

For metastatic hormone-sensitive prostate cancer, irrespective of the concurrent use of prednisone or prednisolone, the recommended premedication regimen is oral dexamethasone 8 mg 12 hours, 3 hours, and 1 hour before docetaxel infusion (see section 4.4).

Prophylactic G-CSF may be used to mitigate the risk of haematological toxicities.

Docetaxel is administered as a one-hour infusion every three weeks.

#### Breast cancer

In the adjuvant treatment of operable node-positive and node-negative breast cancer, the recommended dose of docetaxel is 75 mg/m<sup>2</sup> administered 1-hour after doxorubicin 50 mg/m<sup>2</sup> and cyclophosphamide 500 mg/m<sup>2</sup> every 3 weeks for 6 cycles. For the treatment of patients with locally advanced or metastatic breast cancer, the recommended dosage of docetaxel is 100 mg/m<sup>2</sup> in monotherapy. In firstline treatment, docetaxel 75 mg/m<sup>2</sup> is given in combination therapy with doxorubicin (50 mg/m<sup>2</sup>).

In combination with trastuzumab the recommended dose of docetaxel is 100 mg/m<sup>2</sup> every three weeks, with trastuzumab administered weekly. In a pivotal trial the initial docetaxel infusion was started the day following the first dose of trastuzumab. The subsequent docetaxel doses were administered immediately after completion of the trastuzumab infusion, if the preceding dose of trastuzumab was well tolerated. For trastuzumab dosage and administration, see trastuzumab summary of product characteristics.

In combination with capecitabine, the recommended dose of docetaxel is 75 mg/m<sup>2</sup> every three weeks, combined with capecitabine at 1250 mg/m<sup>2</sup> twice daily (within 30 minutes after a meal) for 2 weeks followed by 1-week rest period. For capecitabine dose calculation according to body surface area, see capecitabine summary of product characteristics.

### Non-small cell lung cancer

In chemotherapy naive patients treated for non-small cell lung cancer, the recommended dose regimen is docetaxel 75 mg/m<sup>2</sup> immediately followed by cisplatin 75 mg/m<sup>2</sup> over 30-60 minutes. For treatment after failure of prior platinum based chemotherapy, the recommended dosage is 75 mg/m<sup>2</sup> as a single agent.

### Prostate cancer

#### *Metastatic castration-resistant prostate cancer*

The recommended dose of docetaxel is 75 mg/m<sup>2</sup>. Prednisone or prednisolone 5 mg orally twice daily is administered continuously (see section 5.1).

#### *Metastatic hormone-sensitive prostate cancer*

The recommended dose of docetaxel is 75 mg/m<sup>2</sup> every 3 weeks for 6 cycles. Prednisone or prednisolone 5 mg orally twice daily may be administered continuously.

### Gastric adenocarcinoma

The recommended dose of docetaxel is 75 mg/m<sup>2</sup> as a 1 hour infusion, followed by cisplatin 75 mg/m<sup>2</sup>, as a 1 to 3 hour infusion (both on day 1 only), followed by 5-fluorouracil 750 mg/m<sup>2</sup> per day given as a 24-hour continuous infusion for 5 days, starting at the end of the cisplatin infusion. Treatment is repeated every three weeks. Patients must receive premedication with antiemetics and appropriate hydration for cisplatin administration. Prophylactic G-CSF should be used to mitigate the risk of haematological toxicities (see also Dose adjustments during treatment).

### Head and neck cancer

Patients must receive premedication with antiemetics and appropriate hydration (prior to and after cisplatin administration). Prophylactic G-CSF may be used to mitigate the risk of hematological toxicities. All patients on the docetaxel-containing arm of the TAX 323 and TAX 324 studies received prophylactic antibiotics.

- Induction chemotherapy followed by radiotherapy (TAX 323)

For the induction treatment of inoperable locally advanced squamous cell carcinoma of the head and neck (SCCHN), the recommended dose of docetaxel

is 75 mg/m<sup>2</sup> as a 1 hour infusion followed by cisplatin 75 mg/m<sup>2</sup> over 1 hour, on day one, followed by 5-fluorouracil as a continuous infusion at 750 mg/m<sup>2</sup> per day for five days. This regimen is administered every 3 weeks for 4 cycles. Following chemotherapy, patients should receive radiotherapy.

- Induction chemotherapy followed by chemoradiotherapy (TAX 324)

For the induction treatment of patients with locally advanced (technically unresectable, low probability of surgical cure, and aiming at organ preservation) squamous cell carcinoma of the head and neck (SCCHN), the recommended dose of docetaxel is 75 mg/m<sup>2</sup> as a 1 hour intravenous infusion on day 1,

followed by cisplatin 100 mg/m<sup>2</sup> administered as a 30-minute to 3 hour infusion, followed by 5-fluorouracil 1000 mg/m<sup>2</sup>/day as a continuous infusion from day 1 to day 4. This regimen is administered every 3 weeks for 3 cycles. Following chemotherapy, patients should receive chemoradiotherapy.

For cisplatin and 5-fluorouracil dose modifications, see the corresponding summary of product characteristics.

### Dosage adjustments during treatment

#### General

Docetaxel should be administered when the neutrophil count is  $\geq 1,500$  cells/mm<sup>3</sup>. In patients who experienced either febrile neutropenia, neutrophil  $< 500$  cells/mm<sup>3</sup> for more than one week, severe or cumulative cutaneous reactions or severe peripheral neuropathy during docetaxel therapy, the dose of docetaxel should be reduced from 100 mg/m<sup>2</sup> to 75 mg/m<sup>2</sup> and/or 75 mg/m<sup>2</sup> to 60 mg/m<sup>2</sup>. If the patient continues to experience these reactions at 60 mg/m<sup>2</sup>, the treatment should be discontinued.

#### Adjuvant therapy for breast cancer

Primary G-CSF prophylaxis should be considered in patients who receive docetaxel, doxorubicin and cyclophosphamide (TAC) adjuvant therapy for breast cancer. Patients who experience febrile neutropenia and/or neutropenic infection should have their docetaxel dose reduced to 60 mg/m<sup>2</sup> in all subsequent cycles (see sections 4.4 and 4.8). Patients who experience Grade 3 or 4 stomatitis should have their dose decreased to 60 mg/m<sup>2</sup>.

#### *In combination with cisplatin*

For patients who are dosed initially at docetaxel 75 mg/m<sup>2</sup> in combination with cisplatin and whose nadir of platelet count during the previous course of therapy is <25000 cells/mm<sup>3</sup>, or in patients who experience febrile neutropenia, or in patients with serious non-hematologic toxicities, the docetaxel dosage in subsequent cycles should be reduced to 65 mg/m<sup>2</sup>. For cisplatin dosage adjustments, see the corresponding summary of product characteristics.

#### *In combination with capecitabine*

- For capecitabine dose modifications, see capecitabine summary of product characteristics.
- For patients developing the first appearance of a Grade 2 toxicity, which persists at the time of the next docetaxel / capecitabine treatment, delay treatment until resolved to Grade 0- 1, and resume at 100% of the original dose.
- For patients developing the second appearance of a Grade 2 toxicity, or the first appearance of a Grade 3 toxicity, at any time during the treatment cycle, delay treatment until resolved to Grade 0- 1, then resume treatment with docetaxel 55 mg/m<sup>2</sup>.
- For any subsequent appearances of toxicities, or any Grade 4 toxicities, discontinue the docetaxel dose.

For trastuzumab dose modifications, see trastuzumab summary of product characteristics.

#### *In combination with cisplatin and 5-fluorouracil:*

If an episode of febrile neutropenia, prolonged neutropenia or neutropenic infection occurs despite GCSF use, the docetaxel dose should be reduced from 75 to 60 mg/m<sup>2</sup>. If subsequent episodes of complicated neutropenia occur the docetaxel dose should be reduced from 60 to 45 mg/m<sup>2</sup>. In case of Grade 4 thrombocytopenia the docetaxel dose should be reduced from 75 to 60 mg/m<sup>2</sup>. Patients should not be retreated with subsequent cycles of docetaxel until neutrophils recover to a level > 1,500 cells/mm<sup>3</sup> and platelets recover to a level > 100,000 cells/mm<sup>3</sup>. Discontinue treatment if these toxicities persist (see section 4.4).

Recommended dose modifications for gastrointestinal toxicities in patients treated with docetaxel in combination with cisplatin and 5-fluorouracil (5-FU):

| <b>Toxicity</b>              | <b>Dosage adjustment</b>                                                                                                                             |
|------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| Diarrhoea grade 3            | First episode: reduce 5-FU dose by 20%.<br>Second episode: then reduce docetaxel dose by 20%.                                                        |
| Diarrhoea grade 4            | First episode: reduce docetaxel and 5-FU doses by 20%. Second episode: discontinue treatment.                                                        |
| Stomatitis/mucositis grade 3 | First episode: reduce 5-FU dose by 20%.<br>Second episode: stop 5-FU only, at all subsequent cycles.<br>Third episode: reduce docetaxel dose by 20%. |
| Stomatitis/mucositis grade 4 | First episode: stop 5-FU only, at all subsequent cycles. Second episode: reduce docetaxel dose by 20%.                                               |

For cisplatin and 5-fluorouracil dose adjustments, see the corresponding summary of product characteristics.

In the pivotal SCCHN studies patients who experienced complicated neutropenia (including prolonged neutropenia, febrile neutropenia, or infection), it was recommended to use G-CSF to provide prophylactic coverage (eg, day 6-15) in all subsequent cycles.

#### Special populations:

##### *Patients with hepatic impairment:*

Based on pharmacokinetic data with docetaxel at 100 mg/m<sup>2</sup> as single agent, patients who have both elevations of transaminase (ALT and/or AST) greater than 1.5 times the upper limit of the normal range (ULN) and alkaline phosphatase greater than 2.5 times the ULN, the recommended dose of docetaxel is 75 mg/m<sup>2</sup> (see sections 4.4 and 5.2). For those patients with serum bilirubin > ULN and/or ALT and AST > 3.5 times the ULN associated with alkaline phosphatase > 6 times the ULN, no dose-reduction can be recommended and docetaxel should not be used unless strictly indicated.

In combination with cisplatin and 5-fluorouracil for the treatment of patients with gastric adenocarcinoma, the pivotal clinical study excluded patients with ALT and/or AST > 1.5 × ULN associated with alkaline phosphatase > 2.5 × ULN, and bilirubin > 1 × ULN; for these patients, no dose-reductions can be

recommended and docetaxel should not be used unless strictly indicated. No data are available in patients with hepatic impairment treated by docetaxel in combination in the other indications.

#### Paediatric population

The safety and efficacy of Docetaxel in nasopharyngeal carcinoma in children aged 1 month to less than 18 years have not yet been established.

There is no relevant use of Docetaxel in the paediatric population in the indications breast cancer, nonsmall cell lung cancer, prostate cancer, gastric carcinoma and head and neck cancer, not including type II and III less differentiated nasopharyngeal carcinoma.

#### Older people

Based on a population pharmacokinetic analysis, there are no special instructions for use in the older people.

In combination with capecitabine, for patients 60 years of age or more, a starting dose reduction of capecitabine to 75% is recommended (see capecitabine summary of product characteristics).

#### Method of administration

For instructions on preparation and administration of the product, see section 6.6.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Docetaxel must not be used in patients with baseline neutrophil count of  $< 1,500$  cells/mm<sup>3</sup>.

Docetaxel must not be used in patients with severe liver impairment since there is no data available (see sections 4.2 and 4.4).

Contraindications for other medicinal products also apply, when combined with docetaxel.

#### **4.4 Special warnings and precautions for use**

For breast and non-small cell lung cancers, premedication consisting of an oral corticosteroid, such as dexamethasone 16 mg per day (e.g. 8 mg BID) for 3 days starting 1 day prior to docetaxel administration, unless contraindicated, can reduce the incidence and severity of fluid retention as well as the severity of hypersensitivity reactions. For prostate cancer, the premedication is oral dexamethasone 8 mg, 12 hours, 3 hours and 1 hour before the docetaxel infusion (see section 4.2).

##### Haematology

Neutropenia is the most frequent adverse reaction of docetaxel. Neutrophil nadirs occurred at a median of 7 days but this interval may be shorter in heavily pre-treated patients. Frequent monitoring of complete blood counts should be conducted on all patients receiving docetaxel. Patients should be retreated with docetaxel when neutrophils recover to a level  $\geq 1,500$  cells/mm<sup>3</sup> (see section 4.2).

In the case of severe neutropenia ( $< 500$  cells/mm<sup>3</sup> for seven days or more) during a course of docetaxel therapy, a reduction in dose for subsequent courses of therapy or the use of appropriate symptomatic measures are recommended (see section 4.2).

In patients treated with docetaxel in combination with cisplatin and 5-fluorouracil (TCF), febrile neutropenia and neutropenic infection occurred at lower rates when patients received prophylactic G-CSF. Patients treated with TCF should receive prophylactic G-CSF to mitigate the risk of complicated neutropenia (febrile neutropenia, prolonged neutropenia or neutropenic infection). Patients receiving TCF should be closely monitored (see sections 4.2 and 4.8).

In patients treated with docetaxel in combination with doxorubicin and cyclophosphamide (TAC), febrile neutropenia and/or neutropenic infection occurred at lower rates when patients received primary G-CSF prophylaxis. Primary G-CSF prophylaxis should be considered in patients who receive adjuvant therapy with TAC for breast cancer to mitigate the risk of complicated neutropenia (febrile neutropenia, prolonged neutropenia or neutropenic

infection). Patients receiving TAC should be closely monitored (see sections 4.2 and 4.8).

#### Gastrointestinal reactions

Caution is recommended for patients with neutropenia, particularly at risk for developing gastrointestinal complications. Although majority of cases occurred during the first or second cycle of docetaxel containing regimen, enterocolitis could develop at any time, and could lead to death as early as on the first day of onset. Patients should be closely monitored for early manifestations of serious gastrointestinal toxicity (see sections 4.2, 4.4 Haematology, and 4.8).

#### Hypersensitivity reactions

Patients should be observed closely for hypersensitivity reactions especially during the first and second infusions. Hypersensitivity reactions may occur within a few minutes following the initiation of the infusion of docetaxel, thus facilities for the treatment of hypotension and bronchospasm should be available. If hypersensitivity reactions occur, minor symptoms such as flushing or localized cutaneous reactions do not require interruption of therapy. However, severe reactions, such as severe hypotension, bronchospasm or generalised rash/erythema require immediate discontinuation of docetaxel and appropriate therapy. Patients who have developed severe hypersensitivity reactions should not be re-challenged with docetaxel. Patients who have previously experienced a hypersensitivity reaction to paclitaxel may be at risk to develop hypersensitivity reaction to docetaxel, including more severe hypersensitivity reaction. These patients should be closely monitored during initiation of docetaxel therapy.

#### Cutaneous reactions

Localised skin erythema of the extremities (palms of the hands and soles of the feet) with oedema followed by desquamation has been observed. Severe symptoms such as eruptions followed by desquamation which lead to interruption or discontinuation of docetaxel treatment were reported (see section 4.2).

Severe Cutaneous Adverse Reactions (SCARs) such as Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported with docetaxel treatment. Patients should be informed about the signs and symptoms of serious skin manifestations and closely monitored. If signs and symptoms suggestive of these reactions appear discontinuation of docetaxel should be considered.

#### Fluid retention

Patients with severe fluid retention such as pleural effusion, pericardial effusion and ascites should be monitored closely.

#### Respiratory disorders

Acute respiratory distress syndrome, interstitial pneumonia/pneumonitis, interstitial lung disease, pulmonary fibrosis and respiratory failure have been reported and may be associated with fatal outcome. Cases of radiation pneumonitis have been reported in patients receiving concomitant radiotherapy.

If new or worsening pulmonary symptoms develop, patients should be closely monitored, promptly investigated, and appropriately treated. Interruption of docetaxel therapy is recommended until diagnosis is available. Early use of supportive care measures may help improve the condition. The benefit of resuming docetaxel treatment must be carefully evaluated.

#### Patients with liver impairment

In patients treated with docetaxel at 100 mg/m<sup>2</sup> as single agent who have serum transaminase levels (ALT and/or AST) greater than 1.5 times the ULN concurrent with serum alkaline phosphatase levels greater than 2.5 times the ULN, there is a higher risk of developing severe adverse reactions such as toxic deaths including sepsis and gastrointestinal haemorrhage which can be fatal, febrile neutropenia, infections, thrombocytopenia, stomatitis and asthenia. Therefore, the recommended dose of docetaxel in those patients with elevated liver function test (LFTs) is 75 mg/m<sup>2</sup> and LFTs should be measured at baseline and before each cycle (see section 4.2).

For patients with serum bilirubin levels > ULN and/or ALT and AST > 3.5 times the ULN concurrent with serum alkaline phosphatase levels > 6 times the ULN, no dose-reduction can be recommended and docetaxel should not be used unless strictly indicated.

In combination with cisplatin and 5-fluorouracil for the treatment of patients with gastric adenocarcinoma, the pivotal clinical study excluded patients with ALT and/or AST > 1.5 × ULN associated with alkaline phosphatase > 2.5 × ULN, and bilirubin > 1 × ULN; for these patients, no dose-reductions can be recommended and docetaxel should not be used unless strictly indicated. No data are available in patients with hepatic impairment treated by docetaxel in combination in the other indications.

#### Patients with renal impairment

There are no data available in patients with severely impaired renal function treated with docetaxel.

#### Nervous system

The development of severe peripheral neurotoxicity requires a reduction of dose (see section 4.2).

#### Cardiac toxicity

Heart failure has been observed in patients receiving docetaxel in combination with trastuzumab, particularly following anthracycline (doxorubicin or epirubicin)-containing chemotherapy. This may be moderate to severe and has been associated with death (see section 4.8).

When patients are candidates for treatment with docetaxel in combination with trastuzumab, they should undergo baseline cardiac assessment. Cardiac function should be further monitored during treatment (e.g. every three months) to help identify patients who may develop cardiac dysfunction. For more details see summary of product characteristics of trastuzumab.

Ventricular arrhythmia including ventricular tachycardia (sometimes fatal) has been reported in patients treated with docetaxel in combination regimens including doxorubicin, 5-fluorouracil and/ or cyclophosphamide (see section 4.8). Baseline cardiac assessment is recommended.

### Eye disorders

Cystoid macular oedema (CMO) has been reported in patients treated with docetaxel. Patients with impaired vision should undergo a prompt and complete ophthalmologic examination. In case CMO is diagnosed, docetaxel treatment should be discontinued and appropriate treatment initiated (see section 4.8).

### Second primary malignancies

Second primary malignancies have been reported when docetaxel was given in combination with anticancer treatments known to be associated with second primary malignancies. Second primary malignancies (including acute myeloid leukaemia, myelodysplastic syndrome and non-Hodgkin lymphoma) may occur several months or years after docetaxel-containing therapy. Patients should be monitored for second primary malignancies (see section 4.8).

### Tumour Lysis Syndrome

Tumour lysis syndrome has been reported with docetaxel after the first or the second cycle (see section 4.8). Patients at risk of tumour lysis syndrome (e.g. with renal impairment, hyperuricemia, bulky tumour, rapid progression) should be closely monitored. Correction of dehydration and treatment of high uric acid levels are recommended prior to initiation of treatment.

### Others

Contraceptive measures must be taken by both men and women during treatment and for men at least 6 months after cessation of therapy (see section 4.6).

The concomitant use of docetaxel with strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin and voriconazole) should be avoided (see section 4.5).

### Additional cautions for use in adjuvant treatment of breast cancer

### Complicated neutropenia

For patients who experience complicated neutropenia (prolonged neutropenia, febrile neutropenia or infection), G-CSF and dose reduction should be considered (see section 4.2).

#### Gastrointestinal reactions

Symptoms such as early abdominal pain and tenderness, fever, diarrhoea, with or without neutropenia, may be early manifestations of serious gastrointestinal toxicity and should be evaluated and treated promptly.

#### Congestive heart failure (CHF)

Patients should be monitored for symptoms of congestive heart failure during therapy and during the follow up period. In patients treated with the TAC regimen for node positive breast cancer, the risk of CHF has been shown to be higher during the first year after treatment (see sections 4.8 and 5.1).

#### Patients with 4+ nodes

As the benefit observed in patient with 4+ nodes was not statistically significant on disease-free survival (DFS) and overall survival (OS), the positive benefit/risk ratio for TAC in patients with 4+ nodes was not fully demonstrated at the final analysis (see section 5.1).

#### Elderly

#### Cautions for use in adjuvant treatment of breast cancer

There are limited data available in patients > 70 years of age on docetaxel use in combination with doxorubicin and cyclophosphamide.

#### Cautions for use in castration-resistant prostate cancer

Of the 333 patients treated with docetaxel every three weeks in a prostate cancer study (TAX 327), 209 patients were 65 years of age or greater and 68 patients were older than 75 years. In patients treated with docetaxel every three weeks, the incidence of related nail changes occurred at a rate  $\geq$  10% higher in patients who were 65 years of age or greater compared to younger patients. The incidence of related fever, diarrhoea, anorexia, and peripheral oedema occurred at rates  $\geq$  10% higher in patients who were 75 years of age or greater versus less than 65 years.

#### Cautions for use in hormone-sensitive prostate cancer

Of the 545 patients treated with docetaxel every 3 weeks in a hormone-sensitive prostate cancer study (STAMPEDE), 296 patients were 65 years of age or older, and 48 patients were 75 years of age or older. More patients aged  $\geq 65$  years in the docetaxel arm reported hypersensitivity reaction, neutropenia, anaemia, fluid retention, dyspnea, and nail changes when compared to the patients aged less than 65 years. None of these increases in frequency reached 10% difference with the control arm. In patients who were 75 years of age or older, when compared to younger patients, neutropenia, anaemia, diarrhea, dyspnea and upper respiratory tract infection were reported with a greater incidence (at least 10% higher).

#### Cautions for use in gastric adenocarcinoma cancer

Among the 300 (221 patients in the phase III part of the study and 79 patients in the phase II part) patients treated with docetaxel in combination with cisplatin and 5-fluorouracil in the gastric cancer study, 74 were 65 years of age or older and 4 patients were 75 years of age or older. The incidence of serious adverse events was higher in older people compared to younger patients. The incidence of the following adverse events (all grades): lethargy, stomatitis, neutropenic infection occurred at rates  $\geq 10\%$  higher in patients who were 65 years of age or older compared to younger patients. Older people treated with TCF should be closely monitored.

#### Excipients

This medicinal product contains 13% (w/v) Alcohol (95% v/v) (Absolute Alcohol Content USP 15.25% v/v) in water for injection USP q.s. to 9 mL

Harmful for those suffering from alcoholism.

To be taken into account in pregnant or breast-feeding women, in children and in high-risk groups such as patients with liver disease, or epilepsy.

Consideration should be given to possible effects on the central nervous system.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

The amount of alcohol in this medicinal product may alter the effects of other medicinal products.

*In vitro* studies have shown that the metabolism of docetaxel may be modified by the concomitant administration of compounds which induce, inhibit or are metabolised by (and thus may inhibit the enzyme competitively) cytochrome P450-3A such as ciclosporine, ketoconazole and erythromycin. As a result, caution should be exercised when treating patients with these medicinal products as concomitant therapy since there is a potential for a significant interaction.

In case of combination with CYP3A4 inhibitors, the occurrence of docetaxel adverse reactions may increase, as a result of reduced metabolism. If the concomitant use of a strong CYP3A4 inhibitor (e.g., ketoconazole, itraconazole, clarithromycin, indinavir, nefazodone, nelfinavir, ritonavir, saquinavir, telithromycin and voriconazole) cannot be avoided, a close clinical surveillance is warranted and a dose-adjustment of docetaxel may be suitable during the treatment with the strong CYP3A4 inhibitor (see section 4.4). In a pharmacokinetic study with 7 patients, the co-administration of docetaxel with the strong CYP3A4 inhibitor ketoconazole leads to a significant decrease in docetaxel clearance by 49%.

Docetaxel pharmacokinetics in the presence of prednisone was studied in patients with metastatic prostate cancer. Docetaxel is metabolised by CYP3A4 and prednisone is known to induce CYP3A4. No statistically significant effect of prednisone on the pharmacokinetics of docetaxel was observed.

Docetaxel is highly protein bound (> 95%). Although the possible *in vivo* interaction of docetaxel with concomitantly administered medicinal product has not been investigated formally, *in vitro* interactions with tightly protein-bound agents such as erythromycin, diphenhydramine, propranolol, propafenone, phenytoin, salicylate, sulfamethoxazole and sodium valproate did not affect protein binding of docetaxel. In addition, dexamethasone did not affect protein binding of docetaxel. Docetaxel did not influence the binding of digitoxin.

The pharmacokinetics of docetaxel, doxorubicin and cyclophosphamide were not influenced by their co-administration. Limited data from a single uncontrolled study were suggestive of an interaction between docetaxel and carboplatin. When combined to docetaxel, the clearance of carboplatin was about 50% higher than values previously reported for carboplatin monotherapy.

## **4.6 Fertility, pregnancy and lactation**

### Pregnancy

There is no information on the use of docetaxel in pregnant women. Docetaxel has been shown to be both embryotoxic and foetotoxic in rabbits and rats, and to reduce fertility in rats. As with other cytotoxic medicinal products, docetaxel may cause foetal harm when administered to pregnant women. Therefore, docetaxel must not be used during pregnancy unless clearly indicated.

Women of childbearing age receiving docetaxel should be advised to avoid becoming pregnant, and to inform the treating physician immediately should this occur.

### Breast-feeding

Docetaxel is a lipophilic substance but it is not known whether it is excreted in human milk.

Consequently, because of the potential for adverse reactions in nursing infants, breast feeding must be discontinued for the duration of docetaxel therapy.

### Contraception in males and females

An effective method of contraception should be used during treatment.

### Fertility

In non clinical studies, docetaxel has genotoxic effects and may alter male fertility (see section 5.3). Therefore, men being treated with docetaxel are advised not to father a child during and up to 6 months after treatment and to seek advice on conservation of sperm prior to treatment. **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed. The amount of alcohol in this medicinal product and the side effects of the product may impair the ability to drive or use machines (see

sections 4.4 and 4.8). Therefore, patients should be warned of the potential impact of the amount of alcohol and the side effects of this medicinal product on the ability to drive or use machines, and be advised not to drive or use machines if they experience these side effects during treatment.

#### **4.8 Undesirable effects**

##### Summary of the safety profile for all indications

The adverse reactions considered to be possibly or probably related to the administration of docetaxel have been obtained in:

- 1312 and 121 patients who received 100 mg/m<sup>2</sup> and 75 mg/m<sup>2</sup> of docetaxel as a single agent respectively.
- 258 patients who received docetaxel in combination with doxorubicin.
- 406 patients who received docetaxel in combination with cisplatin.
- 92 patients treated with docetaxel in combination with trastuzumab.
- 255 patients who received docetaxel in combination with capecitabine.
- 332 patients (TAX327) who received docetaxel in combination with prednisone or prednisolone (clinically important treatment related adverse events are presented).
- 1276 patients (744 and 532 in TAX 316 and GEICAM 9805 respectively) who received docetaxel in combination with doxorubicin and cyclophosphamide (clinically important treatment related adverse events are presented).
- 300 gastric adenocarcinoma patients (221 patients in the phase III part of the study and 79 patients in the phase II part) who received docetaxel in combination with cisplatin and 5fluorouracil (clinically important treatment related adverse events are presented).
- 174 and 251 head and neck cancer patients who received docetaxel in combination with cisplatin and 5-fluorouracil (clinically important treatment related adverse events are presented).
- 545 patients (STAMPEDE study) who received docetaxel in combination with prednisone or prednisolone and ADT.

These reactions were described using the NCI Common Toxicity Criteria (grade 3 = G3; grade 3-4 = G3/4; grade 4 = G4), the COSTART and the MedDRA terms. Frequencies are defined as: very common ( $\geq 1/10$ ), common

( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ); not known (cannot be estimated from available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. The most commonly reported adverse reactions of docetaxel alone are: neutropenia (which was reversible and not cumulative; the median day to nadir was 7 days and the median duration of severe neutropenia ( $< 500$  cells/mm<sup>3</sup>) was 7 days), anaemia, alopecia, nausea, vomiting, stomatitis, diarrhoea and asthenia. The severity of adverse events of docetaxel may be increased when docetaxel is given in combination with other chemotherapeutic agents.

For combination with trastuzumab, adverse events (all grades) reported in  $\geq 10\%$  are displayed. There was an increased incidence of SAEs (40% vs. 31%) and Grade 4 AEs (34% vs. 23%) in the trastuzumab combination arm compared to docetaxel monotherapy.

For combination with capecitabine, the most frequent treatment-related undesirable effects ( $\geq 5\%$ ) reported in a phase III study in breast cancer patients failing anthracycline treatment are presented (see capecitabine summary of product characteristics).

For combination with ADT and with prednisone or prednisolone (STAMPEDE study), adverse events occurring over the 6 cycles of treatment with docetaxel and having at least 2% higher incidence in the docetaxel treatment arm by comparison to the control arm, are presented, using the CTCAE grading scale.

The following adverse reactions are frequently observed with docetaxel:

#### Immune system disorders

Hypersensitivity reactions have generally occurred within a few minutes following the start of the infusion of docetaxel and were usually mild to moderate. The most frequently reported symptoms were flushing, rash with or without pruritus, chest tightness, back pain, dyspnoea and fever or chills. Severe reactions were characterised by hypotension and/or bronchospasm or generalized rash/erythema (see section 4.4).

### Nervous system disorders

The development of severe peripheral neurotoxicity requires a reduction of dose (see sections 4.2 and 4.4). Mild to moderate neuro-sensory signs are characterised by paresthesia, dysesthesia or pain including burning. Neuro-motor events are mainly characterised by weakness. Skin and subcutaneous tissue disorders

Reversible cutaneous reactions have been observed and were generally considered as mild to moderate. Reactions were characterised by a rash including localised eruptions mainly on the feet and hands (including severe hand and foot syndrome), but also on the arms, face or thorax, and frequently associated with pruritus. Eruptions generally occurred within one week after the docetaxel infusion. Less frequently, severe symptoms such as eruptions followed by desquamation which rarely lead to interruption or discontinuation of docetaxel treatment were reported (see sections 4.2 and 4.4). Severe nail disorders are characterised by hypo- or hyperpigmentation and sometimes pain and onycholysis.

### General disorders and administration site conditions

Infusion site reactions were generally mild and consisted of hyper pigmentation, inflammation, redness or dryness of the skin, phlebitis or extravasation and swelling of the vein.

Fluid retention includes events such as peripheral oedema and less frequently pleural effusion, pericardial effusion, ascites and weight gain. The peripheral oedema usually starts at the lower extremities and may become generalised with a weight gain of 3 kg or more. Fluid retention is cumulative in incidence and severity (see section 4.4).

### Tabulated list of adverse reactions in breast cancer for Docetaxel 100 mg/m<sup>2</sup> single agent

| <b>MedDRA<br/>system<br/>organ classes</b> | <b>Very common<br/>adverse reactions</b> | <b>Common adverse<br/>reactions</b> | <b>Uncommon<br/>adverse<br/>reactions</b> |
|--------------------------------------------|------------------------------------------|-------------------------------------|-------------------------------------------|
|--------------------------------------------|------------------------------------------|-------------------------------------|-------------------------------------------|

|                                                 |                                                                                                             |                                                       |                 |
|-------------------------------------------------|-------------------------------------------------------------------------------------------------------------|-------------------------------------------------------|-----------------|
| Infections and infestations                     | Infections (G3/4: 5.7%; including sepsis and pneumonia, fatal in 1.7%)                                      | Infection associated with G4 neutropenia (G3/4: 4.6%) |                 |
| Blood and lymphatic system disorders            | Neutropenia (G4: 76.4%); Anaemia (G3/4: 8.9%); Febrile neutropenia                                          | Thrombocytopenia (G4: 0.2%)                           |                 |
| Immune system disorders                         | Hypersensitivity (G3/4: 5.3%)                                                                               |                                                       |                 |
| Metabolism and nutrition disorders              | Anorexia                                                                                                    |                                                       |                 |
| Nervous system disorders                        | Peripheral sensory neuropathy (G3: 4.1%); Peripheral motor neuropathy (G3/4: 4%); Dysgeusia (severe: 0.07%) |                                                       |                 |
| Cardiac disorders                               |                                                                                                             | Arrhythmia (G3/4: 0.7%)                               | Cardiac failure |
| Vascular disorders                              |                                                                                                             | Hypotension; Hypertension; Haemorrhage                |                 |
| Respiratory, thoracic and mediastinal disorders | Dyspnoea (severe: 2.7%)                                                                                     |                                                       |                 |

|                                                      |                                                                                                |                                                                                                                                                        |                             |
|------------------------------------------------------|------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|
| Gastrointestinal disorders                           | Stomatitis (G3/4: 5.3%);<br>Diarrhoea (G3/4: 4%);<br>Nausea (G3/4: 4%);<br>Vomiting (G3/4: 3%) | Constipation (severe: 0.2%);<br>Abdominal pain (severe: 1%);<br>Gastrointestinal haemorrhage (severe: 0.3%)                                            | Oesophagitis (severe: 0.4%) |
| Skin and subcutaneous tissue disorders               | Alopecia;<br>Skin reaction (G3/4: 5.9%);<br>Nail disorders (severe: 2.6%)                      |                                                                                                                                                        |                             |
| Musculoskeletal and connective tissue disorders      | Myalgia (severe: 1.4%)                                                                         | Arthralgia                                                                                                                                             |                             |
| General disorders and administration site conditions | Fluid retention (severe: 6.5%);<br>Asthenia (severe: 11.2%);<br>Pain                           | Infusion site reaction;<br>Non-cardiac chest pain (severe: 0.4%)                                                                                       |                             |
| Investigations                                       |                                                                                                | G3/4 Blood bilirubin increased (< 5%);<br>G3/4 Blood alkaline phosphatase increased (< 4%);<br>G3/4 AST increased (< 3%);<br>G3/4 ALT increased (< 2%) |                             |

Description of selected adverse reactions in breast cancer for Docetaxel 100 mg/m<sup>2</sup> single agent

*Blood and lymphatic system disorders*

Rare: bleeding episodes associated with grade 3/4 thrombocytopenia.

*Nervous system disorders*

Reversibility data are available among 35.3% of patients who developed neurotoxicity following docetaxel treatment at 100 mg/m<sup>2</sup> as single agent. The events were spontaneously reversible within 3 months.

*Skin and subcutaneous tissue disorders*

Very rare: one case of alopecia non-reversible at the end of the study. 73% of the cutaneous reactions were reversible within 21 days.

*General disorders and administration site conditions*

The median cumulative dose to treatment discontinuation was more than 1,000 mg/m<sup>2</sup> and the median time to fluid retention reversibility was 16.4 weeks (range 0 to 42 weeks). The onset of moderate and severe retention is delayed (median cumulative dose: 818.9 mg/m<sup>2</sup>) in patients with premedication compared with patients without premedication (median cumulative dose: 489.7 mg/m<sup>2</sup>); however, it has been reported in some patients during the early courses of therapy.

Tabulated list of adverse reactions in non-small cell lung cancer for Docetaxel 75 mg/m<sup>2</sup> single Agent

| <b>MedDRA system organ classes</b>   | <b>Very common adverse reactions</b>               | <b>Common adverse reactions</b> |
|--------------------------------------|----------------------------------------------------|---------------------------------|
| Infections and infestations          | Infections (G3/4: 5%)                              |                                 |
| Blood and lymphatic system disorders | Neutropenia (G4: 54.2%);<br>Anaemia (G3/4: 10.8%); | Febrile neutropenia             |

|                                                      |                                                                                             |                                          |
|------------------------------------------------------|---------------------------------------------------------------------------------------------|------------------------------------------|
|                                                      | Thrombocytopenia (G4: 1.7%)                                                                 |                                          |
| Immune system disorders                              |                                                                                             | Hypersensitivity (no severe)             |
| Metabolism and nutrition disorders                   | Anorexia                                                                                    |                                          |
| Nervous system disorders                             | Peripheral sensory neuropathy (G3/4: 0.8%)                                                  | Peripheral motor neuropathy (G3/4: 2.5%) |
| Cardiac disorders                                    |                                                                                             | Arrhythmia (no severe)                   |
| Vascular disorders                                   |                                                                                             | Hypotension                              |
| Gastrointestinal disorders                           | Nausea (G3/4: 3.3%); Stomatitis (G3/4: 1.7%); Vomiting (G3/4: 0.8%); Diarrhoea (G3/4: 1.7%) | Constipation                             |
| Skin and subcutaneous tissue disorders               | Alopecia; Skin reaction (G3/4: 0.8%)                                                        | Nail disorders (severe: 0.8%)            |
| Musculoskeletal and connective tissue disorders      |                                                                                             | Myalgia                                  |
| General disorders and administration site conditions | Asthenia (severe: 12.4%); Fluid retention (severe: 0.8%); Pain                              |                                          |
| Investigations                                       |                                                                                             | G3/4 Blood bilirubin increased (< 2%)    |

Tabulated list of adverse reactions in breast cancer for Docetaxel 75 mg/m<sup>2</sup> in combination with Doxorubicin

| <b>MedDRA system organ classes</b> | <b>Very common adverse reactions</b> | <b>Common adverse reactions</b> | <b>Uncommon adverse reactions</b> |
|------------------------------------|--------------------------------------|---------------------------------|-----------------------------------|
| Infections and infestations        | Infection (G3/4: 7.8%)               |                                 |                                   |

|                                        |                                                                                                          |                                            |             |
|----------------------------------------|----------------------------------------------------------------------------------------------------------|--------------------------------------------|-------------|
| Blood and lymphatic system disorders   | Neutropenia (G4: 91.7%);<br>Anaemia (G3/4: 9.4%);<br>Febrile neutropenia;<br>Thrombocytopenia (G4: 0.8%) |                                            |             |
| Immune system disorders                |                                                                                                          | Hypersensitivity (G3/4: 1.2%)              |             |
| Metabolism and nutrition disorders     |                                                                                                          | Anorexia                                   |             |
| Nervous system disorders               | Peripheral sensory neuropathy (G3: 0.4%)                                                                 | Peripheral motor neuropathy (G3/4: 0.4%)   |             |
| Cardiac disorders                      |                                                                                                          | Cardiac failure;<br>Arrhythmia (no severe) |             |
| Vascular disorders                     |                                                                                                          |                                            | Hypotension |
| Gastrointestinal disorders             | Nausea (G3/4: 5%); Stomatitis (G3/4: 7.8%);<br>Diarrhoea (G3/4: 6.2%);                                   |                                            |             |
|                                        | Vomiting (G3/4: 5%); Constipation                                                                        |                                            |             |
| Skin and subcutaneous tissue disorders | Alopecia;<br>Nail disorders (severe: 0.4%);<br>Skin reaction (no severe)                                 |                                            |             |

|                                                      |                                                                     |                                                                                             |                                                         |
|------------------------------------------------------|---------------------------------------------------------------------|---------------------------------------------------------------------------------------------|---------------------------------------------------------|
| Musculoskeletal and connective tissue disorders      |                                                                     | Myalgia                                                                                     |                                                         |
| General disorders and administration site conditions | Asthenia (severe: 8.1%);<br>Fluid retention (severe: 1.2%);<br>Pain | Infusion site reaction                                                                      |                                                         |
| Investigations                                       |                                                                     | G3/4 Blood bilirubin increased (< 2.5%); G3/4 Blood alkaline phosphatase increased (< 2.5%) | G3/4 AST increased (< 1%);<br>G3/4 ALT increased (< 1%) |

Tabulated list of adverse reactions in non-small cell lung cancer for Docetaxel 75 mg/m<sup>2</sup> in combination with cisplatin

| <b>MedDRA system organ classes</b>   | <b>Very common adverse reactions</b>                                             | <b>Common adverse reactions</b> | <b>Uncommon adverse reactions</b> |
|--------------------------------------|----------------------------------------------------------------------------------|---------------------------------|-----------------------------------|
| Infections and infestations          | Infection (G3/4: 5.7%)                                                           |                                 |                                   |
| Blood and lymphatic system disorders | Neutropenia (G4: 51.5%);<br>Anaemia (G3/4: 6.9%);<br>Thrombocytopenia (G4: 0.5%) | Febrile neutropenia             |                                   |
| Immune system disorders              | Hypersensitivity (G3/4: 2.5%)                                                    |                                 |                                   |
| Metabolism and nutrition disorders   | Anorexia                                                                         |                                 |                                   |

|                                                      |                                                                                                    |                              |                 |
|------------------------------------------------------|----------------------------------------------------------------------------------------------------|------------------------------|-----------------|
| Nervous system disorders                             | Peripheral sensory neuropathy (G3: 3.7%);<br>Peripheral motor neuropathy (G3/4: 2%)                |                              |                 |
| Cardiac disorders                                    |                                                                                                    | Arrhythmia (G3/4: 0.7%)      | Cardiac failure |
| Vascular disorders                                   |                                                                                                    | Hypotension (G3/4: 0.7%)     |                 |
| Gastrointestinal disorders                           | Nausea (G3/4: 9.6%);<br>Vomiting (G3/4: 7.6%);<br>Diarrhoea (G3/4: 6.4%);<br>Stomatitis (G3/4: 2%) | Constipation                 |                 |
| Skin and subcutaneous tissue disorders               | Alopecia;<br>Nail disorders (severe: 0.7%);<br>Skin reaction (G3/4: 0.2%)                          |                              |                 |
| Musculoskeletal and connective tissue disorders      | Myalgia (severe: 0.5%)                                                                             |                              |                 |
| General disorders and administration site conditions | Asthenia (severe: 9.9%);<br>Fluid retention (severe: 0.7%);<br>Fever (G3/4: 1.2%)                  | Infusion site reaction; Pain |                 |

|                |  |                                                                     |                                                                                |
|----------------|--|---------------------------------------------------------------------|--------------------------------------------------------------------------------|
| Investigations |  | G3/4 Blood bilirubin increased (2.1%);<br>G3/4 ALT increased (1.3%) | G3/4 AST increased (0.5%);<br>G3/4 Blood alkaline phosphatase increased (0.3%) |
|----------------|--|---------------------------------------------------------------------|--------------------------------------------------------------------------------|

Tabulated list of adverse reactions in breast cancer for Docetaxel 100 mg/m<sup>2</sup> in combination with Trastuzumab

| <b>MedDRA system organ classes</b>              | <b>Very common adverse reactions</b>                                                                                               | <b>Common adverse reactions</b> |
|-------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|---------------------------------|
| Blood and lymphatic system disorders            | Neutropenia (G3/4: 32%); Febrile neutropenia (includes neutropenia associated with fever and antibiotic use) or neutropenic sepsis |                                 |
| Metabolism and nutrition disorders              | Anorexia                                                                                                                           |                                 |
| Psychiatric disorders                           | Insomnia                                                                                                                           |                                 |
| Nervous system disorders                        | Paresthesia; Headache; Dysgeusia; Hypoaesthesia                                                                                    |                                 |
| Eye disorders                                   | Lacrimation increased; Conjunctivitis                                                                                              |                                 |
| Cardiac disorders                               |                                                                                                                                    | Cardiac failure                 |
| Vascular disorders                              | Lymphoedema                                                                                                                        |                                 |
| Respiratory, thoracic and mediastinal disorders | Epistaxis; Pharyngolaryngeal pain; Nasopharyngitis; Dyspnoea; Cough; Rhinorrhoea                                                   |                                 |

|                                                      |                                                                                                                       |          |
|------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|----------|
| Gastrointestinal disorders                           | Nausea; Diarrhoea; Vomiting; Constipation; Stomatitis; Dyspepsia; Abdominal pain                                      |          |
| Skin and subcutaneous tissue disorders               | Alopecia; Erythema; Rash; Nail disorders                                                                              |          |
| Musculoskeletal and connective tissue disorders      | Myalgia; Arthralgia; Pain in extremity; Bone pain; Back pain                                                          |          |
| General disorders and administration site conditions | Asthenia; Oedema peripheral; Pyrexia; Fatigue; Mucosal inflammation; Pain; Influenza like illness; Chest pain; Chills | Lethargy |
| Investigations                                       | Weight increased                                                                                                      |          |

Description of selected adverse reactions in breast cancer for docetaxel 100 mg/m<sup>2</sup> in combination with trastuzumab

#### *Cardiac disorders*

Symptomatic cardiac failure was reported in 2.2% of the patients who received docetaxel plus trastuzumab compared to 0% of patients given docetaxel alone. In the docetaxel plus trastuzumab arm, 64% had received a prior anthracycline as adjuvant therapy compared with 55% in the docetaxel arm alone.

#### *Blood and lymphatic system disorders*

Very common: Haematological toxicity was increased in patients receiving trastuzumab and docetaxel, compared with docetaxel alone (32% grade 3/4 neutropenia versus 22%, using NCI-CTC criteria). Note that this is likely to be an underestimate since docetaxel alone at a dose of 100 mg/m<sup>2</sup> is known to result in neutropenia in 97% of patients, 76% grade 4, based on nadir blood counts. The incidence of febrile neutropenia/neutropenic sepsis was also increased in patients treated with Herceptin plus docetaxel (23% versus 17% for patients treated with docetaxel alone).

Tabulated list of adverse reactions in breast cancer for TAXOTERE 75 mg/m<sup>2</sup> in combination with capecitabine

| <b>MedDRA system organ classes</b>              | <b>Very common adverse reactions</b>                                                                                                                 | <b>Common adverse reactions</b>                                                         |
|-------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| Infections and infestations                     |                                                                                                                                                      | Oral candidiasis (G3/4: < 1%)                                                           |
| Blood and lymphatic system disorders            | Neutropenia (G3/4: 63%); Anaemia (G3/4: 10%)                                                                                                         | Thrombocytopenia (G3/4: 3%)                                                             |
| Metabolism and nutrition disorders              | Anorexia (G3/4: 1%); Decreased appetite                                                                                                              | Dehydration (G3/4: 2%)                                                                  |
| Nervous system disorders                        | Dysgeusia (G3/4: < 1%); Paresthesia (G3/4: < 1%)                                                                                                     | Dizziness; Headache (G3/4: < 1%); Neuropathy peripheral                                 |
| Eye disorders                                   | Lacrimation increased                                                                                                                                |                                                                                         |
| Respiratory, thoracic and mediastinal disorders | Pharyngolaryngeal pain (G3/4: 2%)                                                                                                                    | Dyspnoea (G3/4: 1%); Cough (G3/4: < 1%); Epistaxis (G3/4: < 1%)                         |
| Gastrointestinal disorders                      | Stomatitis (G3/4: 18%); Diarrhoea (G3/4: 14%); Nausea (G3/4: 6%); Vomiting (G3/4: 4%); Constipation (G3/4: 1%); Abdominal pain (G3/4: 2%); Dyspepsia | Abdominal pain upper; Dry mouth                                                         |
| Skin and subcutaneous tissue disorders          | Hand-foot syndrome (G3/4: 24%); Alopecia (G3/4: 6%); Nail disorders (G3/4: 2%)                                                                       | Dermatitis; Rash erythematous (G3/4: < 1%); Nail discolouration; Onycholysis (G3/4: 1%) |
| Musculoskeletal and connective tissue disorders | Myalgia (G3/4: 2%); Arthralgia (G3/4: 1%)                                                                                                            | Pain in extremity (G3/4: < 1%);                                                         |

|                                                      |                                                                                                             |                                                          |
|------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|----------------------------------------------------------|
|                                                      |                                                                                                             | Back pain (G3/4: 1%)                                     |
| General disorders and administration site conditions | Asthenia (G3/4: 3%);<br>Pyrexia (G3/4: 1%);<br>Fatigue/weakness (G3/4: 5%);<br>Oedema peripheral (G3/4: 1%) | Lethargy;<br>Pain                                        |
| Investigations                                       |                                                                                                             | Weight decreased;<br>G3/4 Blood bilirubin increased (9%) |

Tabulated list of adverse reactions in metastatic castration-resistant prostate cancer for docetaxel 75 mg/m<sup>2</sup> in combination with prednisone or prednisolone

| <b>MedDRA system organ classes</b>   | <b>Very common adverse reactions</b>                                | <b>Common adverse reactions</b>                       |
|--------------------------------------|---------------------------------------------------------------------|-------------------------------------------------------|
| Infections and infestations          | Infection (G3/4: 3.3%)                                              |                                                       |
| Blood and lymphatic system disorders | Neutropenia (G3/4: 32%); Anaemia (G3/4: 4.9%)                       | Thrombocytopenia (G3/4: 0.6%);<br>Febrile neutropenia |
| Immune system disorders              |                                                                     | Hypersensitivity (G3/4: 0.6%)                         |
| Metabolism and nutrition disorders   | Anorexia (G3/4: 0.6%)                                               |                                                       |
| Nervous system disorders             | Peripheral sensory neuropathy (G3/4: 1.2%);<br>Dysgeusia (G3/4: 0%) | Peripheral motor neuropathy (G3/4: 0%)                |
| Eye disorders                        |                                                                     | Lacrimation increased (G3/4: 0.6%)                    |

|                                                      |                                                                                                                  |                                                                     |
|------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|
| Cardiac disorders                                    |                                                                                                                  | Cardiac left ventricular function decrease (G3/4: 0.3%)             |
| Respiratory, thoracic and mediastinal disorders      |                                                                                                                  | Epistaxis (G3/4: 0%);<br>Dyspnoea (G3/4: 0.6%);<br>Cough (G3/4: 0%) |
| Gastrointestinal disorders                           | Nausea (G3/4: 2.4%);<br>Diarrhoea (G3/4: 1.2%);<br>Stomatitis/Pharyngitis (G3/4: 0.9%);<br>Vomiting (G3/4: 1.2%) |                                                                     |
| Skin and subcutaneous tissue disorders               | Alopecia;<br>Nail disorders (no severe)                                                                          | Exfoliative rash (G3/4: 0.3%)                                       |
| Musculoskeletal and connective bone disorders        |                                                                                                                  | Arthralgia (G3/4: 0.3%); Myalgia (G3/4: 0.3%)                       |
| General disorders and administration site conditions | Fatigue (G3/4: 3.9%);<br>Fluid retention (severe: 0.6%)                                                          |                                                                     |

Tabulated list of adverse reactions in high-risk locally advanced or metastatic hormone-sensitive prostate cancer for docetaxel 75 mg/m<sup>2</sup> in combination with prednisone or prednisolone and ADT (STAMPEDE study)

| <b>MedDRA system organ classes</b>   | <b>Very common adverse reactions</b> | <b>Common adverse reactions</b> |
|--------------------------------------|--------------------------------------|---------------------------------|
| Blood and lymphatic system disorders | Neutropenia (G3-4: 12 %)<br>Anaemia  |                                 |
|                                      | Febrile neutropenia (G3-4: 15%)      |                                 |
| Immune system disorders              |                                      | Hypersensitivity (G3-4: 1%)     |
| Endocrine disorders                  |                                      | Diabetes (G3-4: 1%)             |

|                                                      |                                                                                                                                            |                                                                                                                       |
|------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------|
| Metabolism and nutrition disorders                   |                                                                                                                                            | Anorexia                                                                                                              |
| Psychiatric disorders                                | Insomnia (G3: 1%)                                                                                                                          |                                                                                                                       |
| Nervous system disorders                             | Peripheral sensory neuropathy<br>(≥G3: 2%) <sup>a</sup><br>Headache                                                                        | Dizziness                                                                                                             |
| Eye disorders                                        |                                                                                                                                            | Blurred vision                                                                                                        |
| Cardiac disorders                                    |                                                                                                                                            | Hypotension (G3: 0%)                                                                                                  |
| Respiratory, thoracic and mediastinal disorders      | Dyspnea (G3: 1%)<br>Coughing (G3: 0%)<br>Upper respiratory tract infection (G3: 1%)                                                        | Pharyngitis (G3: 0%)                                                                                                  |
| Gastrointestinal disorders                           | Diarrhea (G3: 3%)<br>Stomatitis (G3: 0%)<br>Constipation (G3: 0%)<br>Nausea (G3: 1%)<br>Dyspepsia<br>Abdominal pain (G3: 0%)<br>Flatulence | Vomiting (G3: 1%)                                                                                                     |
| Skin and subcutaneous tissue disorders               | Alopecia (G3: 3%) <sup>a</sup><br>Nail changes (G3: 1%)                                                                                    | Rash                                                                                                                  |
| Musculoskeletal and connective tissue disorders      | Myalgia                                                                                                                                    |                                                                                                                       |
| General disorders and administration site conditions | Lethargy (G3-4: 2%)<br>Flu-like symptoms (G3: 0%)<br>Asthenia (G3: 0%)<br>Fluid retention                                                  | Fever (G3: 1%)<br>Oral candidiasis<br>Hypocalcaemia (G3: 0%)<br>Hypophosphataemia (G3-4: 1%)<br>Hypokalaemia (G3: 0%) |

<sup>a</sup> From the GETUG AFU15 study

Tabulated list of adverse reactions for adjuvant therapy with docetaxel 75 mg/m<sup>2</sup> in combination with doxorubicin and cyclophosphamide in patients with node-positive (TAX 316) and node-negative (GEICAM 9805) breast cancer - pooled data

| <b>MedDRA system organ classes</b>   | <b>Very common adverse reactions</b>                                                                               | <b>Common adverse reactions</b>        | <b>Uncommon adverse reactions</b>                                         |
|--------------------------------------|--------------------------------------------------------------------------------------------------------------------|----------------------------------------|---------------------------------------------------------------------------|
| Infections and infestations          | Infection (G3/4: 2.4%);<br>Neutropenic infection (G3/4: 2.6%)                                                      |                                        |                                                                           |
| Blood and lymphatic system disorders | Anaemia (G3/4: 3%); Neutropenia (G3/4: 59.2%);<br>Thrombocytopenia (G3/4: 1.6%);<br>Febrile neutropenia (G3/4: NA) |                                        |                                                                           |
| Immune system disorders              |                                                                                                                    | Hypersensitivity (G3/4: 0.6%)          |                                                                           |
| Metabolism and nutrition disorders   | Anorexia (G3/4: 1.5%)                                                                                              |                                        |                                                                           |
| Nervous system disorders             | Dysgeusia (G3/4: 0.6%);<br>Peripheral sensory neuropathy (G3/4: <0.1%)                                             | Peripheral motor neuropathy (G3/4: 0%) | Syncope (G3/4: 0%);<br>Neurotoxicity (G3/4: 0%);<br>Somnolence (G3/4: 0%) |

|                                                 |                                                                                                                        |                                              |                        |
|-------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|------------------------|
| Eye disorders                                   | Conjunctivitis (G3/4: <0.1%)                                                                                           | Lacrimation increased (G3/4: <0.1%)          |                        |
| Cardiac disorders                               |                                                                                                                        | Arrhythmia (G3/4: 0.2%)                      |                        |
| Vascular disorders                              | Hot flush (G3/4: 0.5%)                                                                                                 | Hypotension (G3/4: 0%); Phlebitis (G3/4: 0%) | Lymphoedema (G3/4: 0%) |
| Respiratory, thoracic and mediastinal disorders |                                                                                                                        | Cough (G3/4: 0%)                             |                        |
| Gastrointestinal disorders                      | Nausea (G3/4: 5.0%); Stomatitis (G3/4: 6.0%); Vomiting (G3/4: 4.2%); Diarrhoea (G3/4: 3.4%); Constipation (G3/4: 0.5%) | Abdominal pain (G3/4: 0.4%)                  |                        |
| Skin and subcutaneous tissue disorders          | Alopecia (persisting: <3%); Skin disorder (G3/4: 0.6%); Nail disorders (G3/4: 0.4%)                                    |                                              |                        |
| Musculoskeletal and connective tissue disorders | Myalgia (G3/4: 0.7%); Arthralgia (G3/4: 0.2%)                                                                          |                                              |                        |

|                                                      |                                                                                  |                                                               |  |
|------------------------------------------------------|----------------------------------------------------------------------------------|---------------------------------------------------------------|--|
| Reproductive system and breast disorders             | Amenorrhoea (G3/4: NA)                                                           |                                                               |  |
| General disorders and administration site conditions | Asthenia (G3/4: 10.0%);<br>Pyrexia (G3/4: NA);<br>Oedema peripheral (G3/4: 0.2%) |                                                               |  |
| Investigations                                       |                                                                                  | Weight increased (G3/4: 0%);<br>Weight decreased (G3/4: 0.2%) |  |

Description of selected adverse reactions for adjuvant therapy with docetaxel 75 mg/m<sup>2</sup> in combination with doxorubicin and cyclophosphamide in patients with node-positive (TAX 316) and node-negative (GEICAM 9805) breast cancer

#### *Nervous system disorders*

In study TAX316 peripheral sensory neuropathy started during the treatment period and persisted into the follow-up period in 84 patients (11.3%) in TAC arm and 15 patients (2 %) in FAC arm. At the end of the follow-up period (median follow-up time of 8 years), peripheral sensory neuropathy was observed to be ongoing in 10 patients (1.3%) in TAC arm, and in 2 patients (0.3%) in FAC arm. In GEICAM 9805 study peripheral sensory neuropathy that started during the treatment period persisted into the follow-up period in 10 patients (1.9%) in TAC arm and 4 patients (0.8 %) in FAC arm. At the end of the follow-up period (median follow-up time of 10 years and 5 months), peripheral sensory neuropathy was observed to be ongoing in 3 patients (0.6%) in TAC arm, and in 1 patient (0.2%) in FAC arm.

#### *Cardiac disorders*

In study TAX316, 26 patients (3.5%) in the TAC arm and 17 patients (2.3%) in the FAC arm experienced congestive heart failure. All except one patient in each arm were diagnosed with CHF more than 30 days after the treatment

period. Two patients in the TAC arm and 4 patients in the FAC arm died because of cardiac failure.

In GEICAM 9805 study, 3 patients (0.6 %) in TAC arm and 3 patients (0.6 %) in FAC arm developed congestive heart failure during the follow-up period. At the end of the follow-up period (actual median follow-up time of 10 years and 5 months), no patients had CHF in TAC arm and 1 patient in TAC arm died because of dilated cardiomyopathy, and CHF was observed to be ongoing in 1 patient (0.2%) in FAC arm.

#### *Skin and subcutaneous tissue disorders*

In study TAX316 alopecia persisting into the follow-up period after the end of chemotherapy was reported in 687 of 744 TAC patients (92.3%) and 645 of 736 FAC patients (87.6%).

At the end of the follow-up period (actual median follow-up time of 8 years), alopecia was observed to be ongoing in 29 TAC patients (3.9%) and 16 FAC patients (2.2%).

In GEICAM 9805 study alopecia that started during the treatment period and persisted into the follow up period was observed to be ongoing in 49 patients (9.2 %) in TAC arm and 35 patients (6.7 %) in FAC arm. Alopecia related to study drug started or worsened during the follow-up period in 42 patients (7.9 %) in TAC arm and 30 patients (5.8 %) in FAC arm. At the end of the follow-up period (median follow-up time of 10 years and 5 months), alopecia was observed to be ongoing in 3 patients (0.6%) in TAC arm, and in 1 patient (0.2%) in FAC arm.

#### *Reproductive system and breast disorders*

In TAX316 amenorrhoea that started during the treatment period and persisted into the follow-up period after the end of chemotherapy was reported in 202 of 744 TAC patients (27.2%) and 125 of 736 FAC patients (17.0%). Amenorrhea was observed to be ongoing at the end of the follow-up period (median follow-up time of 8 years) in 121 of 744 TAC patients (16.3%) and 86 FAC patients (11.7%).

In GEICAM 9805 study amenorrhoea that started during the treatment period and persisted into the follow-up period was observed to be ongoing in 18 patients (3.4 %) in TAC arm and 5 patients (1.0 %) in FAC arm. At the end of the follow-up period (median follow-up time of 10 years and 5 months),

amenorrhoea was observed to be ongoing in 7 patients (1.3%) in TAC arm, and in 4 patients (0.8%) in FAC arm.

*General disorders and administration site conditions*

In study TAX316 peripheral oedema that started during the treatment period and persisted into the follow-up period after the end of chemotherapy was observed in 119 of 744 TAC patients (16.0%) and 23 of 736 FAC patients (3.1%). At the end of the follow-up period (actual median follow-up time of 8 years), peripheral oedema was ongoing in 19 TAC patients (2.6%) and 4 FAC patients (0.5%). In study TAX316 lymphoedema that started during the treatment period and persisted into the follow up period after the end of chemotherapy was reported in 11 of 744 TAC patients (1.5%) and 1 of 736 FAC patients (0.1%). At the end of the follow-up period (actual median follow-up time of 8 years), lymphoedema was observed to be ongoing in 6 TAC patients (0.8%) and 1 FAC patient (0.1%).

In study TAX316 asthenia that started during the treatment period and persisted into the follow-up period after the end of chemotherapy was reported in 236 of 744 TAC patients (31.7%) and 180 of 736 FAC patients (24.5%). At the end of the follow-up period (actual median follow-up time of 8 years), asthenia was observed to be ongoing in 29 TAC patients (3.9%) and 16 FAC patients (2.2%).

In study GEICAM 9805 peripheral oedema that started during the treatment period persisted into the follow-up period in 4 patients (0.8%) in TAC arm and in 2 patients (0.4%) in FAC arm. At the end of the follow-up period (median follow-up time of 10 years and 5 months), no patients (0%) in TAC arm had peripheral oedema and it was observed to be ongoing in 1 patient (0.2%) in FAC arm.

Lymphoedema that started during the treatment period persisted into the follow-up period in 5 patients (0.9%) in TAC arm and 2 patients (0.4 %) in FAC arm. At the end of the follow-up period, lymphoedema was observed to be ongoing in 4 patients (0.8%) in TAC arm, and in 1 patient (0.2%) in FAC arm. Asthenia that started during the treatment period and persisted into the follow-up period was observed to be ongoing in 12 patients (2.3 %) in TAC arm and 4 patients (0.8 %) in FAC arm. At the end of the follow-up period, asthenia was observed to be ongoing in 2 patients (0.4%) in TAC arm, and in 2 patients (0.4%) in FAC arm.

*Acute leukaemia / Myelodysplastic syndrome.*

After 10 years of follow up in study TAX316, acute leukaemia was reported in 3 of 744 TAC patients (0.4%) and in 1 of 736 FAC patients (0.1%). One TAC patient (0.1%) and 1 FAC patient (0.1%) died due to AML during the follow-up period (median follow-up time of 8 years). Myelodysplastic syndrome was reported in 2 of 744 TAC patients (0.3%) and in 1 of 736 FAC patients (0.1%).

After 10 years of follow-up in GEICAM 9805 study, acute leukaemia occurred in 1 of 532 (0.2%) patients in TAC arm. No cases were reported in patients in FAC arm. No patient was diagnosed with myelodysplastic syndrome in either treatment groups.

**Neutropenic complications**

Table below shows that the incidence of Grade 4 neutropenia, febrile neutropenia and neutropenic infection was decreased in patients who received primary G-CSF prophylaxis after it was made mandatory in the TAC arm - GEICAM study.

Neutropenic complications in patients receiving TAC with or without primary G-CSF prophylaxis (GEICAM 9805)

|                                   | <b>Without<br/>primary<br/>G-CSF<br/>prophylaxis<br/>(n =<br/>111) n<br/>(%)</b> | <b>With primary<br/>G-CSF<br/>prophylaxis<br/>(n =<br/>421) n<br/>(%)</b> |
|-----------------------------------|----------------------------------------------------------------------------------|---------------------------------------------------------------------------|
| Neutropenia (Grade 4)             | 104 (93.7)                                                                       | 135 (32.1)                                                                |
| Febrile neutropenia               | 28 (25.2)                                                                        | 23 (5.5)                                                                  |
| Neutropenic infection             | 14 (12.6)                                                                        | 21 (5.0)                                                                  |
| Neutropenic infection (Grade 3-4) | 2 (1.8)                                                                          | 5 (1.2)                                                                   |

Tabulated list of adverse reactions in gastric adenocarcinoma cancer for TAXOTERE 75 mg/m<sup>2</sup> in combination with cisplatin and 5-fluorouracil

| <b>MedDRA<br/>system organ</b> | <b>Very common<br/>adverse reactions</b> | <b>Common adverse reactions</b> |
|--------------------------------|------------------------------------------|---------------------------------|
|--------------------------------|------------------------------------------|---------------------------------|

| <b>classes</b>                       |                                                                                                               |                                                                                                         |
|--------------------------------------|---------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------|
| Infections and infestations          | Neutropenic infection; Infection (G3/4: 11.7%)                                                                |                                                                                                         |
| Blood and lymphatic system disorders | Anaemia (G3/4: 20.9%);<br>Neutropenia (G3/4: 83.2%);<br>Thrombocytopenia (G3/4: 8.8%);<br>Febrile neutropenia |                                                                                                         |
| Immune system disorders              | Hypersensitivity (G3/4: 1.7%)                                                                                 |                                                                                                         |
| Metabolism and nutrition disorders   | Anorexia (G3/4: 11.7%)                                                                                        |                                                                                                         |
| Nervous system disorders             | Peripheral sensory neuropathy (G3/4: 8.7%)                                                                    | Dizziness (G3/4: 2.3%);<br>Peripheral motor neuropathy (G3/4: 1.3%)                                     |
| Eye disorders                        |                                                                                                               | Lacrimation increased (G3/4: 0%)                                                                        |
| Ear and labyrinth disorders          |                                                                                                               | Hearing impaired (G3/4: 0%)                                                                             |
| Cardiac disorders                    |                                                                                                               | Arrhythmia (G3/4: 1.0%)                                                                                 |
| Gastrointestinal disorders           | Diarrhoea (G3/4: 19.7%); Nausea (G3/4: 16%);<br>Stomatitis (G3/4: 23.7%);<br>Vomiting (G3/4: 14.3%)           | Constipation (G3/4: 1.0%);<br>Gastrointestinal pain (G3/4: 1.0%);<br>Oesophagitis/dysphagia/odynophagia |

|                                                      |                                                                                           |                                                                                            |
|------------------------------------------------------|-------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|
|                                                      |                                                                                           | hagia (G3/4: 0.7%)                                                                         |
| Skin and subcutaneous tissue disorders               | Alopecia (G3/4: 4.0%)                                                                     | Rash pruritus (G3/4: 0.7%);<br>Nail disorders (G3/4: 0.7%);<br>Skin exfoliation (G3/4: 0%) |
| General disorders and administration site conditions | Lethargy (G3/4: 19.0%); Fever (G3/4: 2.3%); Fluid retention (severe/life-threatening: 1%) |                                                                                            |

Description of selected adverse reactions in gastric adenocarcinoma cancer for docetaxel 75 mg/m<sup>2</sup> in combination with cisplatin and 5-fluorouracil

*Blood and lymphatic system disorders*

Febrile neutropenia and neutropenic infection occurred in 17.2% and 13.5% of patients respectively, regardless of G-CSF use. G-CSF was used for secondary prophylaxis in 19.3% of patients (10.7% of the cycles). Febrile neutropenia and neutropenic infection occurred respectively in 12.1% and 3.4% of patients when patients received prophylactic G-CSF, in 15.6% and 12.9% of patients without prophylactic G-CSF (see section 4.2).

Tabulated list of adverse reactions in head and neck cancer for docetaxel 75 mg/m<sup>2</sup> in combination with cisplatin and 5-fluorouracil

- Induction chemotherapy followed by radiotherapy (TAX 323)

| <b>MedDRA system organ classes</b>                                  | <b>Very common adverse reactions</b>          | <b>Common adverse reactions</b> | <b>Uncommon adverse reactions</b> |
|---------------------------------------------------------------------|-----------------------------------------------|---------------------------------|-----------------------------------|
| Infections and infestations                                         | Infection (G3/4: 6.3%); Neutropenic infection |                                 |                                   |
| Neoplasms benign, malignant and unspecified (incl cysts and polyps) |                                               | Cancer pain (G3/4: 0.6%)        |                                   |

|                                      |                                                                                                     |                                                                                                                                                |                         |
|--------------------------------------|-----------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|
| Blood and lymphatic system disorders | Neutropenia (G3/4: 76.3%);<br>Anaemia (G3/4: 9.2%);<br>Thrombocytopenia (G3/4: 5.2%)                | Febrile neutropenia                                                                                                                            |                         |
| Immune system disorders              |                                                                                                     | Hypersensitivity (no severe)                                                                                                                   |                         |
| Metabolism and nutrition disorders   | Anorexia (G3/4: 0.6%)                                                                               |                                                                                                                                                |                         |
| Nervous system disorders             | Dysgeusia/Parosmia; Peripheral sensory neuropathy (G3/4: 0.6%)                                      | Dizziness                                                                                                                                      |                         |
| Eye disorders                        |                                                                                                     | Lacrimation increased; Conjunctivitis                                                                                                          |                         |
| Ear and labyrinth disorders          |                                                                                                     | Hearing impaired                                                                                                                               |                         |
| Cardiac disorders                    |                                                                                                     | Myocardial ischemia (G3/4:1.7%)                                                                                                                | Arrhythmia (G3/4: 0.6%) |
| Vascular disorders                   |                                                                                                     | Venous disorder (G3/4: 0.6%)                                                                                                                   |                         |
| Gastrointestinal disorders           | Nausea (G3/4: 0.6%);<br>Stomatitis (G3/4:4.0%);<br>Diarrhoea (G3/4: 2.9%);<br>Vomiting (G3/4: 0.6%) | Constipation;<br>Esophagitis/dysphagia/odynophagia (G3/4: 0.6%);<br>Abdominal pain;<br>Dyspepsia;<br>Gastrointestinal haemorrhage (G3/4: 0.6%) |                         |

|                                                      |                                                                               |                                                              |  |
|------------------------------------------------------|-------------------------------------------------------------------------------|--------------------------------------------------------------|--|
| Skin and subcutaneous tissue disorders               | Alopecia (G3/4: 10.9%)                                                        | Rash pruritic;<br>Dry skin;<br>Skin exfoliative (G3/4: 0.6%) |  |
| Musculoskeletal and connective tissue disorders      |                                                                               | Myalgia (G3/4: 0.6%)                                         |  |
| General disorders and administration site conditions | Lethargy (G3/4: 3.4%);<br>Pyrexia (G3/4: 0.6%);<br>Fluid retention;<br>Oedema |                                                              |  |
| Investigations                                       |                                                                               | Weight increased                                             |  |

- Induction chemotherapy followed by chemoradiotherapy (TAX 324)

| <b>MedDRA system organ classes</b> | <b>Very common adverse reactions</b> | <b>Common adverse reactions</b> | <b>Uncommon adverse reactions</b> |
|------------------------------------|--------------------------------------|---------------------------------|-----------------------------------|
| Infections and infestations        | Infection (G3/4: 3.6%)               | Neutropenic infection           |                                   |

|                                                                     |                                                                                                            |                          |                  |
|---------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------|------------------|
| Neoplasms benign, malignant and unspecified (incl cysts and polyps) |                                                                                                            | Cancer pain (G3/4: 1.2%) |                  |
| Blood and lymphatic system disorders                                | Neutropenia (G3/4: 83.5%); Anaemia (G3/4: 12.4%);<br>Thrombocytopenia (G3/4: 4.0%);<br>Febrile neutropenia |                          |                  |
| Immune system disorders                                             |                                                                                                            |                          | Hypersensitivity |

|                                                 |                                                                                                                                                                                        |                                                                                                           |                     |
|-------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------|---------------------|
| Metabolism and nutrition disorders              | Anorexia (G3/4:12.0%)                                                                                                                                                                  |                                                                                                           |                     |
| Nervous system disorders                        | Dysgeusia/Parosmia (G3/4: 0.4%);<br>Peripheral sensory neuropathy (G3/4:1.2%)                                                                                                          | Dizziness (G3/4:2.0%);<br>Peripheral motor neuropathy (G3/4:0.4%)                                         |                     |
| Eye disorders                                   |                                                                                                                                                                                        | Lacrimation increased                                                                                     | Conjunctivitis      |
| Ear and labyrinth disorders                     | Hearing impaired (G3/4: 1.2%)                                                                                                                                                          |                                                                                                           |                     |
| Cardiac disorders                               |                                                                                                                                                                                        | Arrhythmia (G3/4:2.0%)                                                                                    | Ischemia myocardial |
| Vascular disorders                              |                                                                                                                                                                                        |                                                                                                           | Venous disorder     |
| Gastrointestinal disorders                      | Nausea (G3/4: 13.9%);<br>Stomatitis (G3/4: 20.7%);<br>Vomiting (G3/4:8.4%);<br>Diarrhoea (G3/4:6.8%);<br>Esophagitis/dysphagia/odynophagia (G3/4: 12.0%);<br>Constipation (G3/4: 0.4%) | Dyspepsia (G3/4:0.8%);<br>Gastrointestinal pain (G3/4: 1.2%);<br>Gastrointestinal haemorrhage (G3/4:0.4%) |                     |
| Skin and subcutaneous tissue disorders          | Alopecia (G3/4: 4.0%);<br>Rash pruritic                                                                                                                                                | Dry skin ;<br>Desquamation                                                                                |                     |
| Musculoskeletal and connective tissue disorders |                                                                                                                                                                                        | Myalgia (G3/4: 0.4%)                                                                                      |                     |
| General disorders and                           | Lethargy (G3/4: 4.0%);                                                                                                                                                                 |                                                                                                           |                     |

|                                |                                                                         |  |                  |
|--------------------------------|-------------------------------------------------------------------------|--|------------------|
| administration site conditions | Pyrexia (G3/4: 3.6%); Fluid retention (G3/4: 1.2%); Oedema (G3/4: 1.2%) |  |                  |
| Investigations                 | Weight decreased                                                        |  | Weight increased |

#### Post-marketing experience

##### *Neoplasms benign, malignant and unspecified (incl cysts and polyps)*

Second primary malignancies (frequency not known), including non-Hodgkin lymphoma have been reported in association with docetaxel when used in combination with other anticancer treatments known to be associated with second primary malignancies. Acute myeloid leukaemia and myelodysplastic syndrome have been reported (frequency uncommon) in pivotal clinical studies in breast cancer with TAC regimen.

##### *Blood and lymphatic system disorders*

Bone marrow suppression and other haematologic adverse reactions have been reported. Disseminated intravascular coagulation (DIC), often in association with sepsis or multi-organ failure, has been reported.

##### *Immune system disorders*

Some cases of anaphylactic shock, sometimes fatal, have been reported. Hypersensitivity reactions (frequency not known) have been reported with docetaxel in patients who previously experienced hypersensitivity reactions to paclitaxel.

##### *Nervous system disorders*

Rare cases of convulsion or transient loss of consciousness have been observed with docetaxel administration. These reactions sometimes appear during the infusion of the medicinal product.

##### *Eye disorders*

Very rare cases of transient visual disturbances (flashes, flashing lights, scotomata) typically occurring during infusion of the medicinal product and in association with hypersensitivity reactions have been reported. These were reversible upon discontinuation of the infusion. Cases of lacrimation with or without conjunctivitis, as cases of lacrimal duct obstruction resulting in

excessive tearing have been rarely reported. Cases of cystoid macular oedema (CMO) have been reported in patients treated with docetaxel.

#### *Ear and labyrinth disorders*

Rare cases of ototoxicity, hearing impaired and/or hearing loss have been reported.

#### *Cardiac disorders*

Rare cases of myocardial infarction have been reported.

Ventricular arrhythmia including ventricular tachycardia (frequency not known), sometimes fatal, has been reported in patients treated with docetaxel in combination regimens including doxorubicin, 5- fluorouracil and/ or cyclophosphamide.

#### *Vascular disorders*

Venous thromboembolic events have rarely been reported.

#### *Respiratory, thoracic and mediastinal disorders*

Acute respiratory distress syndrome and cases of interstitial pneumonia/ pneumonitis, interstitial lung disease, pulmonary fibrosis and respiratory failure sometimes fatal have rarely been reported. Rare cases of radiation pneumonitis have been reported in patients receiving concomitant radiotherapy.

#### *Gastrointestinal disorders*

Rare cases of enterocolitis, including colitis, ischemic colitis, and neutropenic enterocolitis, have been reported with a potential fatal outcome (frequency not known).

Rare occurrences of dehydration have been reported as a consequence of gastrointestinal events including enterocolitis and gastrointestinal perforation. Rare cases of ileus and intestinal obstruction have been reported.

#### *Hepatobiliary disorders*

Very rare cases of hepatitis, sometimes fatal primarily in patients with pre-existing liver disorders, have been reported.

#### *Skin and subcutaneous tissue disorders*

Cases of cutaneous lupus erythematosus, bullous eruptions such as erythema multiforme and severe cutaneous adverse reactions such as Stevens-Johnson Syndrome (SJS), Toxic Epidermal Necrolysis (TEN) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported with docetaxel. Scleroderma-like changes usually preceded by peripheral lymphoedema have been reported with docetaxel. Cases of permanent alopecia (frequency not known) have been reported.

#### *Renal and urinary disorders*

Renal insufficiency and renal failure have been reported. In about 20% of these cases there were no risk factors for acute renal failure such as concomitant nephrotoxic medicinal products and gastrointestinal disorders.

#### *General disorders and administration site conditions*

Radiation recall phenomena have rarely been reported.

Injection site recall reaction (recurrence of skin reaction at a site of previous extravasation following administration of docetaxel at a different site) has been observed at the site of previous extravasation (frequency not known).

Fluid retention has not been accompanied by acute episodes of oliguria or hypotension. Dehydration and pulmonary oedema have rarely been reported.

#### *Metabolism and nutrition disorders*

Cases of electrolyte imbalance have been reported. Cases of hyponatraemia have been reported, mostly associated with dehydration, vomiting and pneumonia. Hypokalaemia, hypomagnesaemia, and hypocalcaemia were observed, usually in association with gastrointestinal disorders and in particular with diarrhoea. Tumour lysis syndrome, potentially fatal, has been reported (frequency not known).

#### *Musculoskeletal disorder*

Myositis has been reported with docetaxel (frequency not known).

#### *Reporting of suspected adverse reactions*

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk

balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

#### **4.9 Overdose**

There were a few reports of overdose. There is no known antidote for docetaxel overdose. In case of overdose, the patient should be kept in a specialised unit and vital functions closely monitored. In cases of overdose, exacerbation of adverse events may be expected. The primary anticipated complications of overdose would consist of bone marrow suppression, peripheral neurotoxicity and mucositis. Patients should receive therapeutic G-CSF as soon as possible after discovery of overdose.

Other appropriate symptomatic measures should be taken, as needed.

### **5. PHARMACOLOGICAL PROPERTIES**

#### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Taxanes, ATC code: L01CD 02.

##### Mechanism of action

Docetaxel is an antineoplastic agent which acts by promoting the assembly of tubulin into stable microtubules and inhibits their disassembly which leads to a marked decrease of free tubulin. The binding of docetaxel to microtubules does not alter the number of protofilaments.

Docetaxel has been shown in vitro to disrupt the microtubular network in cells which is essential for vital mitotic and interphase cellular functions.

##### Pharmacodynamic effects

Docetaxel was found to be cytotoxic in vitro against various murine and human tumour cell lines and against freshly excised human tumour cells in clonogenic assays. Docetaxel achieves high intracellular concentrations with a long cell residence time. In addition, docetaxel was found to be active on some but not all cell lines over expressing the p-glycoprotein which is encoded by the multidrug resistance gene. In vivo, docetaxel is schedule independent

and has a broad spectrum of experimental anti-tumour activity against advanced murine and human grafted tumours.

### Clinical efficacy and safety

#### Breast cancer

##### *Docetaxel in combination with doxorubicin and cyclophosphamide: adjuvant therapy*

#### Patients with operable node-positive breast cancer (TAX 316)

Data from a multi-centre open label randomised study support the use of docetaxel for the adjuvant treatment of patients with operable node-positive breast cancer and KPS  $\geq$  80%, between 18 and 70 years of age. After stratification according to the number of positive lymph nodes (1-3, 4+), 1491 patients were randomised to receive either docetaxel 75 mg/m<sup>2</sup> administered 1-hour after doxorubicin 50 mg/m<sup>2</sup> and cyclophosphamide 500 mg/m<sup>2</sup> (TAC arm), or doxorubicin 50 mg/m<sup>2</sup> followed by fluorouracil 500 mg/m<sup>2</sup> and cyclophosphamide 500 mg/m<sup>2</sup> (FAC arm). Both regimens were administered once every 3 weeks for 6 cycles. Docetaxel was administered as a 1-hour infusion, all other medicinal products were given as intravenous bolus on day one. G-CSF was administered as secondary prophylaxis to patients who experienced complicated neutropenia (febrile neutropenia, prolonged neutropenia, or infection). Patients on the TAC arm received antibiotic prophylaxis with ciprofloxacin 500 mg orally twice daily for 10 days starting on day 5 of each cycle, or equivalent. In both arms, after the last cycle of chemotherapy, patients with positive oestrogen and/or progesterone receptors received tamoxifen 20 mg daily for up to 5 years. Adjuvant radiation therapy was prescribed according to guidelines in place at participating institutions and was given to 69% of patients who received TAC and 72% of patients who received FAC. Two interim analyses and one final analysis were performed. The first interim analysis was planned 3 years after the date when half of study enrolment was done. The second interim analysis was done after 400 DFS events had been recorded overall, which led to a median follow-up of 55 months. The final analysis was performed when all patients had reached their 10-year follow-up visit (unless they had a DFS event or were lost to followup

before). Disease-free survival (DFS) was the primary efficacy endpoint and Overall survival (OS) was the secondary efficacy endpoint.

A final analysis was performed with an actual median follow up of 96 months. Significantly longer disease-free survival for the TAC arm compared to the FAC arm was demonstrated. Incidence of relapses at 10 years was reduced in patients receiving TAC compared to those who received FAC (39% versus 45%, respectively) i.e. an absolute risk reduction by 6% ( $p = 0.0043$ ). Overall survival at 10 years was also significantly increased with TAC compared to FAC (76% versus 69%, respectively) i.e. an absolute reduction of the risk of death by 7% ( $p = 0.002$ ). As the benefit observed in patients with 4+ nodes was not statistically significant on DFS and OS, the positive benefit/risk ratio for TAC in patients with 4+ nodes was not fully demonstrated at the final analysis.

Overall, the study results demonstrate a positive benefit risk ratio for TAC compared to FAC.

TAC-treated patient subsets according to prospectively defined major prognostic factors were analysed:

| Patient subset              | Number Of patients | Disease free survival |        |        | Overall survival |        |        |
|-----------------------------|--------------------|-----------------------|--------|--------|------------------|--------|--------|
|                             |                    | Hazard ratio*         | 95% CI | p =    | Hazard ratio*    | 95% CI | p =    |
| <b>No of positive nodes</b> |                    |                       |        |        |                  |        |        |
| Overall                     | 745                | 0.80                  | 0.68-  | 0.0043 | 0.74             | 0.61-  | 0.0020 |
| 1-3                         | 467                | 0.72                  | 0.93   | 0.0047 | 0.62             | 0.90   | 0.0008 |
| 4+                          | 278                | 0.87                  | 0.58-  | 0.2290 | 0.87             | 0.46-  | 0.2746 |
|                             |                    |                       | 0.91   |        |                  | 0.82   |        |
|                             |                    |                       | 0.70-  |        |                  | 0.67-  |        |
|                             |                    |                       | 1.09   |        |                  | 1.12   |        |

\*a hazard ratio of less than 1 indicates that TAC is associated with a longer disease-free survival and overall survival compared to FAC

Patients with operable node-negative breast cancer eligible to receive chemotherapy (GEICAM 9805)

Data from a multi-centre open label randomised trial support the use of Docetaxel for the adjuvant treatment of patients with operable node-negative breast cancer eligible to receive chemotherapy.

1060 patients were randomised to receive either Docetaxel 75 mg/m<sup>2</sup> administered 1-hour after doxorubicin 50 mg/m<sup>2</sup> and cyclophosphamide 500 mg/m<sup>2</sup> (539 patients in TAC arm), or doxorubicin 50 mg/m<sup>2</sup> followed by fluorouracil 500 mg/m<sup>2</sup> and cyclophosphamide 500 mg/m<sup>2</sup> (521 patients in FAC arm), as adjuvant treatment of operable node-negative breast cancer patients with high risk of relapse according to 1998 St. Gallen criteria (tumour size >2 cm and/or negative ER and PR and/or high histological/nuclear grade (grade 2 to 3) and /or age <35 years). Both regimens were administered once every 3 weeks for 6 cycles. TAXOTERE was administered as a 1-hour infusion, all other medicinal products were given intravenously on day 1 every three weeks. Primary prophylactic G-CSF was made mandatory in TAC arm after 230 patients were randomised. The incidence of Grade 4 neutropenia, febrile neutropenia and neutropenic infection was decreased in patients who received primary G-CSF prophylaxis (see section 4.8). In both arms, after the last cycle of chemotherapy, patients with ER+ and/or PgR+ tumours received tamoxifen 20 mg once a day for up to 5 years. Adjuvant radiation therapy was administered according to guidelines in place at participating institutions and was given to 57.3% of patients who received TAC and 51.2% of patients who received FAC.

One primary analysis and one updated analysis were performed. The primary analysis was done when all patients had a follow-up of greater than 5 years (median follow-up time of 77 months). The updated analysis was performed when all patients had reached their 10-year (median follow up time of 10 years and 5 months) follow-up visit (unless they had a DFS event or were lost to follow-up previously). Disease-free survival (DFS) was the primary efficacy endpoint and Overall survival (OS) was the secondary efficacy endpoint.

At the median follow-up time of 77 months, significantly longer disease-free survival for the TAC arm compared to the FAC arm was demonstrated. TAC-treated patients had a 32% reduction in the risk of relapse compared to those treated with FAC (hazard ratio = 0.68, 95% CI (0.49-0.93), p = 0.01). At the median follow up time of 10 years and 5 months, TAC-treated patients had a 16,5% reduction in the risk of relapse compared to those treated with FAC (hazard ratio = 0.84, 95% CI (0.65-1.08), p=0.1646). DFS data were not

statistically significant but were still associated with a positive trend in favour of TAC.

At the median follow-up time of 77 months, overall survival (OS) was longer in the TAC arm with TAC-treated patients having a 24% reduction in the risk of death compared to FAC (hazard ratio = 0.76, 95% CI (0.46-1.26, p = 0.29). However, the distribution of OS was not significantly different between the 2 groups.

At the median follow up time of 10 years and 5 months, TAC-treated patients had a 9% reduction in the risk of death compared to FAC-treated patients (hazard ratio = 0.91, 95% CI (0.63-1.32)). The survival rate was 93.7% in the TAC arm and 91.4 % in the FAC arm, at the 8-year follow-up time point, and 91.3 % in the TAC arm and 89 % in the FAC arm, at the 10-year follow-up time point.

The positive benefit risk ratio for TAC compared to FAC remained unchanged.

TAC-treated patient subsets according to prospectively defined major prognostic factors were analysed in the primary analysis (at the median follow-up time of 77 months) (see table below):

Subset Analyses-Adjuvant Therapy in Patients with Node-negative Breast Cancer Study (Intent-toTreat Analysis)

| Patient subset                  | Number of patients in TAC group | Disease Free Survival |           |
|---------------------------------|---------------------------------|-----------------------|-----------|
|                                 |                                 | Hazard ratio*         | 95% CI    |
| <b>Overall</b>                  | 539                             | 0.68                  | 0.49-0.93 |
| <b>Age category 1</b>           |                                 |                       |           |
| <50 years                       | 260                             | 0.67                  | 0.43-1.05 |
| ≥50 years                       | 279                             | 0.67                  | 0.43-1.05 |
| <b>Age category 2</b>           |                                 |                       |           |
| <35 years                       | 42                              | 0.31                  | 0.11-0.89 |
| ≥35 years                       | 497                             | 0.73                  | 0.52-1.01 |
| <b>Hormonal receptor status</b> |                                 |                       |           |
| Negative                        | 195                             | 0.7                   | 0.45-1.1  |
|                                 | 344                             | 0.62                  | 0.4-0.97  |

|                                       |     |      |           |
|---------------------------------------|-----|------|-----------|
| Positive                              |     |      |           |
| <b>Tumour size</b>                    |     |      |           |
| ≤2 cm                                 | 285 | 0.69 | 0.43-1.1  |
| >2 cm                                 | 254 | 0.68 | 0.45-1.04 |
| <b>Histological grade</b>             |     |      |           |
| Grade 1 (includes grade not assessed) | 64  | 0.79 | 0.24-2.6  |
| Grade 2                               | 216 | 0.77 | 0.46-1.3  |
| Grade 3                               | 259 | 0.59 | 0.39-0.9  |
| <b>Menopausal status</b>              |     |      |           |
| Pre-Menopausal                        | 285 | 0.64 | 0.40-1    |
| Post-Menopausal                       | 254 | 0.72 | 0.47-1.12 |

\*a hazard ratio (TAC/FAC) of less than 1 indicates that TAC is associated with a longer disease free survival compared to FAC.

Exploratory subgroup analyses for disease-free survival for patients who meet the 2009 St. Gallen chemotherapy criteria – (ITT population) were performed and presented here below:

|                                                           | <b>TAC</b>        | <b>FAC</b>        | <b>Hazard ratio (TAC/FAC)</b> |                |
|-----------------------------------------------------------|-------------------|-------------------|-------------------------------|----------------|
| <b>Subgroups</b>                                          | (n=539)           | (n=521)           | (95% CI)                      | <b>p-value</b> |
| Meeting relative indication for chemotherapy <sup>a</sup> |                   |                   |                               |                |
| No                                                        | 18/214<br>(8.4%)  | 26/227<br>(11.5%) | 0.796 (0.434 - 1.459)         | 0.4593         |
| Yes                                                       | 48/325<br>(14.8%) | 69/294<br>(23.5%) | 0.606 (0.42 - 0.877)          | 0.0072         |

TAC = docetaxel, doxorubicin and cyclophosphamide  
FAC = 5-fluorouracil, doxorubicin and cyclophosphamide  
CI = confidence interval; ER =  
oestrogen receptor PR = progesterone  
receptor <sup>a</sup> ER/PR-negative or Grade 3  
or tumour size >5 cm

The estimated hazard ratio was using Cox proportional hazard model with treatment group as the factor.

#### *Docetaxel as single agent*

Two randomised phase III comparative studies, involving a total of 326 alkylating or 392 anthracycline failure metastatic breast cancer patients, have been performed with docetaxel at the recommended dose and regimen of 100 mg/m<sup>2</sup> every 3 weeks.

In alkylating-failure patients, docetaxel was compared to doxorubicin (75 mg/m<sup>2</sup> every 3 weeks).

Without affecting overall survival time (docetaxel 15 months vs. doxorubicin 14 months,  $p = 0.38$ ) or time to progression (docetaxel 27 weeks vs. doxorubicin 23 weeks,  $p = 0.54$ ), docetaxel increased response rate (52% vs. 37%,  $p = 0.01$ ) and shortened time to response (12 weeks vs. 23 weeks,  $p = 0.007$ ). Three docetaxel patients (2%) discontinued the treatment due to fluid retention, whereas 15 doxorubicin patients (9%) discontinued due to cardiac toxicity (three cases of fatal congestive heart failure).

In anthracycline-failure patients, docetaxel was compared to the combination of mitomycin C and vinblastine (12 mg/m<sup>2</sup> every 6 weeks and 6 mg/m<sup>2</sup> every 3 weeks). Docetaxel increased response rate (33% vs. 12%,  $p < 0.0001$ ), prolonged time to progression (19 weeks vs. 11 weeks,  $p = 0.0004$ ) and prolonged overall survival (11 months vs. 9 months,  $p = 0.01$ ).

During these two phase III studies, the safety profile of docetaxel was consistent with the safety profile observed in phase II studies (see section 4.8).

An open-label, multi-centre, randomised phase III study was conducted to compare docetaxel monotherapy and paclitaxel in the treatment of advanced

breast cancer in patients whose previous therapy should have included an anthracycline. A total of 449 patients were randomised to receive either docetaxel monotherapy 100 mg/m<sup>2</sup> as a 1 hour infusion or paclitaxel 175 mg/m<sup>2</sup> as a 3 hour infusion. Both regimens were administered every 3 weeks.

Without affecting the primary endpoint, overall response rate (32% vs 25%,  $p = 0.10$ ), docetaxel prolonged median time to progression (24.6 weeks vs 15.6 weeks;  $p < 0.01$ ) and median survival (15.3 months vs 12.7 months;  $p = 0.03$ ). More grade 3/4 adverse events were observed for docetaxel monotherapy (55.4%) compared to paclitaxel (23.0%).

#### *Docetaxel in combination with doxorubicin*

One large randomised phase III study, involving 429 previously untreated patients with metastatic disease, has been performed with doxorubicin (50 mg/m<sup>2</sup>) in combination with docetaxel (75 mg/m<sup>2</sup>) (AT arm) versus doxorubicin (60 mg/m<sup>2</sup>) in combination with cyclophosphamide (600 mg/m<sup>2</sup>) (AC arm). Both regimens were administered on day 1 every 3 weeks.

- Time to progression (TTP) was significantly longer in the AT arm versus AC arm,  $p = 0.0138$ . The median TTP was 37.3 weeks (95% CI: 33.4 - 42.1) in AT arm and 31.9 weeks (95% CI: 27.4 - 36.0) in AC arm.
- Overall response rate (ORR) was significantly higher in the AT arm versus AC arm,  $p = 0.009$ . The ORR was 59.3% (95% CI: 52.8 - 65.9) in AT arm versus 46.5% (95% CI: 39.8 - 53.2) in AC arm.

In this study, AT arm showed a higher incidence of severe neutropenia (90% versus 68.6%), febrile neutropenia (33.3% versus 10%), infection (8% versus 2.4%), diarrhoea (7.5% versus 1.4%), asthenia (8.5% versus 2.4%), and pain (2.8% versus 0%) than AC arm. On the other hand, AC arm showed a higher incidence of severe anaemia (15.8% versus 8.5%) than AT arm, and, in addition, a higher incidence of severe cardiac toxicity: congestive heart failure (3.8% versus 2.8%), absolute LVEF decrease  $\geq 20\%$  (13.1% versus 6.1%), absolute LVEF decrease  $\geq 30\%$  (6.2% versus 1.1%). Toxic deaths occurred in 1 patient in the AT arm (congestive heart failure) and in 4 patients in the AC arm (1 due to septic shock and 3 due to congestive heart failure).

In both arms, quality of life measured by the EORTC questionnaire was comparable and stable during treatment and follow-up.

#### *Docetaxel in combination with trastuzumab*

Docetaxel in combination with trastuzumab was studied for the treatment of patients with metastatic breast cancer whose tumours over express HER2, and who previously had not received chemotherapy for metastatic disease. One hundred eighty six patients were randomised to receive docetaxel (100 mg/m<sup>2</sup>) with or without trastuzumab; 60% of patients received prior anthracycline-based adjuvant chemotherapy. Docetaxel plus trastuzumab was efficacious in patients whether or not they had received prior adjuvant anthracyclines. The main test method used to determine HER2 positivity in this pivotal study was immunohistochemistry (IHC). A minority of patients were tested using fluorescence in-situ hybridization (FISH). In this study, 87% of patients had disease that was IHC 3+, and 95% of patients entered had disease that was IHC 3+ and/or FISH positive. Efficacy results are summarized in the following table:

| Parameter                                              | Docetaxel plus<br>trastuzumab<br>n = 92 | Docetaxel<br>n = 94              |
|--------------------------------------------------------|-----------------------------------------|----------------------------------|
| Response rate<br>(95% CI)                              | 61%<br>(50-71)                          | 34%<br>(25-45)                   |
| Median duration of<br>response<br>(months)<br>(95% CI) | 11.4<br>(9.2-15.0)                      | 5.1<br>(4.4-6.2)                 |
| Median TTP<br>(months) (95% CI)                        | 10.6<br>(7.6-12.9)                      | 5.7<br>(5.0-6.5)                 |
| Median survival<br>(months) (95% CI)                   | 30.5 <sup>2</sup><br>(26.8-ne)          | 22.1 <sup>2</sup><br>(17.6-28.9) |

TTP = time to progression; “ne” indicates that it could not be estimated or it was not yet reached.

<sup>1</sup>Full analysis set (intent-to-treat)

2 Estimated median survival

### Docetaxel in combination with capecitabine

Data from one multi-centre, randomized, controlled phase III clinical study support the use of docetaxel in combination with capecitabine for treatment of patients with locally advanced or metastatic breast cancer after failure of cytotoxic chemotherapy, including an anthracycline. In this study, 255 patients were randomized to treatment with docetaxel (75 mg/m<sup>2</sup> as a 1 hour intravenous

infusion every 3 weeks) and capecitabine (1250 mg/m<sup>2</sup> twice daily for 2 weeks followed by 1-week rest period). 256 patients were randomised to treatment with docetaxel alone (100 mg/m<sup>2</sup> as a 1 hour intravenous infusion every 3 weeks). Survival was superior in the docetaxel + capecitabine combination arm (p = 0.0126). Median survival was 442 days (docetaxel + capecitabine) vs. 352 days (docetaxel alone). The overall objective response rates in the all-randomised population (investigator assessment) were 41.6% (docetaxel + capecitabine) vs. 29.7% (docetaxel alone); p = 0.0058. Time to progressive disease was superior in the docetaxel + capecitabine combination arm (p < 0.0001). The median time to progression was 186 days (docetaxel + capecitabine) vs. 128 days (docetaxel alone).

### *Non-small cell lung cancer*

#### *Patients previously treated with chemotherapy with or without radiotherapy*

In a phase III study, in previously treated patients, time to progression (12.3 weeks versus 7 weeks) and overall survival were significantly longer for docetaxel at 75 mg/m<sup>2</sup> compared to Best Supportive Care. The 1-year survival rate was also significantly longer in docetaxel (40%) versus BSC (16%). There was less use of morphinic analgesic (p < 0.01), non-morphinic analgesics (p < 0.01), other disease-related medications (p = 0.06) and radiotherapy (p < 0.01) in patients treated with docetaxel at 75 mg/m<sup>2</sup> compared to those with BSC.

The overall response rate was 6.8% in the evaluable patients, and the median duration of response was 26.1 weeks.

*Docetaxel in combination with platinum agents in chemotherapy-naïve patients*

In a phase III study, 1218 patients with unresectable stage IIIB or IV NSCLC, with KPS of 70% or greater, and who did not receive previous chemotherapy for this condition, were randomised to either docetaxel (T) 75 mg/m<sup>2</sup> as a 1 hour infusion immediately followed by cisplatin (Cis) 75 mg/m<sup>2</sup> over 30-60 minutes every 3 weeks (TCis), docetaxel 75 mg/m<sup>2</sup> as a 1 hour infusion in combination with carboplatin (AUC 6 mg/ml.min) over 30-60 minutes every 3 weeks, or vinorelbine (V) 25 mg/m<sup>2</sup> administered over 6-10 minutes on days 1, 8, 15, 22 followed by cisplatin 100 mg/m<sup>2</sup> administered on day 1 of cycles repeated every 4 weeks (VCis).

Survival data, median time to progression and response rates for two arms of the study are illustrated in the following table:

|                                          | TCis<br>n =<br>408 | VCis<br>n =<br>404 | Statistical analysis                                                 |
|------------------------------------------|--------------------|--------------------|----------------------------------------------------------------------|
| Overall survival<br>(Primary end-point): |                    |                    |                                                                      |
| Median survival<br>(months)              | 11.3               | 10.1               | Hazard ratio: 1.122<br>[97.2% CI: 0.937; 1.342]*                     |
| 1-year Survival<br>(%)                   | 46                 | 41                 | Treatment difference:<br>5.4%                                        |
| 2-year Survival<br>(%)                   | 21                 | 14                 | [95% CI: -1.1; 12.0]<br>Treatment difference:<br>6.2%                |
| Median time to<br>progression (weeks):   | 22.0               | 23.0               | [95% CI: 0.2; 12.3]<br>Hazard ratio: 1.032<br>[95% CI: 0.876; 1.216] |
| Overall response rate (%):               | 31.6               | 24.5               | Treatment difference:<br>7.1% [95% CI: 0.7; 13.5]                    |

\*: Corrected for multiple comparisons and adjusted for stratification factors (stage of disease and region of treatment), based on evaluable patient population.

Secondary end-points included change of pain, global rating of quality of life by EuroQoL-5D, Lung Cancer Symptom Scale, and changes in Karnofsky performance status. Results on these end-points were supportive of the primary end-points results.

For docetaxel/carboplatin combination, neither equivalent nor non-inferior efficacy could be proven compared to the reference treatment combination VCis.

### Prostate cancer

#### Metastatic castration-resistant prostate cancer

The safety and efficacy of docetaxel in combination with prednisone or prednisolone in patients with metastatic castration-resistant prostate cancer were evaluated in a randomised multi-centre phase III study (TAX 327). A total of 1006 patients with KPS  $\geq$  60 were randomised to the following treatment groups:

- Docetaxel 75 mg/m<sup>2</sup> every 3 weeks for 10 cycles.
- Docetaxel 30 mg/m<sup>2</sup> administered weekly for the first 5 weeks in a 6 week cycle for 5 cycles.
- Mitoxantrone 12 mg/m<sup>2</sup> every 3 weeks for 10 cycles.

All 3 regimens were administered in combination with prednisone or prednisolone 5 mg twice daily, continuously.

Patients who received docetaxel every three weeks demonstrated significantly longer overall survival compared to those treated with mitoxantrone. The increase in survival seen in the docetaxel weekly arm was not statistically significant compared to the mitoxantrone control arm. Efficacy endpoints for the docetaxel arms versus the control arm are summarized in the following table:

| Endpoint | Docetaxel<br>every 3<br>weeks | Docetaxel every<br>week | Mitoxantrone<br>every 3<br>weeks |
|----------|-------------------------------|-------------------------|----------------------------------|
|----------|-------------------------------|-------------------------|----------------------------------|

|                          |                       |                       |                     |
|--------------------------|-----------------------|-----------------------|---------------------|
| Number of patients       | 335                   | 334                   | 337                 |
| Median survival (months) | 18.9<br>(17.0-21.2)   | 17.4<br>(15.7-19.0)   | 16.5<br>(14.4-18.6) |
| 95% CI                   | 0.761                 | 0.912                 | --                  |
| Hazard ratio             | (0.619-0.936)         | (0.747-1.113)         | --                  |
| 95% CI p-value†*         | 0.0094                | 0.3624                | --                  |
| Number of patients       | 291                   | 282                   | 300                 |
| PSA** response rate (%)  | 45.4<br>(39.5-51.3)   | 47.9<br>(41.9-53.9)   | 31.7<br>(26.4-37.3) |
| 95% CI p-value*          | 0.0005                | <0.0001               | --                  |
| Number of patients       | 153                   | 154                   | 157                 |
| Pain response rate (%)   | 34.6                  | 31.2                  | 21.7                |
| 95% CI p-value*          | (27.1-42.7)<br>0.0107 | (24.0-39.1)<br>0.0798 | (15.5-28.9)<br>--   |
| Number of patients       | 141                   | 134                   | 137                 |
| Tumour response rate (%) | 12.1<br>(7.2-18.6)    | 8.2<br>(4.2-14.2)     | 6.6<br>(3.0-12.1)   |
| 95% CI p-value*          | 0.1112                | 0.5853                | --                  |

†Stratified log-rank test

\*Threshold for statistical significance = 0.0175

\*\*PSA: Prostate-Specific Antigen

Given the fact that docetaxel every week presented a slightly better safety profile than docetaxel every 3 weeks, it is possible that certain patients may benefit from docetaxel every week.

No statistical differences were observed between treatment groups for Global Quality of Life.

#### Metastatic hormone-sensitive prostate cancer

##### *STAMPEDE study*

The safety and efficacy of docetaxel administered concomitantly with standard of care (ADT) in patients with high-risk locally advanced or metastatic hormone-sensitive prostate cancer were evaluated in a randomised multi-centre, multi-arm multi-stage (MAMS) study with a seamless phase II/III

design (STAMPEDE – MRC PR08). A total of 1776 male patients were allocated to the treatment arms of interest:

- Standard of care + docetaxel 75 mg/m<sup>2</sup>, administered every 3 weeks for 6 cycles
- Standard of care alone

Docetaxel regimen was administered in combination with prednisone or prednisolone 5 mg twice daily continuously.

Among the 1776 randomised patients 1086 (61%) had metastatic disease, 362 were randomised to docetaxel in combination with standard of care, 724 received standard of care alone.

In these metastatic prostate cancer patients, the median overall survival was significantly longer in docetaxel treatment groups than in the standard of care alone group, with a median overall survival 19 months longer with the addition of docetaxel to standard of care (HR = 0.76, 95% CI = 0.62-0.92, p=0.005).

Efficacy results in metastatic prostate cancer patients for docetaxel arm versus control arm are summarized in the following table:

Efficacy of docetaxel in combination with prednisone or prednisolone and standard of care in the treatment of patients with metastatic hormone-sensitive prostate cancer (STAMPEDE)

| Endpoint                                      | Docetaxel + standard of care | Standard of care alone |
|-----------------------------------------------|------------------------------|------------------------|
| Number of metastatic prostate cancer patients | 362                          | 724                    |
| Median overall survival (months)              | 62<br>51-73                  | 43<br>40-48            |
| 95% CI                                        | 0.76                         |                        |
| Adjusted hazard ratio                         | (0.62-0.92)                  |                        |
| 95% CI                                        | 0.005                        |                        |
| p-value <sup>a</sup>                          |                              |                        |
| Failure-Free survival <sup>b</sup>            |                              |                        |
| Median (months)                               | 20.4<br>16.8-25.2            | 12<br>9.6-12           |

|                       |             |
|-----------------------|-------------|
| 95% CI                | 0.66        |
| Adjusted hazard ratio | (0.57-0.76) |
| 95% CI                | <0.001      |
| p-value <sup>a</sup>  |             |

<sup>a</sup> p-value calculated from the likelihood ratio test and adjusted for all stratification factors (except center and planned hormone therapy) and stratified by trial period

<sup>b</sup> Failure-free survival: time from randomization to first evidence of at least one of: biochemical failure (defined as a rise in PSA of 50% above the within-24-week nadir and above 4 ng/mL and confirmed by retest or treatment); progression either locally, in lymph nodes, or in distant metastases; skeletal-related event; or death from prostate cancer.

#### *CHAARTED study*

The safety and efficacy of docetaxel administered at the beginning of androgen-deprivation therapy (ADT) in patients with metastatic hormone-sensitive prostate cancer were evaluated in a randomized Phase III multi-centre study (CHAARTED). A total of 790 male patients were allocated to the 2 treatment groups.

- ADT + docetaxel 75 mg/m<sup>2</sup> given at the beginning of ADT, administered every 3 weeks for 6 cycles
- ADT alone

The median overall survival was significantly longer in docetaxel treatment group than in the ADT alone group, with a median overall survival 13.6 months longer with the addition of docetaxel to ADT (hazard ratio (HR) = 0.61, 95% confidence interval (CI) = 0.47-0.80, p=0.0003).

Efficacy results of the docetaxel arm versus the control arm are summarized in the following table:

Efficacy of docetaxel and ADT in the treatment of patients with metastatic hormone-sensitive prostate cancer (CHAARTED)

| Endpoint | Docetaxel +ADT | ADT alone |
|----------|----------------|-----------|
|----------|----------------|-----------|

|                                                            |                        |                 |
|------------------------------------------------------------|------------------------|-----------------|
| Number of patients                                         | 397                    | 393             |
| Median overall survival (months) All patients              | 57.6                   | 44.0            |
| 95% CI                                                     | 49.1-72.8              | 34.4-49.1       |
| Adjusted hazard ratio                                      | 0.61                   | --              |
| 95% CI                                                     | (0.47-0.80)            | --              |
| p-value <sup>a</sup>                                       | 0.0003                 | --              |
| Progression Free Survival                                  |                        |                 |
| Median (months)                                            | 19.8                   | 11.6            |
| 95% CI                                                     | 16.7-22.8              | 10.8-14.3       |
| Adjusted hazard ratio                                      | 0.60                   | --              |
| 95% CI p-value*                                            | 0.51-0.72              | --              |
|                                                            | P<0.0001               | --              |
| PSA response** at 6 months – N (%) p-value <sup>a</sup> *  | 127 (32.0)<br><0.0001  | 77 (19.6)<br>-- |
| PSA response** at 12 months – N (%) p-value <sup>a</sup> * | 110 (27.7)<br><0.0001  | 66 (16.8)<br>-- |
| Time to castration-resistant prostate cancer <sup>b</sup>  |                        |                 |
| Median (months)                                            | 20.2                   | 11.7            |
| 95% CI                                                     | (17.2-23.6)            | (10.8-14.7)     |
| Adjusted hazard ratio                                      | 0.61                   | --              |
| 95% CI p-value <sup>a</sup> *                              | (0.51-0.72)<br><0.0001 | --<br>--        |
| Time to clinical progression <sup>c</sup>                  |                        |                 |
| Median (months)                                            | 33.0                   | 19.8            |
| 95% CI                                                     | (27.3-41.2)            | (17.9-22.8)     |
| Adjusted hazard ratio                                      | 0.61                   | --              |
| 95% CI p-value <sup>a</sup> *                              | (0.50-0.75)<br><0.0001 | --<br>--        |

<sup>a</sup> Time to event variables: Stratified log-rank test. Response rate variables: Fisher's Exact test \* p-value for descriptive purpose.

\*\* PSA response: Prostate-Specific Antigen response: PSA level <0.2 ng/mL measured for two consecutive measurements at least 4 weeks apart.

<sup>b</sup> Time to castration-resistant prostate cancer = time from randomization to PSA progression or clinical progression (i.e., increasing symptomatic bone

metastases, progression per Response Evaluation Criteria in Solid Tumours (RECIST) criteria, or clinical deterioration due to cancer per the Investigator's opinion), whichever occurred first.

<sup>c</sup> The time to clinical progression = the time from randomization until clinical progression (i.e., increased symptoms of bone metastases; progression according to RECIST; or clinical deterioration due to cancer according to the investigator's opinion).

#### Gastric adenocarcinoma

A multi-centre, open-label, randomized study was conducted to evaluate the safety and efficacy of docetaxel for the treatment of patients with metastatic gastric adenocarcinoma, including adenocarcinoma of the gastro esophageal junction, who had not received prior chemotherapy for metastatic disease. A total of 445 patients with KPS > 70 were treated with either docetaxel (T) (75 mg/m<sup>2</sup> on day 1) in combination with cisplatin (C) (75 mg/m<sup>2</sup> on day 1) and 5-fluorouracil (F) (750 mg/m<sup>2</sup> per day for 5 days) or cisplatin (100 mg/m<sup>2</sup> on day 1) and 5-fluorouracil (1000 mg/m<sup>2</sup> per day for 5 days). The length of a treatment cycle was 3 weeks for the TCF arm and 4 weeks for the CF arm. The median number of cycles administered per patient was 6 (with a range of 1-16) for the TCF arm compared to 4 (with a range of 1-12) for the CF arm. Time to progression (TTP) was the primary endpoint. The risk reduction of progression was 32.1% and was associated with a significantly longer TTP (p = 0.0004) in favour of the TCF arm. Overall survival was also significantly longer (p = 0.0201) in favour of the TCF arm with a risk reduction of mortality of 22.7%. Efficacy results are summarized in the following table:

#### Efficacy of docetaxel in the treatment of patients with gastric adenocarcinoma

| <b>Endpoint</b>     | <b>TCF<br/>n = 221</b> | <b>CF<br/>n = 224</b> |
|---------------------|------------------------|-----------------------|
| Median TTP (months) | 5.6                    | 3.7                   |
| 3.7                 | (4.86-5.91)            | (3.45-4.47)           |
| (95% CI)            | 1.473                  |                       |
| Hazard ratio        | (1.189-                |                       |
| (95% CI)            | 1.825)                 |                       |

|                                                     |                                  |                    |
|-----------------------------------------------------|----------------------------------|--------------------|
| *p-value                                            | 0.0004                           |                    |
| Median survival (months)<br>(95% CI)                | 9.2<br>(8.38-10.58)              | 8.6<br>(7.16-9.46) |
| 2-year estimate (%)                                 | 18.4                             | 8.8                |
| Hazard ratio<br>(95% CI)<br>*p-value                | 1.293<br>(1.041-1.606)<br>0.0201 |                    |
| Overall response rate<br>(CR+PR) (%) p-value        | 36.7                             | 25.4               |
|                                                     | 0.0106                           |                    |
| Progressive disease as best<br>overall Response (%) | 16.7                             | 25.9               |

\*Unstratified logrank test

Subgroup analyses across age, gender and race consistently favoured the TCF arm compared to the CF arm.

A survival update analysis conducted with a median follow-up time of 41.6 months no longer showed a statistically significant difference although always in favour of the TCF regimen and showed that the benefit of TCF over CF is clearly observed between 18 and 30 months of follow up.

Overall, quality of life (QoL) and clinical benefit results consistently indicated improvement in favour of the TCF arm. Patients treated with TCF had a longer time to 5% definitive deterioration of global health status on the QLQ-C30 questionnaire (p = 0.0121) and a longer time to definitive worsening of Karnofsky performance status (p = 0.0088) compared to patients treated with CF.

### Head and neck cancer

- Induction chemotherapy followed by radiotherapy (TAX323)

The safety and efficacy of docetaxel in the induction treatment of patients with squamous cell carcinoma of the head and neck (SCCHN) was evaluated in a

phase III, multicenter, open-label, randomised study (TAX323). In this study, 358 patients with inoperable locally advanced SCCHN, and WHO performance status 0 or 1, were randomised to one of two treatment arms. Patients on the docetaxel arm received docetaxel (T) 75 mg/m<sup>2</sup> followed by cisplatin (P) 75 mg/m<sup>2</sup> followed by 5fluorouracil (F) 750 mg/m<sup>2</sup> per day as a continuous infusion for 5 days. This regimen was administered

every three weeks for 4 cycles in case at least a minor response ( $\geq 25\%$  reduction in bidimensionally measured tumour size) was observed after 2 cycles. At the end of chemotherapy, with a minimal interval of 4 weeks and a maximal interval of 7 weeks, patients whose disease did not progress received radiotherapy (RT) according to institutional guidelines for 7 weeks (TPF/RT). Patients on the comparator arm received cisplatin (P) 100 mg/m<sup>2</sup> followed by 5-fluorouracil (F) 1000 mg/m<sup>2</sup> per day for 5 days. This regimen was administered every three weeks for 4 cycles in case at least a minor

response ( $\geq 25\%$  reduction in bidimensionally measured tumour size) was observed after 2 cycles. At the end of chemotherapy, with a minimal interval of 4 weeks and a maximal interval of 7 weeks, patients whose disease did not progress received radiotherapy (RT) according to institutional guidelines for 7 weeks (PF/RT). Locoregional therapy with radiation was delivered either with a conventional fraction (1.8 Gy - 2.0 Gy once a day, 5 days per week for a total dose of 66 to 70 Gy), or accelerated/hyperfractionated regimens of radiation therapy (twice a day, with a minimum interfraction interval of 6 hours, 5 days per week). A total of 70 Gy was recommended for accelerated regimens and 74 Gy for hyperfractionated schemes. Surgical resection was allowed following chemotherapy, before or after radiotherapy. Patients on the TPF arm received antibiotic prophylaxis with ciprofloxacin 500 mg orally twice daily for 10 days starting on day 5 of each cycle, or equivalent. The primary endpoint in this study, progression-free survival (PFS), was significantly longer in the TPF arm compared to the PF arm,  $p = 0.0042$  (median PFS: 11.4 vs. 8.3 months respectively) with an overall median follow up time of 33.7 months. Median overall survival was also significantly longer in favour of the TPF arm compared to the PF arm (median OS: 18.6 vs. 14.5 months respectively) with a 28% risk reduction of mortality,  $p = 0.0128$ . Efficacy results are presented in the table below:

Efficacy of docetaxel in the induction treatment of patients with inoperable locally advanced SCCHN (Intent-to-Treat Analysis)

| <b>Endpoint</b>                                                                    | <b>Docetaxel +<br/>Cis + 5-<br/>FU n =<br/>177</b> | <b>Cis + 5-<br/>FU n =<br/>181</b> |
|------------------------------------------------------------------------------------|----------------------------------------------------|------------------------------------|
| Median progression free survival<br>(months) (95% CI)                              | 11.4<br>(10.1-14.0)                                | 8.3<br>(7.4-9.1)                   |
| Adjusted hazard ratio<br>(95% CI)                                                  | 0.70<br>(0.55-0.89)                                |                                    |
| *p-value                                                                           | 0.0042                                             |                                    |
| Median survival (months)<br>(95% CI)                                               | 18.6<br>(15.7-24.0)                                | 14.5<br>(11.6-18.7)                |
| Hazard ratio<br>(95% CI)                                                           | 0.72<br>(0.56-0.93)                                |                                    |
| **p-value                                                                          | 0.0128                                             |                                    |
| Best overall response to<br>chemotherapy (%) (95% CI)                              | 67.8<br>(60.4-74.6)                                | 53.6<br>(46.0-61.0)                |
| ***p-value                                                                         | 0.006                                              |                                    |
| Best overall response to study<br>treatment<br>[chemotherapy +/- radiotherapy] (%) | 72.3<br>(65.1-78.8)                                | 58.6<br>(51.0-65.8)                |
| (95% CI)                                                                           | 0.006                                              |                                    |
| ***p-value                                                                         |                                                    |                                    |
| Median duration of response to<br>chemotherapy ± radiotherapy<br>(months)          | n = 128<br>15.7<br>(13.4-24.6)                     | n = 106<br>11.7<br>(10.2-17.4)     |
| (95% CI)                                                                           | 0.72<br>(0.52-0.99)                                |                                    |
| Hazard ratio<br>(95% CI)                                                           | 0.0457                                             |                                    |
| **p-value                                                                          |                                                    |                                    |

A hazard ratio of less than 1 favours docetaxel + cisplatin + 5-FU

\*Cox model (adjustment for Primary tumour site, T and N clinical stages and PSWHO)

\*\*Log rank test

\*\*\* Chi-square test

### *Quality of life parameters*

Patients treated with TPF experienced significantly less deterioration of their Global health score compared to those treated with PF ( $p = 0.01$ , using the EORTC QLQ-C30 scale).

### *Clinical benefit parameters*

The performance status scale, for head and neck (PSS-HN) subscales designed to measure understandability of speech, ability to eat in public, and normalcy of diet, was significantly in favour of TPF as compared to PF.

Median time to first deterioration of WHO performance status was significantly longer in the TPF arm compared to PF. Pain intensity score improved during treatment in both groups indicating adequate pain management.

- Induction chemotherapy followed by chemoradiotherapy (TAX324)

The safety and efficacy of docetaxel in the induction treatment of patients with locally advanced squamous cell carcinoma of the head and neck (SCCHN) was evaluated in a randomised, multi-centre open-label, phase III study (TAX324). In this study, 501 patients, with locally advanced SCCHN, and a WHO performance status of 0 or 1, were randomised to one of two arms. The study population comprised patients with technically unresectable disease, patients with low probability of surgical cure and patients aiming at organ preservation. The efficacy and safety evaluation solely addressed survival endpoints and the success of organ preservation was not formally addressed. Patients on the docetaxel arm received docetaxel (T) 75 mg/m<sup>2</sup> by intravenous infusion on day 1 followed by cisplatin (P) 100 mg/m<sup>2</sup> administered as a 30-minute to three-hour intravenous infusion, followed by the continuous intravenous infusion of 5-fluorouracil (F) 1000 mg/m<sup>2</sup>/day from day 1 to day 4. The cycles were repeated every 3 weeks for 3 cycles. All patients who did not have progressive disease were to receive chemoradiotherapy (CRT) as per protocol (TPF/CRT). Patients on the comparator arm received cisplatin (P) 100 mg/m<sup>2</sup> as a 30-minute to three-hour intravenous infusion on day 1 followed by the continuous intravenous infusion of 5-fluorouracil (F) 1000 mg/m<sup>2</sup>/day from day 1 to day 5. The cycles were repeated every 3 weeks for 3 cycles. All patients who did not have progressive disease were to receive CRT as per protocol (PF/CRT).

Patients in both treatment arms were to receive 7 weeks of CRT following induction chemotherapy with a minimum interval of 3 weeks and no later than

8 weeks after start of the last cycle (day 22 to day 56 of last cycle). During radiotherapy, carboplatin (AUC 1.5) was given weekly as a one-hour intravenous infusion for a maximum of 7 doses. Radiation was delivered with megavoltage equipment using once daily fractionation (2 Gy per day, 5 days per week for 7 weeks, for a total dose of 70-72 Gy). Surgery on the primary site of disease and/or neck could be considered at any time following completion of CRT. All patients on the docetaxel-containing arm of the study received prophylactic antibiotics. The primary efficacy endpoint in this study, overall survival (OS) was significantly longer (log-rank test,  $p = 0.0058$ ) with the docetaxel-containing regimen compared to PF (median OS: 70.6 versus 30.1 months respectively), with a 30% risk reduction in mortality compared to PF (hazard ratio (HR) = 0.70, 95% confidence interval (CI) = 0.54-0.90) with an overall median follow up time of 41.9 months. The secondary endpoint, PFS, demonstrated a 29% risk reduction of progression or death and a 22 month improvement in median PFS (35.5 months for TPF and 13.1 for PF). This was also statistically significant with an HR of 0.71; 95% CI 0.56-0.90; log-rank test  $p = 0.004$ . Efficacy results are presented in the table below:

Efficacy of docetaxel in the induction treatment of patients with locally advanced SCCHN (Intent-to-Treat Analysis)

| <b>Endpoint</b>                                 | <b>Docetaxel + Cis +<br/>5-FU n = 255</b> | <b>Cis + 5-<br/>FU n =<br/>246</b> |
|-------------------------------------------------|-------------------------------------------|------------------------------------|
| Median overall survival<br>(months)<br>(95% CI) | 70.6<br>(49.0-NA)                         | 30.1<br>(20.9-<br>51.5)            |
| Hazard ratio:<br>(95% CI)<br>*p-value           | 0.70<br>(0.54-0.90)<br>0.0058             |                                    |
| Median PFS (months)<br>(95% CI)                 | 35.5<br>(19.3-NA)                         | 13.1<br>(10.6 -<br>20.2)           |
| Hazard ratio:<br>(95% CI)<br>**p-value          | 0.71<br>(0.56 - 0.90)<br>0.004            |                                    |

|                                                                                                                 |                     |                     |
|-----------------------------------------------------------------------------------------------------------------|---------------------|---------------------|
| Best overall response (CR + PR) to chemotherapy (%) (95% CI) ***p-value                                         | 71.8<br>(65.8-77.2) | 64.2<br>(57.9-70.2) |
|                                                                                                                 | 0.070               |                     |
| Best overall response (CR + PR) to study treatment [chemotherapy +/- chemoradiotherapy] (%) (95% CI) ***p-value | 76.5<br>(70.8-81.5) | 71.5<br>(65.5-77.1) |
|                                                                                                                 | 0.209               |                     |

A hazard ratio of less than 1 favors docetaxel + cisplatin + fluorouracil

\*un-adjusted log-rank test

\*\*un-adjusted log-rank test, not adjusted for multiple comparisons \*\*\*Chi square test, not adjusted for multiple comparisons NA-not applicable

### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with TAXOTERE in all subsets of the paediatric population in breast cancer, non-small cell lung cancer, prostate cancer, gastric carcinoma and head and neck cancer, not including type II and III less differentiated nasopharyngeal carcinoma (see section 4.2 for information on paediatric use). **5.2 Pharmacokinetic properties**

### Absorption

The pharmacokinetics of docetaxel have been evaluated in cancer patients after administration of 20115 mg/m<sup>2</sup> in phase I studies. The kinetic profile of docetaxel is dose independent and consistent with a three-compartment pharmacokinetic model with half-lives for the  $\alpha$ ,  $\beta$  and  $\gamma$  phases of 4 min, 36

min and 11.1 h, respectively. The late phase is due, in part, to a relatively slow efflux of docetaxel from the peripheral compartment.

### Distribution

Following the administration of a 100 mg/m<sup>2</sup> dose given as a one-hour infusion a mean peak plasma level of 3.7 µg/ml was obtained with a corresponding AUC of 4.6 h·µg/ml. Mean values for total body clearance and steady-state volume of distribution were 21 l/h/m<sup>2</sup> and 113 l, respectively. Inter individual variation in total body clearance was approximately 50%. Docetaxel is more than 95% bound to plasma proteins.

### Elimination

A study of <sup>14</sup>C-docetaxel has been conducted in three cancer patients. Docetaxel was eliminated in both the urine and faeces following cytochrome P450-mediated oxidative metabolism of the tert-butyl ester group, within seven days, the urinary and faecal excretion accounted for about 6% and 75% of the administered radioactivity, respectively. About 80% of the radioactivity recovered in faeces is excreted during the first 48 hours as one major inactive metabolite and 3 minor inactive metabolites and very low amounts of unchanged medicinal product.

### Special populations

#### *Age and gender*

A population pharmacokinetic analysis has been performed with docetaxel in 577 patients.

Pharmacokinetic parameters estimated by the model were very close to those estimated from phase I studies. The pharmacokinetics of docetaxel were not altered by the age or sex of the patient.

#### *Hepatic impairment*

In a small number of patients (n = 23) with clinical chemistry data suggestive of mild to moderate liver function impairment (ALT, AST ≥ 1.5 times the ULN associated with alkaline phosphatase ≥ 2.5 times the ULN), total clearance was lowered by 27% on average (see section 4.2).

#### *Fluid retention*

Docetaxel clearance was not modified in patients with mild to moderate fluid retention and there are no data available in patients with severe fluid retention.

#### Combination therapy

##### *Doxorubicin*

When used in combination, docetaxel does not influence the clearance of doxorubicin and the plasma levels of doxorubicinol (a doxorubicin metabolite). The pharmacokinetics of docetaxel, doxorubicin and cyclophosphamide were not influenced by their coadministration.

##### *Capecitabine*

Phase I study evaluating the effect of capecitabine on the pharmacokinetics of docetaxel and vice versa showed no effect by capecitabine on the pharmacokinetics of docetaxel (C<sub>max</sub> and AUC) and no effect by docetaxel on the pharmacokinetics of a relevant capecitabine metabolite 5'-DFUR.

*Cisplatin* Clearance of docetaxel in combination therapy with cisplatin was similar to that observed following monotherapy. The pharmacokinetic profile of cisplatin administered shortly after docetaxel infusion is similar to that observed with cisplatin alone.

##### *Cisplatin and 5-fluorouracil*

The combined administration of docetaxel, cisplatin and 5-fluorouracil in 12 patients with solid tumours had no influence on the pharmacokinetics of each individual medicinal product.

##### *Prednisone and dexamethasone*

The effect of prednisone on the pharmacokinetics of docetaxel administered with standard dexamethasone premedication has been studied in 42 patients.

##### *Prednisone*

No effect of prednisone on the pharmacokinetics of docetaxel was observed.

### **5.3 Preclinical safety data**

The carcinogenic potential of docetaxel has not been studied.

## **DOCETAXEL INJECTION USP 120mg/3mL**

Docetaxel has been shown to be mutagenic in the in vitro micronucleus and chromosome aberration test in CHO-K1 cells and in the in vivo micronucleus test in the mouse. However, it did not induce mutagenicity in the Ames test or the CHO/HGPRT gene mutation assay. These results are consistent with the pharmacological activity of docetaxel.

Undesirable effects on the testis observed in rodent toxicity studies suggest that docetaxel may impair male fertility.

### **6. PHARMACEUTICAL PARTICULARS**

#### **6.1 List of excipients**

Concentrate vial:

Polysorbate 80

Citric acid  
anhydrous

Solvent vial:

Absolute Alcohol

Water for injections

#### **6.2 Incompatibilities**

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

#### **6.3 Shelf life**

Unopened vial

24 months (as packaged for sale)

After dilution

Premix solution: The premix solution contains 10 mg/ml docetaxel and should be used immediately after preparation. However the chemical and physical stability of the premix solution has been demonstrated for 8 hours when stored either between 2°C and 8°C or at room temperature (below 25°C).

Infusion solution: the infusion solution should be used within 4 hours at room temperature (below 25°C).

---

#### **6.4 Special precautions for storage** Store between 2°C to 8°C.

Protect from Light

---

Keep the vial in the outer carton in order to protect from light.

For storage conditions of the diluted medicinal product, see section 6.3.

#### **6.5 Nature and contents of container**

Each combipack contains:

- one single-dose vial of concentrate (3.0 mL) and,
- one single-dose vial of solvent (9.0 mL).

Docetaxel Injection USP 120 mg/3 mL concentrate for solution for infusion vial

A 20mL clear moulded glass vial USP Type I with 20 mm bromobutyl grey rubber stopper and sealed with purple colour flip off aluminum seal. This vial contains Docetaxel Injection USP 120mg/3mL concentrate for solution for infusion.

Solvent vial

A 10mL clear moulded glass vial USP Type I with 20 mm bromobutyl grey rubber stopper and sealed with orange colour flip off aluminum seal. This vial contain 9 mL Solvent for Docetaxel Injection USP 120mg/3mL concentrate for solution for infusion.

#### **6.6 Special precautions for disposal and other handling**

Docetaxel is an antineoplastic agent and, as with other potentially toxic compounds, caution should be exercised when handling it and preparing Docetaxel solutions. The use of gloves is recommended. If Docetaxel concentrate, premix solution or infusion solution should come into contact with skin, wash immediately and thoroughly with soap and water. If Docetaxel concentrate, premix solution or infusion solution should come into contact with mucous membranes, wash immediately and thoroughly with water.

Preparation for the intravenous administration

a) Preparation of the Docetaxel concentrate for solution for infusion premix solution (10 mg docetaxel/mL)

---

If the vials are stored under refrigeration, allow the required number of Docetaxel boxes to stand at room temperature (below 25°C) for 5 minutes.

Using a syringe fitted with a needle, aseptically withdraw the entire contents of the solvent for Docetaxel vial by partially inverting the vial.

Inject the entire contents of the syringe into the corresponding Docetaxel vial.

Remove the syringe and needle and mix manually by repeated inversions for at least 45 seconds. Do not shake.

Allow the premix vial to stand for 5 minutes at room temperature (below 25°C) and then check that the solution is homogenous and clear (foaming is normal even after 5 minutes due to the presence of polysorbate 80 in the formulation).

The premix solution contains 10 mg/ml docetaxel and should be used immediately after preparation. However the chemical and physical stability of the premix solution has been demonstrated for 8 hours when stored either between 2°C and 8°C or at room temperature (below 25°C).

#### b) Preparation of the infusion solution

More than one premix vial may be necessary to obtain the required dose for the patient. Based on the required dose for the patient expressed in mg, aseptically withdraw the corresponding premix volume containing 10 mg/ml docetaxel from the appropriate number of premix vials using graduated syringes fitted with a needle. For example, a dose of 140 mg docetaxel would require 14 ml docetaxel premix solution.

Inject the required premix volume into a 250 ml infusion bag or bottle containing either 5% glucose solution or sodium chloride 9 mg/ml (0.9%) solution for infusion.

If a dose greater than 200 mg of docetaxel is required, use a larger volume of the infusion vehicle so that a concentration of 0.74 mg/ml docetaxel is not exceeded.

Mix the infusion bag or bottle manually using a rocking motion.

The Docetaxel infusion solution should be used within 4 hours and should be aseptically administered as a 1-hour infusion under room temperature (below 25°C) and normal lighting conditions.

---

As with all parenteral products, Docetaxel premix solution and infusion solution should be visually inspected prior to use, solutions containing a precipitate should be discarded.

---

Any unused medicinal product or waste material should be disposed of in accordance with local requirements

**7. MARKETING AUTHORIZATION HOLDER**

ADMAC LIFESCIENCES

Plot No. 107A & B, EPIP, Jharmajri,  
Baddi, Distt. Solan (H.P) – 174 103 INDIA  
Telephone: 0172-2565161, 2565162  
Fax: 0172-2569286  
E-mail: [admac@admacinternational.com](mailto:admac@admacinternational.com)

**8. MARKETING AUTHORIZATION NUMBER (S)**

H2026/CTD11893/23792

**9. DATE OF FIRST AUTHORIZATION/RENEWAL OF THE AUTHORIZATION**

18/02/2026

**10. DATE OF REVISION OF THE TEXT**

18/02/2026

---