

Summary of Product Characteristics

1. NAME OF THE DRUG PRODUCT :

IOPROST-T EYE DROPS

Latanoprost and Timolol Eye Drops

2. QUALITATIVE AND QUANTITATIVE COMPOSITIONS:

1 ml solution contains latanoprost 50 micrograms and timolol maleate 6.8 mg equivalent to 5 mg timolol.

Excipients with known effect:

Benzalkonium chloride 0.20 mg/mL.

Disodium phosphate (E339ii), sodium dihydrogen phosphate monohydrate (E339i) (containing total phosphate 6.3 mg/mL).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM: Ophthalmic Solution/ Eye Drops

4. CLINICAL PARTICULARS:

4.1 Therapeutical indications :

Reduction of intraocular pressure (IOP), in patients with open angle glaucoma and ocular hypertension who are insufficiently responsive to topical beta-blockers or prostaglandin analogues.

4.2 Posology and method of administration :

Posology

Recommended therapy is one drop in the affected eye(s) once daily.

If one dose is missed, treatment should continue with the next dose as planned. The dose should not exceed one drop in the affected eye(s) daily.

Paediatric population

The safety and efficacy in children and adolescents has not yet been established.

Method of Administration

Precautions to be taken before handling or administering the medicinal product

Contact lenses should be removed before instillation of the eye drops and may be reinserted after 15 minutes.

If more than one topical ophthalmic drug is being used, the drugs should be administered at least five minutes apart.

When using nasolacrimal occlusion or closing the eyelids for 2 minutes, the systemic absorption is reduced. This may result in a decrease in systemic side effects and an increase in local activity.

4.3 Contraindications :

Latanoprost & Timolol Eye Drops is contraindicated in patients with:

Reactive airway disease including bronchial asthma or a history of bronchial asthma, severe chronic obstructive pulmonary disease.

Sinus bradycardia, sick sinus syndrome sino-atrial block, second or third degree atrioventricular block not controlled with a pace-maker. Overt cardiac failure, cardiogenic shock.

Hypersensitivity to the active substances (latanoprost or timolol) or to any of the excipients listed in.

4.4 Special warning and precautions for use :

Systemic effects

Like other topically applied ophthalmic agents, latanoprost and timolol are absorbed systemically. Due to the beta-adrenergic component, timolol, the same types of cardiovascular, pulmonary and other adverse reactions seen with systemic beta-adrenergic blocking agents may occur. The incidence of systemic ADRs after topical ophthalmic administration is lower than for systemic administration. To reduce the systemic absorption.

Cardiac disorders

In patients with cardiovascular diseases (e.g. coronary heart disease, Prinzmetal's angina, and cardiac failure) and hypotension therapy with beta-blockers should be critically assessed and the therapy with other active substances should be considered. Patients with cardiovascular diseases should be watched for signs of deterioration of these diseases and of adverse reactions.

Due to its negative effect on conduction time, beta-blockers should only be given with caution to patients with first degree heart block.

Cardiac reactions, and rarely, death in association with cardiac failures have been reported following administration of timolol

Vascular disorders

Patients with severe peripheral circulatory disturbance/disorders (i.e. severe forms of Raynaud's disease or Raynaud's syndrome) should be treated with caution.

Respiratory disorders

Respiratory reactions including death due to bronchospasm in patients with asthma have been reported following administration of some ophthalmic beta-blockers.

Latanoprost & Timolol Eye Drops should be used with caution, in patients with mild/moderate chronic obstructive pulmonary disease (COPD) and only if the potential benefit outweighs the potential risk.

Hypoglycaemia/diabetes

Beta-blockers should be administered with caution in patients subject to spontaneous hypoglycaemia or to patients with labile diabetes, as beta-blockers may mask the signs and symptoms of acute hypoglycaemia.

Hyperthyroidism

Beta-blockers may also mask the signs of hyperthyroidism.

Corneal diseases

Ophthalmic β -blockers may induce dryness of eyes. Patients with corneal

diseases should be treated with caution.

Other beta-blocking agents

The effect on intra-ocular pressure or the known effects of systemic beta-blockade may be potentiated when timolol is given to the patients already receiving a systemic beta-blocking agent. The response of these patients should be closely observed. The use of two topical beta-adrenergic blocking agents is not recommended.

Anaphylactic reactions

While taking beta-blockers, patients with a history of atopy or a history of severe anaphylactic reaction to a variety of allergens may be more reactive to repeated

challenge with such allergens and unresponsive to the usual doses of adrenaline used to treat anaphylactic reactions.

Surgical anaesthesia

β -blocking ophthalmological preparations may block systemic β -agonist effects e.g. of adrenaline. The anaesthesiologist should be informed when the patient is receiving timolol.

Concomitant therapy

Timolol may interact with other drugs, Interaction with other medicinal products and other forms of interaction.

The effect on intraocular pressure or the known effects of systemic beta-blockade may be potentiated when Latanoprost & Timolol Eye Drops is given to patients already receiving systemic beta-blocking agent. The use of two local beta-blockers or two local prostaglandins is not recommended.

Latanoprost may gradually change the eye colour by increasing the amount of brown pigment in the iris. Similar to experience with latanoprost eye drops, increased iris pigmentation was seen in 16-20% of all patients treated with Latanoprost & Timolol Eye Drops for up to one year (based on photographs). This effect has predominantly been seen in patients with mixed coloured irides, i.e. green-brown, yellow-brown or blue/grey-brown, and is due to

increased melanin content in the stromal melanocytes of the iris. Typically, the brown pigmentation around the pupil spreads concentrically towards the periphery in affected eyes, but the entire iris or parts of it may become more brownish. In patients with homogeneously blue, grey, green or brown eyes, the change has only rarely been seen during two years of treatment in clinical trials with latanoprost.

The change in iris colour occurs slowly and may not be noticeable for several months to years and it has not been associated with any symptom or pathological changes.

No further increase in brown pigment has been observed after discontinuation of treatment, but the resultant colour change may be permanent.

Neither naevi nor freckles of the iris have been affected by treatment.

Accumulation of pigment in the trabecular meshwork or elsewhere in the anterior chamber has not been observed but patients should be examined regularly and, depending on the clinical situation, treatment may be stopped if increased iris pigmentation ensues.

Before treatment is instituted patients should be informed of the possibility of a change in eye colour. Unilateral treatment can result in permanent heterochromia.

There is no documented experience with latanoprost in inflammatory, neovascular, chronic angle closure or congenital glaucoma, in open angle glaucoma of pseudophakic patients and in pigmentary glaucoma. Latanoprost has no or little effect on the pupil but there is no documented experience in acute attacks of closed angle glaucoma. It is recommended, therefore, that Latanoprost & Timolol Eye Drops should be used with caution in these conditions until more experience is obtained.

Latanoprost should be used with caution in patients with a history of herpetic keratitis, and should be avoided in cases of active herpes simplex keratitis and in patients with a history of recurrent herpetic keratitis specifically associated with prostaglandin analogues.

Macular oedema, including cystoid macular oedema, has been reported

during treatment with latanoprost. These reports have mainly occurred in aphakic patients, in pseudophakic patients with a torn posterior lens capsule, or in patients with known risk factors for macular oedema. Latanoprost & Timolol Eye Drops should be used with caution in these patients.

Choroidal detachment has been reported with administration of aqueous suppressant therapy (e.g. timolol, acetazolamide) after filtration procedures.

Use of contact lenses

Latanoprost & Timolol Eye Drops contains benzalkonium chloride, which is commonly used as a preservative in ophthalmic products. Benzalkonium chloride has been reported to cause punctate keratopathy and/or toxic ulcerative keratopathy, may cause eye irritation and is known to discolour soft contact lenses. Close monitoring is required with frequent or prolonged use of Latanoprost & Timolol Eye Drops in dry eye patients, or in conditions where the cornea is compromised. Contact lenses may absorb benzalkonium chloride and these should be removed before applying Latanoprost & Timolol Eye Drops Solution but may be reinserted after 15.

Paediatric population

The safety and efficacy in children and adolescents has not yet been established.

4.5 Interaction with other drugs, other forms of interactions :

No interaction studies have been performed.

There have been reports of paradoxical elevations in intraocular pressure following the concomitant ophthalmic administration of two prostaglandin analogues. Therefore, the use of two or more prostaglandins, prostaglandin analogues, or prostaglandin derivatives is not recommended.

The effect on intraocular pressure or the known effects of systemic beta-blockade may be potentiated when Latanoprost & Timolol Eye Drops is given to patients already receiving an oral beta-adrenergic blocking agent, and the use of two or more topical beta-adrenergic blocking agents is not recommended.

Mydriasis resulting from concomitant use of ophthalmic beta-blockers and adrenaline (epinephrine) has been reported occasionally.

There is a potential for additive effects resulting in hypotension and/or marked bradycardia when ophthalmic beta-blockers solutions are administered concomitantly with oral calcium channel blockers, guanethidine or beta-adrenergic blocking agents, antiarrhythmics (including amiodarone), digitalis glycosides or parasympathomimetics.

Potentiated systemic beta-blockade (e.g. decreased heart rate, depression) has been reported during combined treatment with CYP2D6 inhibitors (e.g. quinidine, fluoxetine, paroxetine) and timolol.

The hypertensive reaction to sudden withdrawal of clonidine can be potentiated when taking beta-blockers.

Beta-blockers may increase the hypoglycaemic effect of antidiabetic agents. Beta-blockers can mask the signs and symptoms of hypoglycaemia.

4.6 Use in pregnancy and lactation :

Latanoprost:

There are no or limited amount of data from the use of latanoprost in pregnant women. Studies in animals have shown reproductive toxicity. The potential risk for humans is unknown.

Timolol:

There are no adequate data for the use of timolol in pregnant women. Timolol should not be used during pregnancy unless clearly necessary. To reduce the systemic absorption.

Epidemiological studies have not revealed malformative effects, but show a risk for intrauterine growth retardation when beta-blockers are administered by the oral route. In addition, signs and symptoms of beta-blockade (e.g. bradycardia, hypotension, respiratory distress and hypoglycaemia) have been observed in the neonate when beta-blockers have been administered until delivery. If Latanoprost & Timolol Eye Drops is administered until delivery,

the neonate should be carefully monitored during the first days of life.

Consequently Latanoprost & Timolol Eye Drops should not be used during pregnancy.

Breastfeeding

Beta- blockers are excreted in breast milk. However, at therapeutic doses of timolol in eye drops it is not likely that sufficient amounts would be present in breast milk to produce clinical symptoms of beta-blockade in the infant. To reduce the systemic absorption.

Consequently Latanoprost and its metabolites are excreted in human milk. Latanoprost & Timolol Eye Drops should not be used during breastfeeding.

Fertility

There are no human data on the effects of Latanoprost & Timolol Eye Drops on fertility.

4.7 Effects on ability to drive and operate machine :

Instillation of eye drops may cause transient blurring of vision. Until this has resolved, patients should not drive or use machines.

4.8 Undesirable effects :

For latanoprost, the majority of adverse events relate to the ocular system. In data from the extension phase of Latanoprost / Timolol pivotal trials, 16 - 20% of patients developed increased iris pigmentation, which may be permanent. In an open 5 year latanoprost safety study, 33% of patients developed iris pigmentation. Other ocular adverse events are generally transient and occur on dose administration.

Like other topically applied ophthalmic drugs, timolol is absorbed into the systemic circulation. This may cause similar undesirable effects as seen with systemic beta-blocking agents. The incidence of systemic ADRs after topical ophthalmic administration is lower than for systemic administration. Listed adverse reactions include reactions seen within the class of ophthalmic beta-blockers.

Adverse events are categorised by frequency as follows: very common ($\geq 1/10$), common ($\geq 1/100$ to $<1/10$), uncommon ($\geq 1/1,000$ to $<1/100$), rare ($\geq 1/10,000$ to $<1/1,000$) and very rare ($<1/10,000$) and not known (cannot be estimated from the available data).

Treatment related adverse events seen in clinical trials with **Latanoprost & Timolol Eye Drops** are listed below.

Nervous System Disorders:

Uncommon: Headache.

Eye Disorders:

Very common: Increased iris pigmentation.

Common: Eye irritation (including stinging, burning and itching), Eye pain.

Uncommon: Eye hyperaemia, Conjunctivitis, Vision blurred, Lacrimation increased, Blepharitis, Corneal disorders.

Skin and Subcutaneous Tissue Disorders:

Uncommon: Skin rash, Pruritus.

Additional adverse events have been reported specific to the use of the individual components of Latanoprost & Timolol Eye Drops in either in clinical studies, spontaneous reports or in the available literature.

For **Latanoprost**, these are:

Infection and Infestations:

Hepatitis Keratitis

Nervous System Disorders

Dizziness

Eye Disorders:

Eyelash and vellus hair changes (increased length, thickness, pigmentation, and number), Punctate epithelial erosions, periorbital oedema, iritis/uveitis, macular oedema (in aphakic, pseudophakic patients with torn posterior lens capsules or in patients with known risk factors for macular oedema), dry eye, keratitis, corneal oedema and erosions, misdirected eyelashes sometimes

resulting in eye irritation and iris cyst.

Cardiac Disorders:

Aggravation of angina in patients with pre-existing disease, palpitations.

Respiratory, Thoracic and Mediastinal Disorders:

Asthma, asthma aggravation,
dyspnea. *Skin and Subcutaneous*

Tissue Disorders: Darkening of
palpebral skin.

Musculoskeletal, Connective Tissue and Bone Disorders:

Joint pain, Muscle pain.

General disorders and Administration Site Conditions:

Chest pain

For **timolol**, these are:

Immune System Disorders:

Signs and symptoms of systemic allergic reactions including angioedema, urticaria, and localized and generalized rash, pruritus, anaphylactic reaction.

Metabolism and nutrition disorders:

Hypoglycaemia.

Psychiatric Disorders:

Depression, memory loss, decreased libido, insomnia, nightmares.

Nervous System Disorders:

Dizziness, paraesthesia, cerebral ischemia, cerebrovascular accident, increase in signs and symptoms of myasthenia gravis, syncope, and headache

Eye Disorders:

Signs and symptoms of ocular irritation (e.g. burning, stinging, itching, tearing, redness), Blepharitis, keratitis, blurred vision, decreased corneal sensitivity, and dry-eyes, corneal erosion, Visual disturbances including refractive changes (due to withdrawal of miotic therapy in some cases), diplopia, ptosis, choroidal detachment (following filtration surgery).

Ear and Labyrinth Disorders:

Tinnitus

Cardiac Disorders:

Palpitations, arrhythmia, bradycardia, cardiac arrest, heart block, congestive heart failure, chest pain, oedema, atrioventricular block, cardiac failure.

Vascular Disorders:

Hypotension, Raynaud's phenomenon, cold hands and feet.

Respiratory, Thoracic and Mediastinal Disorders:

Bronchospasm (predominantly in patients with pre-existing bronchospastic disease), dyspnoea, cough.

Gastrointestinal Disorders:

Dysgeusia, nausea, diarrhoea, dyspepsia, dry mouth, abdominal pain, vomiting.

Skin and Subcutaneous Tissue Disorders:

Alopecia, psoriasiform rash or exacerbation of psoriasis, skin rash.

Musculoskeletal and connective tissue disorders:

Myalgia.

Reproductive system and breast disorders:

Sexual dysfunction, decreased libido.

General Disorders and Administration Site Conditions:

Asthenia/fatigue

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. Healthcare professionals are requested to report any suspected adverse reactions via the Pharmacy and Poisons Reporting System (PvERS) <https://pv.pharmacyboardkenya.org>

.

4.9 Overdoses:

Symptoms

No data are available in humans with regard to overdose with Latanoprost &

Timolol Eye Drops.

Symptoms of systemic timolol overdose are: bradycardia, hypotension, bronchospasm and cardiac arrest. If such symptoms occur the treatment should be symptomatic and supportive. Studies have shown that timolol does not dialyse readily.

Apart from ocular irritation and conjunctival hyperaemia no other ocular or systemic side effects are known if latanoprost is overdosed.

Management

If latanoprost is accidentally ingested orally the following information may be useful:

Treatment: Gastric lavage if needed Symptomatic treatment. Latanoprost is extensively metabolised during the first pass through the liver. Intravenous infusion of 3 micrograms / kg in healthy volunteers induced no symptoms but a dose of 5.5-10 micrograms /kg caused nausea, abdominal pain, dizziness, fatigue, hot flushes and sweating. These events were mild to moderate in severity and resolved without treatment, within 4 hours after terminating the infusion.

5 PHARMACEUTICAL PROPERTIES:

5.1 PHARMACO-KINETIC PROPERTIES:

Pharmacokinetic properties

Latanoprost

Latanoprost is an isopropyl ester prodrug, which per se is inactive, but after hydrolysis by esterases in the cornea to the acid of latanoprost, becomes biologically active. The prodrug is well absorbed through the cornea and all drug that enters the aqueous humor is hydrolysed during the passage through the cornea. Studies in man indicate that the maximum concentration in the aqueous humour, approximately 15-30 ng/ml, is reached about 2 hours after topical administration of latanoprost alone. After topical application in monkeys latanoprost is distributed primarily in the anterior segment, the conjunctiva and the eye lids.

The acid of latanoprost has a plasma clearance of 0.40 l/h/kg and a small volume of distribution, 0.16 l/kg, resulting in a rapid half life in plasma, 17 minutes. After topical ocular administration the systemic bioavailability of the acid of latanoprost is 45%. The acid of latanoprost has a plasma protein binding of 87%.

There is practically no metabolism of the acid of latanoprost in the eye. The main metabolism occurs in the liver. The main metabolites, the 1, 2-dinor and 1, 2, 3, 4- tetranor metabolites, exert no or only weak biological activity in animal studies and are excreted primarily in the urine.

Timolol

The maximum concentration of timolol in the aqueous humor is reached about 1 hour after topical administration of eye drops. Part of the dose is absorbed systemically and a maximum plasma concentration of 1 ng/ml is reached 10-20 minutes after topical administration of one eye drop to each eye once daily (300 micrograms/day). The half-life of timolol in plasma is about 6 hours. Timolol is extensively metabolised in the liver. The metabolites are excreted in the urine together with some unchanged Timolol.

Latanoprost & Timolol Eye Drops:

No pharmacokinetic interactions between latanoprost and timolol were observed although there was an approximate 2-fold increased concentration of the acid of latanoprost in aqueous humour 1-4 hours after administration of *Latanoprost & Timolol Eye Drops* compared to monotherapy.

5.2 PHARMACO-DYNAMICS PROPERTIES :

Pharmacodynamic properties General properties

Pharmacotherapeutic group: Ophthalmological-beta blocking agents -

Timolol, combinations. ATC code: **S01ED51**

Mechanism of action

Latanoprost & Timolol Eye Drops consists of two components: Latanoprost

and Timolol Maleate. These two components decrease elevated intraocular pressure (IOP) by different mechanisms of action and the combined effect results in additional IOP reduction compared to either compound administered alone.

Latanoprost, a prostaglandin $F_{2\alpha}$ analogue, is a selective prostanoid FP receptor agonist that reduces the IOP by increasing the outflow of aqueous humour. The main mechanism of action is increased uveoscleral outflow. Additionally, some increase in outflow facility (decrease in trabecular outflow resistance) has been reported in man. Latanoprost has no significant effect on the production of aqueous humour, the blood-aqueous barrier or the intraocular blood circulation. Chronic treatment with latanoprost in monkey eyes, which had undergone extracapsular lens extraction, did not affect the retinal blood vessels as determined by fluorescein angiography. Latanoprost has not induced fluorescein leakage in the posterior segment of pseudophakic human eyes during short term treatment.

Timolol is a beta-1 and beta-2 (non-selective) adrenergic receptor blocking agent that has no significant intrinsic sympathomimetic, direct myocardial depressant or membrane-stabilising activity. Timolol lowers IOP by decreasing the formation of aqueous in the ciliary epithelium.

The precise mechanism of action is not clearly established, but inhibition of the increased cyclic AMP synthesis caused by endogenous beta-adrenergic stimulation is probable. Timolol has not been found to significantly affect the permeability of the blood-aqueous barrier to plasma proteins. In rabbits, timolol was without effect on the regional ocular blood flow after chronic treatment.

Pharmacodynamic effects

Clinical effects

In dose finding studies, Latanoprost/ Timolol produced significantly greater decreases in mean diurnal IOP compared to latanoprost and timolol administered once daily as monotherapy. In two well controlled, double

masked six-month clinical studies the IOP reducing effect of Latanoprost/ Timolol was compared with latanoprost and timolol monotherapy in patients with an IOP of at least 25 mm Hg or greater. Following a 2-4 week run-in with timolol (mean decrease in IOP from enrollment of 5 mm Hg), additional decreases in mean diurnal IOP of 3.1, 2.0 and 0.6 mm Hg were observed after 6 months of treatment for Latanoprost/ Timolol (twice daily), respectively. The IOP lowering effect of Latanoprost/ Timolol was maintained in 6-month open label extension of these studies.

Existing data suggest that evening dosing may be more effective in IOP lowering than morning dosing. However, when considering a recommendation of either morning or evening dosing, sufficient consideration should be given to the lifestyle of the patient and their likely compliance.

It should be kept in mind that in case of insufficient efficacy of the fixed combination, results from studies indicate that the use of unfixed administration of Timolol bid and latanoprost once a day might be still efficient.

Onset of action of Latanoprost & Timolol Eye Drops is within one hour and maximal effect occurs within six to eight hours. Adequate IOP reducing effect has been shown to be present up to 24 hours post dosage after multiple treatments.

5.3 Pre-clinical safety data

Preclinical safety data

The ocular and systemic safety profile of the individual components is well established. No adverse ocular or systemic effects were seen in rabbits treated topically with the fixed combination or with concomitantly administered latanoprost and timolol ophthalmic

solutions. Safety pharmacology, genotoxicity and carcinogenicity studies with each of the components revealed no special hazards for humans. Latanoprost did not affect corneal wound healing in the rabbit eye, whereas timolol

inhibited the process in the rabbit and the monkey eye when administered more frequently than once a day.

For latanoprost, no effects on male and female fertility in rats and no teratogenic potential in rats and rabbits have been established. No embryotoxicity was observed in rats after intravenous doses of up to 250 micrograms/kg/day. Latanoprost, however, caused embryofetal toxicity, characterised by increased incidence of late resorption and abortion and by reduced fetal weight, in rabbits at intravenous doses of 5 micrograms/kg/day (approximately 100 times the clinical dose) and above. Timolol showed no effects on male and female fertility in rats or teratogenic potential in mice, rats and rabbits.

6 PHARMACEUTICAL PARTICULARS :

6.1 List of excipients :

Sr. No.	Ingredient	Grade
1	Benzalkonium Chloride (50% solution)	Ph. Eur.
2	Sodium Dihydrogen Phosphate Monohydrate	BP
3	Disodium Phosphate, Anhydrous	Ph. Eur.
4	Sodium Chloride	Ph. Eur.
5	Sodium Hydroxide	Ph. Eur.
6	Hydrochloric Acid	Ph. Eur.
7	Water For Injection	Ph. Eur.

6.2 Incompatibilities:

In vitro studies have shown that precipitation occurs when eye drops containing thiomersal are mixed with Latanoprost and Timolol Eye Drops. If such drugs are used concomitantly with Latanoprost & Timolol Eye Drops, they should be administered with an interval of at least five minutes.

6.3 Shelf life:

Before first opening: 2 years

After first opening of bottle: 4 weeks

6.4 Special precaution for storage:

Store between 2°C and

8°C. After opening, store

below 25°C.

Keep the bottle in the outer carton in order to protect from light. Keep out of the sight and reach of children

7. MARKETING AUTHORIZATION HOLDER:

Marketing Authorization Holder	Name: Surgipharm Limited Full Physical address: Nivina Towers - 5th Floor ,71 Westlands Road P.O Box 46043, 00100 , Nairobi, Kenya Telephone Number: +254 20 3749304 / 3749306 Fax Number: +254 20 3741262
Manufacturer of the product	Name: FDC Limited Full Physical address: B-8, MIDC Industrial Area, Waluj, Aurangabad- 431 136, Maharashtra State India.

8. MARKETING AUTHORIZATION NUMBER (LIC. NO):

Kenya : H2022/CTD9747/17718

9. DATE OF FIRST AUTHORIZATION/RENEWAL OF AUTHORIZATION :

---- Mon Sep 18 10:14:39 EAT 2023

10. DATE OF REVISION OF THE TEXT: Mon Sep 18 10:14:39 EAT 2023 ---

-