

SUMMARY OF PRODUCT CHARACTERISTICS (SPCs)

1. Name of the medicinal product:

Lerace 250mg Tablets

2. Qualitative and quantitative composition

Each film coated tablet contains:

Levetiracetam250mg

For the full list of excipients, see Section 6.1

3. Pharmaceutical form

White, Square biconvex film coated tablet “LER 250” Engraved on one side and Break line on other side.

4. Clinical particulars

4.1 Therapeutic indications

Lerace is indicated as monotherapy in the treatment of partial onset seizures with or without secondary generalisation in adults and adolescents from 16 years of age with newly diagnosed epilepsy.

Lerace is indicated as adjunctive therapy

- in the treatment of partial onset seizures with or without secondary generalisation in adults, adolescents, children and infants from 1 month of age with epilepsy.
- in the treatment of myoclonic seizures in adults and adolescents from 12 years of age with Juvenile Myoclonic Epilepsy.
- in the treatment of primary generalised tonic-clonic seizures in adults and adolescents from 12 years of age with Idiopathic Generalised Epilepsy.

4.2 Posology and method of administration

Monotherapy for adults and adolescents from 16 years of age

The recommended starting dose is 250 mg twice daily which should be increased to an initial therapeutic dose of 500 mg twice daily after two weeks. The dose can be further increased by 250 mg twice daily every two weeks depending upon the clinical response. The maximum dose is 1500 mg twice daily.

Add-on therapy for adults (≥18 years) and adolescents (12 to 17 years) weighing 50 kg or more

The initial therapeutic dose is 500 mg twice daily. This dose can be started on the first day of treatment.

Depending upon the clinical response and tolerability, the daily dose can be increased up to 1,500 mg twice daily. Dose changes can be made in 500 mg twice daily increases or decreases every two to four weeks.

Discontinuation

If levetiracetam has to be discontinued it is recommended to withdraw it gradually (e.g. in adults and adolescents weighing more than 50 kg: 500 mg decreases twice daily every two to four weeks; in infants older than 6 months, children and adolescents weighing less than 50 kg: dose decrease should not exceed 10 mg/kg twice daily every two weeks; in infants (less than 6 months): dose decrease should not exceed 7 mg/kg twice daily every two weeks).

Special populations

Elderly (65 years and older)

Adjustment of the dose is recommended in elderly patients with compromised renal function (see “Renal impairment” below).

Renal impairment

The daily dose must be individualised according to renal function.

For adult patients, refer to the following table and adjust the dose as indicated. To use this dosing table, an estimate of the patient's creatinine clearance (CLcr) in ml/min is needed. The CLcr in ml/min may be estimated from serum creatinine (mg/dl) determination, for adults and adolescents weighing 50 kg or more, the following formula:

$$\text{CLcr (ml/min)} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dl)}} \quad (\times 0.85 \text{ for women})$$

Then CLcr is adjusted for body surface area (BSA) as follows

$$\text{CLcr (ml/min/1.73 m}^2\text{)} = \frac{\text{CLcr (ml/min)}}{\text{BSA subject (m}^2\text{)}} \times 1.73$$

Dosing adjustment for adult and adolescents patients weighing more than 50 kg with impaired renal function:

Group	Creatinine clearance (ml/min/1.73m ²)	Dose and frequency
Normal	> 80	500 to 1,500 mg twice daily
Mild	50-79	500 to 1,000 mg twice daily
Moderate	30-49	250 to 750 mg twice daily
Severe	< 30	250 to 500 mg twice daily
End-stage renal disease patients undergoing dialysis ⁽¹⁾	-	500 to 1,000 mg once daily ⁽²⁾

(1) A 750 mg loading dose is recommended on the first day of treatment with levetiracetam.

(2) Following dialysis, a 250 to 500 mg supplemental dose is recommended.

For children with renal impairment, levetiracetam dose needs to be adjusted based on the renal function as levetiracetam clearance is related to renal function. This recommendation is based on a study in adult renally impaired patients.

The CL_{Cr} in ml/min/1.73 m² may be estimated from serum creatinine (mg/dl) determination, for young adolescents, children and infants, using the following formula (Schwartz formula):

$$CL_{Cr} (ml/min/1.73 m^2) = \frac{\text{Height (cm)} \times k_s}{\text{Serum Creatinine (mg/dl)}}$$

k_s= 0.45 in Term infants to 1 year old; k_s= 0.55 in Children to less than 13 years and in adolescent female; k_s= 0.7 in adolescent male

Dosing adjustment for infants, children and adolescents patients weighing less than 50 kg with impaired renal function:

	Creatinine	Dose and frequency (1)
--	------------	------------------------

Group	clearance (ml/min/1.7 3m ²)	Infants 1 to less than 6 months	Infants 6 to 23 months, children and adolescents weighing less than 50 kg
Normal	> 80	7 to 21 mg/kg (0.07 to 0.21 ml/kg) twice daily	10 to 30 mg/kg (0.10 to 0.30 ml/kg) twice daily
Mild	50-79	7 to 14 mg/kg (0.07 to 0.14 ml/kg) twice daily	10 to 20 mg/kg (0.10 to 0.20 ml/kg) twice daily
Moderate	30-49	3.5 to 10.5 mg/kg (0.035 to 0.105 ml/kg) twice daily	5 to 15 mg/kg (0.05 to 0.15 ml/kg) twice daily
Severe	< 30	3.5 to 7 mg/kg (0.035 to 0.07 ml/kg) twice daily	5 to 10 mg/kg (0.05 to 0.10 ml/kg) twice daily
End-stage renal disease patients undergoing dialysis	--	7 to 14 mg/kg (0.07 to 0.14 ml/kg) once daily ⁽²⁾ ⁽⁴⁾	10 to 20 mg/kg (0.10 to 0.20 ml/kg) once daily ⁽³⁾ ⁽⁵⁾

(1) Lerace oral solution should be used for doses under 250 mg, for doses not multiple of 250 mg when dosing recommendation is not achievable by taking multiple tablets and for patients unable to swallow tablets.

(2) A 10.5 mg/kg (0.105 ml/kg) loading dose is recommended on the first day of treatment with levetiracetam.

(3) A 15 mg/kg (0.15 ml/kg) loading dose is recommended on the first day of treatment with levetiracetam.

(4) Following dialysis, a 3.5 to 7 mg/kg (0.035 to 0.07 ml/kg) supplemental dose is recommended.

(5) Following dialysis, a 5 to 10 mg/kg (0.05 to 0.10 ml/kg) supplemental dose is recommended.

Hepatic impairment

No dose adjustment is needed in patients with mild to moderate hepatic impairment. In patients with severe hepatic impairment, the creatinine clearance may underestimate the renal insufficiency. Therefore a 50 % reduction of the daily maintenance dose is recommended when the creatinine clearance is < 60 ml/min/1.73 m².

Paediatric population

The physician should prescribe the most appropriate pharmaceutical form, presentation and strength according to age, weight and dose.

The tablet formulation is not adapted for use in infants and children under the age of 6 years. Lerace oral solution is the preferred formulation for use in this population. In addition, the available dose strengths of the tablets are not appropriate for initial treatment in children weighing less than 25 kg, for patients unable to swallow tablets or for the administration of doses below 250 mg. In all of the above cases Lerace oral solution should be used.

Monotherapy

The safety and efficacy of Lerace in children and adolescents below 16 years as monotherapy treatment have not been established.

No data are available.

Add-on therapy for infants aged 6 to 23 months, children (2 to 11 years) and adolescents (12 to 17 years) weighing less than 50 kg

Lerace oral solution is the preferred formulation for use in infants and children under the age of 6 years.

For children 6 years and above, Lerace oral solution should be used for doses under 250 mg, for doses not multiple of 250 mg when dosing recommendation is not achievable by taking multiple tablets and for patients unable to swallow tablets.

The lowest effective dose should be used. The starting dose for a child or adolescent of 25kg should be 250mg twice daily with a maximum dose of 750mg twice daily. Dose in children 50 kg or greater is the same as in adults.

Add-on therapy for infants aged from 1 month to less than 6 months

The oral solution is the formulation to use in infants.

Method of administration

The film-coated tablets must be taken orally, swallowed with a sufficient quantity of liquid and may be taken with or without food. The daily dose is administered in two equally divided doses.

4.3 Contraindications

Hypersensitivity to the active substance or other pyrrolidone derivatives or to any of the excipients listed in section 6.1.

4.4 Precautions for use

Renal impairment

The administration of levetiracetam to patients with renal impairment may require dose adjustment. In patients with severely impaired hepatic function, assessment of renal function is recommended before dose selection.

Acute Kidney injury

The use of levetiracetam has been very rarely associated with acute kidney injury, with a time to onset ranging from a few days to several months.

Blood cell counts

Rare cases of decreased blood cell counts (neutropenia, agranulocytosis, leucopenia, thrombocytopenia and pancytopenia) have been described in association with levetiracetam administration, generally at the beginning of the treatment. Complete blood cell counts are advised in patients experiencing important weakness, pyrexia, recurrent infections or coagulation disorders.

Suicide

Suicide, suicide attempt, suicidal ideation and behaviour have been reported in patients treated with anti-epileptic agents (including levetiracetam). A meta-analysis of randomized placebo-controlled trials of anti-epileptic medicinal products has shown a small increased risk of suicidal thoughts and behaviour. The mechanism of this risk is not known.

Therefore, patients should be monitored for signs of depression and/or suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of depression and/or suicidal ideation or behaviour emerge.

Worsening of seizures

As with other types of antiepileptic drugs, levetiracetam may rarely exacerbate seizure frequency or severity. This paradoxical effect was mostly reported within the first month after levetiracetam initiation or increase of the dose, and was reversible upon drug discontinuation or dose decrease. Patients should be advised to consult their physician immediately in case of aggravation of epilepsy.

Lack of efficacy or seizure worsening has been reported in patients with epilepsy associated with sodium voltage-gated channel alpha subunit 8 (SCN8A) mutations.

Electrocardiogram QT interval prolongation

Rare cases of ECG QT interval prolongation have been observed during the post-marketing surveillance. Levetiracetam should be used with caution in patients with QTc-interval prolongation, in patients concomitantly treated with drugs affecting the QTc-interval, or in patients with relevant pre-existing cardiac disease or electrolyte disturbances.

Paediatric population

The tablet formulation is not adapted for use in infants and children under the age of 6 years.

Available data in children did not suggest impact on growth and puberty. However, long term effects on learning, intelligence, growth, endocrine function, puberty and childbearing potential in children remain unknown.

4.5 Interaction with other medicinal products and other form of interaction

Antiepileptic medicinal products

Pre-marketing data from clinical studies conducted in adults indicate that levetiracetam did not influence the serum concentrations of existing antiepileptic medicinal products (phenytoin, carbamazepine, valproic acid, phenobarbital, lamotrigine, gabapentin and primidone) and that these antiepileptic medicinal products did not influence the pharmacokinetics of levetiracetam.

As in adults, there is no evidence of clinically significant medicinal product interactions in paediatric patients receiving up to 60 mg/kg/day levetiracetam.

A retrospective assessment of pharmacokinetic interactions in children and adolescents with epilepsy (4 to 17 years) confirmed that adjunctive therapy with orally administered levetiracetam did not influence the steady-state serum concentrations of concomitantly administered carbamazepine and valproate. However, data suggested a 20 % higher

levetiracetam clearance in children taking enzyme-inducing antiepileptic medicinal products. Dose adjustment is not required.

Probenecid

Probenecid (500 mg four times daily), a renal tubular secretion blocking agent, has been shown to inhibit the renal clearance of the primary metabolite, but not of levetiracetam. Nevertheless, the concentration of this metabolite remains low.

Methotrexate

Concomitant administration of levetiracetam and methotrexate has been reported to decrease methotrexate clearance, resulting in increased/prolonged blood methotrexate concentration to potentially toxic levels. Blood methotrexate and levetiracetam levels should be carefully monitored in patients treated concomitantly with the two drugs.

Oral contraceptives and other pharmacokinetics interactions

Levetiracetam 1,000 mg daily did not influence the pharmacokinetics of oral contraceptives (ethinyl-estradiol and levonorgestrel); endocrine parameters (luteinizing hormone and progesterone) were not modified. Levetiracetam 2,000 mg daily did not influence the pharmacokinetics of digoxin and warfarin; prothrombin times were not modified. Co-administration with digoxin, oral contraceptives and warfarin did not influence the pharmacokinetics of levetiracetam.

Laxatives

There have been isolated reports of decreased levetiracetam efficacy when the osmotic laxative macrogol has been concomitantly administered with oral levetiracetam. Therefore, macrogol should not be taken orally for one hour before and for one hour after taking levetiracetam.

Food and alcohol

The extent of absorption of levetiracetam was not altered by food, but the rate of absorption was slightly reduced.

No data on the interaction of levetiracetam with alcohol are available.

4.6 Fertility, pregnancy and lactation

Women of child bearing potential

Specialist advice should be given to women who are of childbearing potential. Treatment with levetiracetam should be reviewed when a woman is planning to become pregnant. As with all antiepileptic medicines, sudden discontinuation of levetiracetam should be avoided as this may lead to breakthrough seizures that could have serious consequences for the woman and the unborn child. Monotherapy should be preferred whenever possible because therapy with multiple antiepileptic medicines AEDs could be associated with a higher risk of congenital malformations than monotherapy, depending on the associated antiepileptics.

Pregnancy

A large amount of post-marketing data on pregnant women exposed to levetiracetam monotherapy (more than 1800, among which in more than 1500 exposure occurred during the 1st trimester) do not suggest an increase in the risk for major congenital malformations. Limited evidence is available on the neurodevelopment of children exposed to levetiracetam monotherapy in utero. Data from two observational population-based registry studies undertaken in largely the same dataset from the Nordic countries and including more than 1000 children born to women with epilepsy prenatally exposed to levetiracetam monotherapy do not suggest an increased risk of autism spectrum disorders or intellectual disability compared to children born to women with epilepsy not exposed to an antiepileptic drug in utero. The mean follow-up time of children in the levetiracetam group was shorter than for the group of children non exposed to any antiepileptic drug (e.g. 4.4 years vs 6.8 years in one of the studies).

Levetiracetam can be used during pregnancy, if after careful assessment it is considered clinically needed. In such case, the lowest effective dose is recommended.

Physiological changes during pregnancy may affect levetiracetam concentration. Decrease in levetiracetam plasma concentrations has been observed during pregnancy. This decrease is more pronounced during the third trimester (up to 60% of baseline concentration before

pregnancy). Appropriate clinical management of pregnant women treated with levetiracetam should be ensured.

Breastfeeding

Levetiracetam is excreted in human breast milk. Therefore, breast-feeding is not recommended.

However, if levetiracetam treatment is needed during breastfeeding, the benefit/risk of the treatment should be weighed considering the importance of breastfeeding.

Fertility

No impact on fertility was detected in animal studies. No clinical data are available, potential risk for human is unknown.

4.7 Effects on ability to drive and use machines

Levetiracetam has minor or moderate influence on the ability to drive and use machines.

Due to possible different individual sensitivity, some patients might experience somnolence or other central nervous system related symptoms, especially at the beginning of treatment or following a dose increase. Therefore, caution is recommended in those patients when performing skilled tasks, e.g. driving vehicles or operating machinery. Patients are advised not to drive or use machines until it is established that their ability to perform such activities is not affected.

4.7 Undesirable effects

Summary of the safety profile

The most frequently reported adverse reactions were nasopharyngitis, somnolence, headache, fatigue and dizziness. The adverse reaction profile presented below is based on the analysis of pooled placebo-controlled clinical trials with all indications studied, with a total of 3,416 patients treated with Levetiracetam. These data are supplemented with the use of Levetiracetam in corresponding open-label extension studies, as well as post-marketing experience. The safety profile of Levetiracetam is generally similar across age groups (adult and pediatric patients) and across the approved

epilepsy indications. Since there was limited exposure for Levetiracetam intravenous use and since oral and intravenous formulations are bioequivalent, the safety information of Levetiracetam intravenous will rely on Levetiracetam oral use.

Tabulated list of adverse reactions

Adverse reactions reported in clinical studies (adults, adolescents, children and infants >1 month) and from post-marketing experience are listed in the following table per System Organ Class and per frequency. Adverse reactions are presented in the order of decreasing seriousness and their frequency is defined as follows: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$) and very rare ($< 1/10,000$).

MedDRA SOC	Frequency category				
	Very common	Common	Uncommon	Rare	Very rare
Infections and infestations	Nasopharyngitis			Infection	
Blood and lymphatic system disorders			Thrombocytopenia, leukopenia	Pancytopenia, neutropenia, agranulocytosis	
Immune system disorders				Drug reaction with eosinophilia and systemic symptoms (DRESS), Hypersensitivity (including angioedema and anaphylaxis)	
Metabolism and nutrition disorders		Anorexia	Weight decreased, weight increase	Hyponatremia	
Psychiatric disorders		Depression, hostility/aggression, anxiety, insomnia, nervousness/irritability	Suicide attempt, suicidal ideation, psychotic disorder, abnormal behavior, hallucination, anger, confusional state, panic attack, affect	Completed suicide, personality disorder, thinking abnormal, delirium	Obsessive compulsive disorder**

MedDRA SOC	Frequency category				
	Very common	Common	Uncommon	Rare	Very rare
			lability/mood swings, agitation		
Nervous system disorders	Somnolence, headache	Convulsion, balance disorder, dizziness, lethargy, tremor	Amnesia, memory impairment, coordination abnormal/ataxia, paresthesia, disturbance in attention	Choreoathetosis, dyskinesia, hyperkinesia, gait disturbance, encephalopathy, seizures aggravated, Neuroleptic malignant syndrome*	
Eye disorders			Diplopia, vision blurred		
Ear and labyrinth disorders		Vertigo			
Cardiac disorders				Electrocardiogram QT prolonged	
Respiratory, thoracic and mediastinal disorders		Cough			
Gastrointestinal disorders		Abdominal pain, diarrhea, dyspepsia, vomiting, nausea		Pancreatitis	
Hepatobiliary disorders			Liver function test abnormal	Hepatic failure, hepatitis	

Renal and urinary disorders				Acute kidney injury	
Skin and subcutaneous tissue disorders		Rash	Alopecia, eczema, pruritus	Toxic epidermal necrolysis, Stevens-Johnson syndrome, erythema multiforme	
Musculoskeletal and connective tissue disorders			Muscular weakness, myalgia	Rhabdomyolysis and blood creatine phosphokinase increased*	
General disorders and administration site conditions		Asthenia/fatigue			
MedDRA SOC	Frequency category				
	Very common	Common	Uncommon	Rare	Very rare
Injury, poisoning and procedural complications			Injury		

* Prevalence is significantly higher in Japanese patients when compared to non-Japanese patients.

**Very rare cases of development of obsessive-compulsive disorders (OCD) in patients with underlying history of OCD or psychiatric disorders have been observed in post-marketing surveillance.

Description of selected adverse reactions

The risk of anorexia is higher when Levetiracetam is co-administered with topiramate.

In several cases of alopecia, recovery was observed when Levetiracetam was discontinued.

Bone marrow suppression was identified in some of the cases of pancytopenia.

Cases of encephalopathy generally occurred at the beginning of the treatment (few days to a few months) and were reversible after treatment discontinuation.

Pediatric population

In patients aged 1 month to less than 4 years, a total of 190 patients have been treated with Levetiracetam in placebo-controlled and open label extension studies. Sixty of these patients were treated with Levetiracetam in placebo-controlled studies. In patients aged 4-16 years, a total of 645 patients have been treated with Levetiracetam in placebo-controlled and open label extension studies. 233 of these patients were treated with Levetiracetam in placebo-controlled studies. In both these pediatric age ranges, these data are supplemented with the post-marketing experience of the use of Levetiracetam.

In addition, 101 infants aged less than 12 months have been exposed in a post authorization safety study. No new safety concerns for levetiracetam were identified for infants less than 12 months of age with epilepsy.

The adverse reaction profile of Levetiracetam is generally similar across age groups and across the approved epilepsy indications. Safety results in pediatric patients in placebo-controlled clinical studies were consistent with the safety profile of Levetiracetam in adults except for behavioral and psychiatric adverse reactions which were more common in children than in adults. In children and adolescents aged 4 to 16 years, vomiting (very common, 11.2%), agitation (common, 3.4%), mood swings (common, 2.1%), affect lability (common, 1.7%), aggression (common, 8.2%), abnormal behavior (common, 5.6%), and lethargy (common, 3.9%) were reported more frequently than in other age ranges or in the overall safety profile. In infants and children aged 1 month to less than 4 years, irritability (very common, 11.7%) and coordination abnormal (common, 3.3%) were reported more frequently than in other age groups or in the overall safety profile.

A double-blind, placebo-controlled pediatric safety study with a non-inferiority design has assessed the cognitive and neuropsychological effects of Levetiracetam in children 4 to 16 years of age with partial onset seizures. It was concluded that Levetiracetam was not different

(non-inferior) from placebo with regard to the change from baseline of the Leiter-R Attention and Memory, Memory Screen Composite score in the per- protocol population. Results related to behavioral and emotional functioning indicated a worsening in Levetiracetam treated patients on aggressive behavior as measured in a standardized and systematic way using a validated instrument (CBCL - Achenbach Child Behavior Checklist). However, subjects, who took Levetiracetam in the longterm open label follow-up study, did not experience a worsening, on average, in their behavioral and emotional functioning; in particular measures of aggressive behavior were not worse than baseline.

4.8 Overdose

Symptoms

Somnolence, agitation, aggression, depressed level of consciousness, respiratory depression and coma were observed with Levetiracetam overdoses.

Management of overdose

After an acute overdose, the stomach may be emptied by gastric lavage or by induction of emesis. There is no specific antidote for levetiracetam. Treatment of an overdose will be symptomatic and may include haemodialysis. The dialyser extraction efficiency is 60 % for levetiracetam and 74 % for the primary metabolite.

Reporting of suspected adverse reactions:

Healthcare professionals are requested to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org>

5. Pharmacology

Pharmacotherapeutic group: antiepileptics, other antiepileptics, ATC code: N03AX14.

The active substance, levetiracetam, is a pyrrolidone derivative (S-enantiomer of α -ethyl-2-oxo-1-pyrrolidine acetamide), chemically unrelated to existing antiepileptic active substances.

Mechanism of action

The mechanism of action of levetiracetam still remains to be fully elucidated. In vitro and in vivo experiments suggest that levetiracetam does not alter basic cell characteristics and normal neurotransmission.

In vitro studies show that levetiracetam affects intraneuronal Ca^{2+} levels by partial inhibition of N-type Ca^{2+} currents and by reducing the release of Ca^{2+} from intraneuronal stores. In addition it partially reverses the reductions in GABA- and glycine-gated currents induced by zinc and β -carbolines. Furthermore, levetiracetam has been shown in in vitro studies to bind to a specific site in rodent brain tissue. This binding site is the synaptic vesicle protein 2A, believed to be involved in vesicle fusion and neurotransmitter exocytosis. Levetiracetam and related analogs show a rank order of affinity for binding to the synaptic vesicle protein 2A which correlates with the potency of their anti-seizure protection in the mouse audiogenic model of epilepsy. This finding suggests that the interaction between levetiracetam and the synaptic vesicle protein 2A seems to contribute to the antiepileptic mechanism of action of the medicinal product.

5.1 Pharmacodynamic effects

Levetiracetam induces seizure protection in a broad range of animal models of partial and primary generalised seizures without having a pro-convulsant effect. The primary metabolite is inactive.

In man, an activity in both partial and generalised epilepsy conditions (epileptiform discharge/photoparoxysmal response) has confirmed the broad spectrum pharmacological profile of levetiracetam.

5.2 Pharmacokinetic properties

Levetiracetam is a highly soluble and permeable compound. The pharmacokinetic profile is linear with low intra- and inter-subject variability. There is no modification of the clearance after repeated administration. There is no evidence for any relevant gender, race or circadian variability. The pharmacokinetic profile is comparable in healthy volunteers and in patients with epilepsy.

Due to its complete and linear absorption, plasma levels can be predicted from the oral dose of levetiracetam expressed as mg/kg bodyweight. Therefore, there is no need for plasma level monitoring of levetiracetam.

A significant correlation between saliva and plasma concentrations has been shown in adults and children (ratio of saliva/plasma concentrations ranged from 1 to 1.7 for oral tablet formulation and after 4 hours post-dose for oral solution formulation).

Adults and adolescents

Absorption

Levetiracetam is rapidly absorbed after oral administration. Oral absolute bioavailability is close to 100 %.

Peak plasma concentrations (C_{max}) are achieved at 1.3 hours after dosing. Steady-state is achieved after two days of a twice daily administration schedule.

Peak concentrations (C_{max}) are typically 31 and 43 µg/ml following a single 1,000 mg dose and repeated 1,000 mg twice daily dose, respectively.

The extent of absorption is dose-independent and is not altered by food.

Distribution

No tissue distribution data are available in humans.

Neither levetiracetam nor its primary metabolite are significantly bound to plasma proteins (< 10 %).

The volume of distribution of levetiracetam is approximately 0.5 to 0.7 l/kg, a value close to the total body water volume.

Biotransformation

Levetiracetam is not extensively metabolised in humans. The major metabolic pathway (24 % of the dose) is an enzymatic hydrolysis of the acetamide group. Production of the primary metabolite, ucb L057, is not supported by liver cytochrome P450 isoforms. Hydrolysis of the acetamide group was measurable in a large number of tissues including blood cells. The metabolite ucb L057 is pharmacologically inactive.

Two minor metabolites were also identified. One was obtained by hydroxylation of the pyrrolidone ring (1.6 % of the dose) and the other one by opening of the pyrrolidone ring (0.9 % of the dose).

Other unidentified components accounted only for 0.6 % of the dose.

No enantiomeric interconversion was evidenced in vivo for either levetiracetam or its primary metabolite.

In vitro, levetiracetam and its primary metabolite have been shown not to inhibit the major human liver cytochrome P450 isoforms (CYP3A4, 2A6, 2C9, 2C19, 2D6, 2E1 and 1A2), glucuronyl transferase (UGT1A1 and UGT1A6) and epoxide hydroxylase activities. In addition, levetiracetam does not affect the in vitro glucuronidation of valproic acid.

In human hepatocytes in culture, levetiracetam had little or no effect on CYP1A2, SULT1E1 or UGT1A1. Levetiracetam caused mild induction of CYP2B6 and CYP3A4. The in vitro data and in vivo interaction data on oral contraceptives, digoxin and warfarin indicate that no significant enzyme induction is expected in vivo. Therefore, the interaction of Lerace with other substances, or vice versa, is unlikely.

Elimination

The plasma half-life in adults was 7 ± 1 hours and did not vary either with dose, route of administration or repeated administration. The mean total body clearance was 0.96 ml/min/kg.

The major route of excretion was via urine, accounting for a mean 95 % of the dose (approximately 93 % of the dose was excreted within 48 hours). Excretion via faeces accounted for only 0.3 % of the dose.

The cumulative urinary excretion of levetiracetam and its primary metabolite accounted for 66 % and 24 % of the dose, respectively during the first 48 hours.

The renal clearance of levetiracetam and ucb L057 is 0.6 and 4.2 ml/min/kg respectively indicating that levetiracetam is excreted by glomerular filtration with subsequent tubular reabsorption and that the primary metabolite is also excreted by active tubular secretion in addition to glomerular filtration. Levetiracetam elimination is correlated to creatinine clearance.

6 Pharmaceutical particulars

6.1 List of excipients

Inactive Ingredients (Core Materials)

Corn Starch (Maize Starch)

Ac-Di-Sol (Cross Carmellose Sodium)

Magnesium Stearate

Aerosil 200 (Silicon Dioxide)

Povidone K-90

Deionized Water

Inactive Ingredients (Coated Materials)

Hydroxy Propyl Methyl Cellulose

Titanium Dioxide USP USP

Talcum Powder

Hydroxypropyl Cellulose

Polyethylene Glycol 6000

Deionized Water BP

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

24 months

6.4 Special precautions for storage

Store below 30oC. Protect from sunlight and moisture. Keep out of reach of children.

6.5 Nature and contents of container

Blister pack of printed Aluminium Foil with cold forming foil Packed In Printed Carton Along With Leaflet

6.6 Special precautions for disposal

Not Applicable

7. Marketing authorization holder

Hilton Pharma (Pvt.) Ltd

8th& 9th Floor, Progressive Plaza, Beaumont Road Karachi, Pakistan

8. Marketing authorization number

H2025/CTD8487/16656

9. Date of first authorization/ Renewal of authorization

21/10/2025

10. Date of revision of the text

21/10/2025