

1.17 Summary Product Characteristics (SPC)

Attached

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

MEROCEL (Meropenem for Injection USP 1 gm)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

MEROCEL (Meropenem for Injection USP 1 gm)

Each Vial Contains:

Meropenem USP eq. to Meropenem Anhydrous 1 gm

(also contains 90.2 mg of sodium as sodium carbonate USP)

3. PHARMACEUTICAL FORM

Powder for solution for injection or infusion

A white to off-white Colour powder, filled & sealed in colorless glass vial.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

MEROCEL is indicated for the treatment of the following infections in adults and children over 3 months of age.

- Pneumonia, including community acquired pneumonia and nosocomial pneumonia.
- Broncho-pulmonary infections in cystic fibrosis
- Complicated urinary tract infections
- Complicated intra-abdominal infections
- Intra- and post-partum infections
- Complicated skin and soft tissue infections
- Acute bacterial meningitis

MEROCEL may be used in the management of neutropenic patients with fever that is suspected to be due to a bacterial infection.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

The tables below provide general recommendations for dosing.

The dose of meropenem administered and the duration of treatment should take into account the type of infection to be treated, including its severity, and the clinical response.

A dose of up to 2,000 mg three times daily in adults and adolescents and a dose of up to 40 mg/kg three times daily in children may be particularly appropriate when treating some types of infections, such as nosocomial infections due to *Pseudomonas aeruginosa* or *Acinetobacter spp.*

Additional considerations for dosing are needed when treating patients with renal insufficiency (see further below).

Adults and adolescents

Infection	Dose to be administered every 8 hours
Pneumonia including community-acquired pneumonia and nosocomial pneumonia.	500 mg or 1,000 mg
Broncho-pulmonary infections in cystic fibrosis	2,000 mg
Complicated urinary tract infections	500 mg or 1,000 mg
Complicated intra-abdominal infections	500 mg or 1,000 mg
Intra- and post-partum infections	500 mg or 1,000 mg
Complicated skin and soft tissue infections	500 mg or 1,000 mg
Acute bacterial meningitis	2,000 mg
Management of febrile neutropenic patients	1,000 mg

Meropenem is usually given by intravenous infusion over approximately 15 to 30 minutes (see section 6.2, 6.3 and 6.6).

Alternatively, doses up to 1,000 mg can be given as an intravenous bolus injection over approximately 5 minutes. There are limited safety data available to support the administration of a 2,000 mg dose in adults as an intravenous bolus injection.

Renal impairment

The dose for adults and adolescents should be adjusted when creatinine clearance is less than 51 ml/min, as shown below. There are limited data to support the application of these dose adjustments for a unit dose of 2,000 mg.

Creatinine clearance (ml/min)	Dose (based on “unit” dose range of 500 mg or 1,000 mg or 2,000 mg, see table above)	Frequency
26-50	one unit dose	every 12 hours
10-25	half of one unit dose	every 12 hours
< 10	half of one unit dose	every 24 hours

Meropenem is cleared by haemodialysis and haemofiltration. The required dose should be administered after completion of the haemodialysis cycle.

There are no established dose recommendations for patients receiving peritoneal dialysis.

Hepatic impairment

No dose adjustment is necessary in patients with hepatic impairment (see section 4.4).

Elderly

No dose adjustment is required for the elderly with normal renal function or creatinine clearance values above 50 ml/min.

Paediatric population

Children under 3 months of age

The safety and efficacy of meropenem in children under 3 months of age have not been established and the optimal dose regimen has not been identified. However, limited pharmacokinetic data suggest that 20 mg/kg every 8 hours may be an appropriate regimen (see section 5.2).

Children from 3 months to 11 years of age and up to 50 kg body weight

The recommended dose regimens are shown in the table below:

Infection	Dose to be administered every 8 hours
Pneumonia including community-acquired pneumonia and nosocomial pneumonia	10 or 20 mg/kg
Broncho-pulmonary infections in cystic fibrosis	40 mg/kg
Complicated urinary tract infections	10 or 20 mg/kg
Complicated intra-abdominal infections	10 or 20 mg/kg
Complicated skin and soft tissue infections	10 or 20 mg/kg
Acute bacterial meningitis	40 mg/kg
Management of febrile neutropenic patients	20 mg/kg

Children over 50 kg body weight

The adult dose should be administered.

There is no experience in children with renal impairment.

Method of administration

Meropenem is usually given by intravenous infusion over approximately 15 to 30 minutes (see sections 6.2, 6.3, and 6.6). Alternatively, meropenem doses of up to 20 mg/kg may be given as an intravenous bolus over approximately 5 minutes. There are limited safety data available to support the administration of a 40 mg/kg dose in children as an intravenous bolus injection.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Hypersensitivity to any other carbapenem antibacterial agent.

Severe hypersensitivity (e.g. anaphylactic reaction, severe skin reaction) to any other type of betalactam antibacterial agent (e.g. penicillins or cephalosporins).

4.4 Special warnings and precautions for use

The selection of meropenem to treat an individual patient should take into account the appropriateness of using a carbapenem antibacterial agent based on factors such as severity of the infection, the prevalence of resistance to other suitable antibacterial agents and the risk of selecting for carbapenem-resistant bacteria.

As with all beta-lactam antibiotics, serious and occasionally fatal hypersensitivity reactions have been reported (see sections 4.3 and 4.8).

Patients who have a history of hypersensitivity to carbapenems, penicillins or other beta-lactam antibiotics may also be hypersensitive to meropenem. Before initiating therapy with meropenem, careful inquiry should be made concerning previous hypersensitivity reactions to beta-lactam antibiotics.

If a severe allergic reaction occurs, the medicinal product should be discontinued and appropriate measures taken.

Antibiotic-associated colitis and pseudomembranous colitis have been reported with nearly all anti-bacterial agents, including meropenem, and may range in severity from mild to life threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhoea during or subsequent to the administration of meropenem (see section 4.8). Discontinuation of therapy with meropenem and the administration of specific treatment for *Clostridium difficile* should be considered. Medicinal products that inhibit peristalsis should not be given.

Seizures have infrequently been reported during treatment with carbapenems, including meropenem (see section 4.8).

Hepatic function should be closely monitored during treatment with meropenem due to the risk of hepatic toxicity (hepatic dysfunction with cholestasis and cytolysis) (see section 4.8).

Use in patients with liver disease: patients with pre-existing liver disorders should have liver function monitored during treatment with meropenem. There is no dose adjustment necessary (see section 4.2).

A positive direct or indirect Coombs test may develop during treatment with meropenem.

The concomitant use of meropenem and valproic acid/sodium valproate is not recommended (see section 4.5).

4.5 Interaction with other medicinal products and other forms of interaction

No specific medicinal product interaction studies other than probenecid were conducted.

Probenecid competes with meropenem for active tubular secretion and thus inhibits the renal excretion of meropenem with the effect of increasing the elimination half-life and plasma concentration of meropenem. Caution is required if probenecid is co-administered with meropenem.

The potential effect of meropenem on the protein binding of other medicinal products or metabolism has not been studied. However, the protein binding is so low that no interactions with other compounds would be expected on the basis of this mechanism.

Decreases in blood levels of valproic acid have been reported when it is co-administered with carbapenem agents resulting in a 60-100 % decrease in valproic acid levels in about two days. Due to the rapid onset and the extent of the decrease, co-administration of valproic acid with carbapenem agents is not considered to be manageable and therefore should be avoided (see section 4.4).

Oral anti-coagulants

Simultaneous administration of antibiotics with warfarin may augment its anti-coagulant effects. There have been many reports of increases in the anti-coagulant effects of orally administered anti-coagulant agents, including warfarin in patients who are concomitantly receiving antibacterial agents. The risk may vary with the underlying infection, age and general status of the patient so that the contribution of the antibiotic to the increase in INR (international normalised ratio) is difficult to assess. It is recommended that the INR should be monitored frequently during and shortly after co-administration of antibiotics with an oral anti-coagulant agent.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no or limited amount of data from the use of meropenem in pregnant women.

Animal studies do not indicate direct or indirect harmful effects with respect to reproductive toxicity (see section 5.3).

As a precautionary measure, it is preferable to avoid the use of meropenem during pregnancy.

Breast-feeding

Small amounts of meropenem have been reported to be excreted in human milk. Meropenem is detectable at very low concentrations in animal milk. Meropenem should not be used in breast-feeding women unless the potential benefit for the mother justifies the potential risk to the baby.

4.7 Effects on ability to drive and use machines

No studies on the effect on the ability to drive and use machines have been performed.

4.8 Undesirable effects

Summary of the safety profile

In a review of 4,872 patients with 5,026 meropenem treatment exposures, meropenem-related adverse reactions most frequently reported were diarrhoea (2.3%), rash (1.4%), nausea/vomiting (1.4 %) and injection site inflammation (1.1%). The most commonly reported meropenem-related laboratory adverse events were thrombocytosis (1.6 %) and increased hepatic enzymes (1.5-4.3 %).

Tabulated risk of adverse reactions

Adverse reactions listed in the table with a frequency of “not known” were not observed in the 2,367 patients who were included in pre-authorisation clinical studies with intravenous and intramuscular meropenem but have been reported during the post-marketing period.

In the table below all adverse reactions are listed by system organ class and frequency: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$) and not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Table 1

System Organ Class	Frequency	Event
Infections and infestations	Uncommon	oral and vaginal candidiasis
Blood and lymphatic system disorders	Common	thrombocythaemia
	Uncommon	eosinophilia, thrombocytopenia, leucopenia, neutropenia
	Not known	agranulocytosis, haemolytic anaemia
Immune system disorders	Not known	angioedema, anaphylaxis (see sections 4.3 and 4.4)

Nervous system disorders	Common	headache
	Uncommon	paraesthesiae
	Rare	convulsions (see section 4.4)
Gastrointestinal disorders	Common	diarrhoea, vomiting, nausea, abdominal pain
	Not known	antibiotic-associated colitis (see section 4.4)
Hepatobiliary disorders	Common	transaminases increased, blood alkaline phosphatase increased, blood lactate dehydrogenase increased.
	Uncommon	blood bilirubin increased
Skin and subcutaneous tissue disorders	Common	rash, pruritis
	Uncommon	urticaria
	Not known	toxic epidermal necrolysis, Stevens Johnson syndrome, erythema multiforme, Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS Syndrome).
Renal and urinary disorders	Uncommon	blood creatinine increased, blood urea increased
General disorders and administration site conditions	Common	inflammation, pain
	Uncommon	thrombophlebitis
	Not known	pain at the injection site

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

4.9 Overdose

Relative overdose may be possible in patients with renal impairment if the dose is not adjusted as described in section 4.2. Limited post-marketing experience indicates that if adverse reactions occur following overdose, they are consistent with the adverse reaction profile described in section 4.8, are generally mild in severity and resolve on withdrawal or dose reduction. Symptomatic treatments should be considered.

In individuals with normal renal function, rapid renal elimination will occur.

Haemodialysis will remove meropenem and its metabolite.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antibacterials for systemic use, carbapenems, ATC code: J01DH02

Mechanism of action

Meropenem exerts its bactericidal activity by inhibiting bacterial cell wall synthesis in Gram-positive and Gram-negative bacteria through binding to penicillin-binding proteins (PBPs).

Pharmacokinetic/Pharmacodynamic (PK/PD) relationship

Similar to other beta-lactam antibacterial agents, the time that meropenem concentrations exceed the MIC (T>MIC) has been shown to best correlate with efficacy. In preclinical models meropenem demonstrated activity when plasma concentrations exceeded the MIC of the infecting organisms for approximately 40 % of the dosing interval. This target has not been established clinically.

Mechanism of resistance

Bacterial resistance to meropenem may result from: (1) decreased permeability of the outer membrane of Gram-negative bacteria (due to diminished production of porins) (2) reduced affinity of the target PBPs (3) increased expression of efflux pump components, and (4) production of beta-lactamases that can hydrolyse carbapenems.

Localised clusters of infections due to carbapenem-resistant bacteria have been reported in the European Union.

There is no target-based cross-resistance between meropenem and agents of the quinolone, aminoglycoside, macrolide and tetracycline classes. However, bacteria may exhibit resistance to more than one class of antibacterials agents when the mechanism involved include impermeability and/or an efflux pump(s).

Breakpoints

European Committee on Antimicrobial Susceptibility Testing (EUCAST) clinical breakpoints for MIC testing are presented below.

EUCAST clinical MIC breakpoints for meropenem (2009-06-05, v 3.1)		
Organism	Susceptible (S) (mg/l)	Resistant (R)(mg/l)
<i>Enterobacteriaceae</i>	≤ 2	> 8
<i>Pseudomonas</i>	≤ 2	> 8
<i>Acinetobacter</i>	≤ 2	> 8
<i>Streptococcus</i> groups A, B, C, G	≤ 2	> 2
<i>Streptococcus pneumoniae</i> ¹	≤ 2	> 2
Other streptococci	2	2
<i>Enterococcus</i>	--	--
<i>Staphylococcus</i> ²	note 3	note 3
<i>Haemophilus influenzae</i> ¹ and <i>Moraxella catarrhalis</i>	≤ 2	> 2
<i>Neisseria meningitidis</i> ^{2,4}	≤ 0.25	> 0.25
Gram-positive anaerobes	≤ 2	> 8
Gram-negative anaerobes	≤ 2	> 8
Non-species related breakpoints ⁵	≤ 2	> 8

¹ Meropenem breakpoints for *Streptococcus pneumoniae* and *Haemophilus influenzae* in meningitis are 0.25/1 mg/L.

² Strains with MIC values above the S/I breakpoint are rare or not yet reported. The identification and antimicrobial susceptibility tests on any such isolate must be repeated and if the result is confirmed the isolate sent to a reference laboratory. Until there is evidence regarding clinical response for confirmed isolates with MIC above the current resistant breakpoint (in italics) they should be reported as resistant.

³ Susceptibility of staphylococci to meropenem is inferred from the methicillin susceptibility.

⁴ Meropenem breakpoints in *Neisseria meningitidis* relates to meningitis only.

⁵ Non-species related breakpoints have been determined mainly from PK/PD data and are independent of the MIC distributions of specific species. They are for use for species not mentioned in the table and footnotes.

-- = Susceptibility testing not recommended as the species is a poor target for therapy with the medicinal product.

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

The following table of pathogens listed is derived from clinical experience and therapeutic guidelines.

Commonly susceptible species

Gram-positive aerobes

Enterococcus faecalis

Staphylococcus aureus (methicillin-susceptible)[‡]

Staphylococcus species (methicillin-susceptible) including *Staphylococcus epidermidis*

Streptococcus agalactiae (Group B)

Streptococcus milleri group (*S. anginosus*, *S. constellatus*, and *S. intermedius*)

Streptococcus pneumoniae

Streptococcus pyogenes (Group A)

Gram-negative aerobes

Citrobacter freundii

Citrobacter koseri

Enterobacter aerogenes

Enterobacter cloacae

Escherichia coli

Haemophilus influenzae

Klebsiella oxytoca

Klebsiella pneumoniae

Morganella morganii

Neisseria meningitidis

Proteus mirabilis

Proteus vulgaris

Serratia marcescens

Gram-positive anaerobes

Clostridium perfringens

Peptoniphilus asaccharolyticus

Peptostreptococcus species (including *P. micros*, *P. anaerobius*, *P. magnus*)

Gram-negative anaerobes

Bacteroides caccae

Bacteroides fragilis group

Prevotella bivia

Prevotella disiens

Species for which acquired resistance may be a problem

Gram-positive aerobes

Enterococcus faecium^{§†}

Gram-negative aerobes

Acinetobacter species

Burkholderia cepacia

Pseudomonas aeruginosa

Inherently resistant organisms

Gram-negative aerobes

Stenotrophomonas maltophilia

Legionella species

Other micro-organisms

Chlamydophila pneumoniae

Chlamydophila psittaci

Coxiella burnetii

Mycoplasma pneumoniae

[§]Species that show natural intermediate susceptibility

[£]All methicillin-resistant staphylococci are resistant to meropenem

[†]Resistance rate $\geq 50\%$ in one or more EU countries

5.2 Pharmacokinetic properties

In healthy subjects the mean plasma half-life is approximately 1 hour; the mean volume of distribution is approximately 0.25 l/kg (11-27 l) and the mean clearance is 287 ml/min at 250 mg falling to 205 ml/min at 2,000 mg. Doses of 500, 1,000 and 2,000 mg doses infused over 30 minutes give mean C_{\max} values of approximately 23, 49 and 115 $\mu\text{g/ml}$ respectively, corresponding AUC values were 39.3, 62.3 and 153 $\mu\text{g}\cdot\text{h/ml}$. After infusion over 5 minutes C_{\max} values are 52 and 112 $\mu\text{g/ml}$ after 500 and 1,000 mg doses respectively. When multiple doses are administered 8-hourly to subjects with normal renal function, accumulation of meropenem does not occur.

A study of 12 patients administered meropenem 1,000 mg 8 hourly post-surgically for intra-abdominal infections showed a comparable C_{max} and half-life to normal subjects but a greater volume of distribution 27 l.

Distribution

The average plasma protein binding of meropenem was approximately 2 % and was independent of concentration. After rapid administration (5 minutes or less) the pharmacokinetics are biexponential but this is much less evident after 30 minutes infusion. Meropenem has been shown to penetrate well into several body fluids and tissues: including lung, bronchial secretions, bile, cerebrospinal fluid, gynaecological tissues, skin, fascia, muscle, and peritoneal exudates.

Biotransformation

Meropenem is metabolised by hydrolysis of the beta-lactam ring generating a microbiologically inactive metabolite. In vitro meropenem shows reduced susceptibility to hydrolysis by human dehydropeptidase-I (DHP-I) compared to imipenem and there is no requirement to co-administer a DHP-I inhibitor.

Elimination

Meropenem is primarily excreted unchanged by the kidneys; approximately 70 % (50 –75 %) of the dose is excreted unchanged within 12 hours. A further 28% is recovered as the microbiologically inactive metabolite. Faecal elimination represents only approximately 2% of the dose. The measured renal clearance and the effect of probenecid show that meropenem undergoes both filtration and tubular secretion.

Renal insufficiency

Renal impairment results in higher plasma AUC and longer half-life for meropenem. There were AUC increases of 2.4 fold in patients with moderate impairment (CrCL 33-74 ml/min), 5 fold in severe impairment (CrCL 4-23 ml/min) and 10 fold in haemodialysis patients (CrCL <2 ml/min) when compared to healthy subjects (CrCL >80 ml/min). The AUC of the microbiologically inactive ring opened metabolite was also considerably increased in patients with renal impairment. Dose adjustment is recommended for patients with moderate and severe renal impairment (see section 4.2).

Meropenem is cleared by haemodialysis with clearance during haemodialysis being approximately 4 times higher than in anuric patients.

Hepatic insufficiency

A study in patients with alcoholic cirrhosis shows no effect of liver disease on the pharmacokinetics of meropenem after repeated doses.

Adult patients

Pharmacokinetic studies performed in patients have not shown significant pharmacokinetic differences versus healthy subjects with equivalent renal function. A population model developed from data in 79 patients with intra-abdominal infection or pneumonia, showed a dependence of the central volume on weight and the clearance on creatinine clearance and age.

Paediatrics

The pharmacokinetics in infants and children with infection at doses of 10, 20 and 40 mg/kg showed C_{max} values approximating to those in adults following 500, 1,000 and 2,000 mg doses, respectively. Comparison showed consistent pharmacokinetics between the doses and half-lives similar to those observed in adults in all but the youngest subjects (< 6 months $t_{1/2}$ 1.6 hours). The mean meropenem clearance values were 5.8 ml/min/kg (6-12 years), 6.2 ml/min/kg (2-5 years), 5.3 ml/min/kg (6-23 months) and 4.3 ml/min/kg (2-5 months). Approximately 60 % of the dose is excreted in urine over 12 hours as meropenem with a further 12 % as metabolite. Meropenem concentrations in the CSF of children with meningitis are approximately 20 % of concurrent plasma levels although there is significant inter-individual variability.

The pharmacokinetics of meropenem in neonates requiring anti-infective treatment showed greater clearance in neonates with higher chronological or gestational age with an overall average half-life of 2.9 hours. Monte Carlo simulation based on a population PK model showed that a dose regimen of 20 mg/kg 8 hourly achieved 60 % $T > MIC$ for *P. aeruginosa* in 95 % of pre-term and 91 % of full term neonates.

Elderly

Pharmacokinetic studies in healthy elderly subjects (65-80 years) have shown a reduction in plasma clearance, which correlated with age-associated reduction in creatinine clearance, and a smaller reduction in non-renal clearance. No dose adjustment is required in elderly patients, except in cases of moderate to severe renal impairment (see section 4.2).

5.3 Preclinical safety data

Animal studies indicate that meropenem is well tolerated by the kidney. Histological evidence of renal tubular damage was seen in mice and dogs only at doses of 2,000 mg/kg and above after a single administration and in monkeys at 500 mg/kg in a 7-day study.

Meropenem is generally well tolerated by the central nervous system. Effects were seen in acute toxicity studies in rodents at doses exceeding 1,000 mg/kg.

The IV LD_{50} of meropenem in rodents is greater than 2,000 mg/kg.

In repeat dose studies of up to 6 months duration only minor effects were seen including a decrease in red cell parameters in dogs.

There was no evidence of mutagenic potential in a conventional test battery and no evidence of reproductive toxicity including teratogenic potential in studies in rats up to 750 mg/kg and in monkeys up to 360 mg/kg.

There was increased evidence of abortions at 500 mg/kg in a preliminary study in monkeys.

There was no evidence of increased sensitivity to meropenem in juveniles compared to adult animals. The intravenous formulation was well tolerated in animal studies.

The sole metabolite of meropenem had a similar profile of toxicity in animal studies.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Sodium carbonate, anhydrous

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

6.3 Shelf life

2 years

After reconstitution:

The reconstituted solutions for intravenous injection or infusion should be used immediately. The time interval between the beginning of reconstitution and the end of intravenous injection or infusion should not exceed one hour.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions
Do not freeze the reconstituted solution.

6.5 Nature and contents of container

MEROCEL (Meropenem for Injection USP 1 gm)

1,348 mg powder in a 30 ml type I colourless glass vials with butyl rubber (type I) closures and sealed with aluminium caps.

Store protected from moisture at a temperature not exceeding 25°C.

6.6 Special precautions for disposal and other handling

Injection

Meropenem to be used for bolus intravenous injection should be constituted with sterile water for injection (50 mg/mL).

Infusion

For intravenous infusion meropenem vials can be reconstituted with sterile water for injection or 0.9 % sodium chloride or 5% glucose solutions for infusion. The solution formed will be further diluted with 0.9 % sodium chloride or 5% glucose solutions (concentration ranging from 1 to 20 mg/mL).

For storage conditions of the reconstituted medicinal product, see section 6.3.

Each vial is for single use only.

Standard aseptic techniques should be used for solution preparation and administration.

Shake well until dissolved completely.

The solutions should be inspected visually for particles prior to administration. Only clear pale yellow solution, free from visible particles should be used.

Any unused product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Applicant:

PROMED INDUSTRIES LIMITED

West lands Avenue

David Osieri Road, Off Waiyaki Way,

Nairobi, 00200

Kenya

Manufacturer:

Systochem Laboratories Ltd.

B-75, Roop Nagar, Industrial Area,

Loni-201 102 (UP) (India)

8. MARKETING AUTHORISATION NUMBER(S)

9. DATE OF FIRST AUTHORISATION/ RENEWAL OF THE AUTHORISATION

10. DATE OF REVISION OF THE TEXT

08/04/2021