

Summary of Product Characteristics

1. Name of the Medical Product

Monoflam Suspension

2. Qualitative & Quantitative Composition:

Each 5ml contains;

Diclofenac free acid.....9mg

Excipients of known effect

Sorbitol 70%

For the full list of excipients, see section 6.1.

3. Pharmaceutical Form:

Oral syrup

Visual Description

A colourless syrup

4. Clinical Particulars

4.1 Therapeutic Indications:

Adults and elderly: Relief of all grades of pain and inflammation in a wide range of conditions, including: (i) arthritic conditions: rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, acute gout, (ii) and other acute musculo-skeletal disorders such as peri-arthritis (for example frozen shoulder), tendinitis, tenosynovitis, bursitis, (iii) other painful conditions resulting from trauma, including fracture, low back pain, sprains, strains, dislocations, orthopaedic, dental and other minor surgery.

Children: Children aged 1-12 years: for treatment of juvenile chronic arthritis

Children aged 9 years and above: for short term treatment of fever related to infection of the ear, nose or throat (ENT), e.g. pharyngotonsillitis, otitis media. As monotherapy or as adjunct therapy with morphine or other opiates (due to its opiate-sparing effect) for the relief of acute post-operative pain.

4.2 Posology and Method of administration:

Posology

Posology

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (see section 4.4).

75 mg to 150 mg daily in two or three divided doses.

The recommended maximum daily dose of diclofenac is 150mg.

Special populations Elderly:

Although the pharmacokinetics of Diclofenac are not impaired to any clinically relevant extent in elderly patients, nonsteroidal anti-inflammatory drugs should be used with particular caution in such patients who generally are more prone to adverse reactions. In particular it is recommended that the lowest effective dosage be used in

frail elderly patients or those with a low body weight (see also precautions) and the patient should be monitored for GI bleeding during NSAID therapy.

Cardiovascular and significant cardiovascular risk factors

Diclofenac is contraindicated in patients with established congestive heart failure (NYHA II-IV), ischemic heart disease, peripheral arterial disease and/or cerebrovascular disease (see section 4.3 Contraindications).

Patients with congestive heart failure (NYHA-I) or significant risk factors for cardiovascular disease should be treated with diclofenac only after careful consideration. Since cardiovascular risks with diclofenac may increase with dose and duration of exposure, the lowest effective daily dose should be used and for the shortest duration possible (see section 4.4 Special warnings and precautions for use).

Renal impairment: Diclofenac is contraindicated in patients with severe renal impairment (see section 4.3). No specific studies have been carried out in patients with renal impairment, therefore, no specific dose adjustment recommendations can be made. Caution is advised when administering diclofenac to patients with mild to moderate renal impairment (see section 4.3 and 4.4).

Hepatic impairment: Diclofenac is contraindicated in patients with severe hepatic impairment (see section 4.3). No specific studies have been carried out in patients with hepatic impairment, therefore, no specific dose adjustment recommendations can be made. Caution is advised when administering diclofenac to patients with mild to moderate hepatic impairment (see section 4.3 and 4.4).

Paediatric Population:

For treatment of Juvenile chronic arthritis the following dosage should be given: Children (aged 1-12 years): 1 to 3-mg/kg per day in divided doses. For the short term treatment of fever related to infection of the ear, nose or throat (ENT) and post-operative pain the following dosage should be given: Children aged 9 years (min. 35 kg body weight) or over and adolescents should be given up to 2 mg/kg body weight per day in 3 divided doses, depending on the severity of the disorder.

Method of administration

For oral administration

To be taken preferably with or after food.

4.3 Contraindications:

- Hypersensitivity to Diclofenac sodium or to any of the excipients listed in section 6.1.
- Active or history of recurrent peptic ulceration / haemorrhage (two or more distinct episodes of proven ulceration or bleeding).
- A history of gastrointestinal bleeding or perforation, related to previous NSAIDs therapy.
- Active, or gastric or intestinal ulcer, bleeding or perforation.
- During the last trimester of pregnancy (see section 4.6 – Pregnancy and lactation)
- Hepatic failure
- Renal failure
- Established congestive heart failure (NYHA-II-IV), ischemic heart disease, peripheral arterial disease and/or cerebrovascular disease.
- Like other non-steroidal anti-inflammatory drugs (NSAIDs), diclofenac is also contraindicated in patients in whom attacks of asthma, angiodema, urticaria or

acute rhinitis are precipitated by ibuprofen, acetylsalicylic acid or other nonsteroidal anti-inflammatory drugs.

4.4 Special warnings & precautions for use

General

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms (see section 4.2, and GI and cardiovascular risks below).

The concomitant use of diclofenac with systemic NSAIDs including cyclooxygenase-2 selective inhibitors should be avoided due to the absence of any evidence demonstrating synergistic benefits and the potential for additive undesirable effects (see section 4.5).

Caution is indicated in the elderly on basic medical grounds. In particular, it is recommended that the lowest effective dose be used in frail elderly patients or those with a low body weight (see section 4.2).

As with other nonsteroidal anti-inflammatory drugs including diclofenac, allergic reactions, including anaphylactic/anaphylactoid reactions, can also occur without earlier exposure to the drug (see section 4.8). Hypersensitivity reactions can also progress to Kounis syndrome, a serious allergic reaction that can result in myocardial infarction. Presenting symptoms of such reactions can include chest pain occurring in association with an allergic reaction to diclofenac.

Like other NSAIDs, diclofenac may mask the signs and symptoms of infection due to its pharmacodynamic properties.

Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine. This medicine contains less than 1 mmol sodium (23 mg) per 25 mg tablet, that is to say essentially 'sodium-free'. This medicine also contains Sunset Yellow (E110) which may cause allergic reactions.

Gastrointestinal effects

Gastrointestinal bleeding (haematemesis, melaena), ulceration or perforation, which can be fatal has been reported with all NSAIDs including diclofenac and may occur at any time during treatment, with or without warning symptoms or a previous history of serious gastrointestinal (GI) events. They generally have more serious consequences in the elderly. If gastrointestinal bleeding or ulceration occurs in patients receiving diclofenac, the medicinal product should be withdrawn. As with all NSAIDs, including diclofenac, close medical surveillance is imperative and particular caution should be exercised when prescribing diclofenac in patients with symptoms indicative of gastrointestinal disorders, or with a history suggestive of gastric or intestinal ulceration, bleeding or perforation (see section 4.8). The risk of GI bleeding, ulceration or perforation is higher with increasing NSAID doses including diclofenac and in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation.

The elderly have an increased frequency of adverse reactions to NSAIDs especially gastrointestinal bleeding and perforation which may be fatal (see section 4.2). To reduce the risk of GI toxicity in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation, and in the elderly, the treatment should be initiated and maintained at the lowest effective dose. Combination therapy with protective agents (e.g. misoprostol or proton pump inhibitors) should be considered for these patients, and also for patients requiring concomitant use of

medicinal products containing low dose acetylsalicylic acid (ASA/aspirin), or other medicinal products likely to increase gastrointestinal risk (see below and section 4.5).

Patients with a history of GI toxicity, particularly the elderly, should report any unusual abdominal symptoms (especially GI bleeding).

Caution is recommended in patients receiving concomitant medications which could increase the risk of ulceration or bleeding, such as systemic corticosteroids, anticoagulants such as warfarin, selective serotonin-reuptake inhibitors (SSRIs) or anti-platelet agents such as acetylsalicylic acid (see section 4.5).

Close medical surveillance and caution should also be exercised in patients with ulcerative colitis or Crohn's disease, as their condition may be exacerbated (see section 4.8).

NSAIDs, including diclofenac, may be associated with increased risk of gastrointestinal anastomotic leak. Close medical surveillance and caution are recommended when using diclofenac after gastro-intestinal surgery. *Hepatic impairment*

Close medical surveillance is required when prescribing diclofenac to patients with impairment of hepatic function, as their condition may be exacerbated.

As with other NSAIDs, including diclofenac, values of one or more liver enzymes may increase. During prolonged treatment with diclofenac, regular monitoring of hepatic function is indicated as a precautionary measure. If abnormal liver function tests persist or worsen, clinical signs or symptoms consistent with liver disease develop or if other manifestations occur (eosinophilia, rash), diclofenac should be discontinued.

Hepatitis may occur with diclofenac without prodromal symptoms.

Caution is called for when using diclofenac in patients with hepatic porphyria, since it may trigger an attack. *Renal impairment*

As fluid retention and oedema have been reported in association with NSAID therapy, including diclofenac, particular caution is called for in patients with impaired cardiac or renal function, history of hypertension, the elderly, patients receiving concomitant treatment with diuretics or medicinal products that can significantly impact renal function, and in those patients with substantial extracellular volume depletion from any cause, e.g. before or after major surgery (see 4.3). Monitoring of renal function is recommended as a precautionary measure when using diclofenac in such cases. Discontinuation of therapy is usually followed by recovery to the pre-treatment state. *Skin reactions*

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, toxic epidermal necrolysis, and generalised bullous fixed drug eruption have been reported very rarely in association with the use of NSAIDs, including diclofenac (see section 4.8). Patients appear to be at highest risk for these reactions early in the course of therapy: the onset of the reaction occurring in the majority of cases within the first month of treatment. Diclofenac sodium tablets should be discontinued at the first appearance of skin rash, mucosal lesions or any other signs of hypersensitivity.

SLE and mixed connective tissue disease

In patients with systemic lupus erythematosus (SLE) and mixed connective tissue disorders there may be an increased risk of aseptic meningitis (see section 4.8).

Cardiovascular and cerebrovascular effects

Patients with congestive heart failure (NYHA-1) or patients with significant risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes

mellitus, smoking) should only be treated with diclofenac after careful consideration.

As the cardiovascular risks of diclofenac may increase with dose and duration of exposure, the shortest duration possible and the lowest effective daily dose should be used. The patient's need for symptomatic relief and response to therapy should be re-evaluated periodically. Appropriate monitoring and advice are required for patients with a history of hypertension and congestive heart failure (NYHA-1) as fluid retention and oedema have been reported in association with NSAID therapy including diclofenac.

Clinical trial and epidemiological data consistently point towards increased risk of arterial thrombotic events (for example myocardial infarction or stroke) associated with the use of diclofenac, particularly at high dose (150mg daily) and in long term treatment.

Patients should remain alert for the signs and symptoms of serious arteriothrombotic events (e.g. chest pain, shortness of breath, weakness, slurring of speech), which can occur without warnings. Patients should be instructed to see a physician immediately in case of such an event

Haematological effects

During prolonged treatment with diclofenac, as with other NSAIDs, monitoring of the blood count is recommended. Diclofenac may reversibly inhibit platelet aggregation (see anticoagulants in section 4.5). Patients with defects of haemostasis, bleeding diathesis or haematological abnormalities should be carefully monitored.

Pre-existing asthma

In patients with asthma, seasonal allergic rhinitis, swelling of the nasal mucosa (i.e. nasal polyps), chronic obstructive pulmonary diseases or chronic infections of the respiratory tract (especially if linked to allergic rhinitis-like symptoms), reactions on NSAIDs like asthma exacerbations (so-called intolerance to analgesics / analgesics-asthma), Quincke's oedema or urticaria are more frequent than in other patients. Therefore, special precaution is recommended in such patients (readiness for emergency). This is applicable as well for patients who are allergic to other substances, e.g. with skin reactions, pruritus or urticaria.

Like other drugs that inhibit prostaglandin synthetase activity, diclofenac sodium and other NSAIDs can precipitate bronchospasm if administered to patients suffering from, or with a previous history of bronchial asthma.

Female fertility:

The use of Diclofenac may impair female fertility and is not recommended in women attempting to conceive. In women who may have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of Diclofenac should be considered (see section 4.6).

4.5 Interactions with other medicinal products and other forms of Interactions

The following interactions include those observed with diclofenac gastroresistant tablets and/or other pharmaceutical forms of diclofenac.

Lithium: If used concomitantly, diclofenac may raise plasma concentrations of lithium. Monitoring of the serum lithium level is recommended.

Digoxin: If used concomitantly, diclofenac may raise plasma concentrations of digoxin. Monitoring of the serum digoxin level is recommended.

Diuretics and Anti-hypertensive agents: Like other NSAIDs, concomitant use of diclofenac with diuretics or antihypertensive agents (e.g. beta-blockers, angiotensin

converting enzyme (ACE) inhibitors) may cause a decrease in their antihypertensive effect via inhibition of vasodilatory prostaglandin synthesis.

Therefore, the combination should be administered with caution and patients, especially the elderly, should have their blood pressure periodically monitored. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy and periodically thereafter, particularly for diuretics and ACE inhibitors due to the increased risk of nephrotoxicity.

Drugs known to cause hyperkalemia: Concomitant treatment with potassium-sparing diuretics, ciclosporin, tacrolimus or trimethoprim may be associated with increased serum potassium levels, which should therefore be monitored frequently (see section 4.4).

Anticoagulants and anti-platelet agents: Caution is recommended since concomitant administration could increase the risk of bleeding (see section 4.4). Although clinical investigations do not appear to indicate that diclofenac affects the action of anticoagulants, there are reports of an increased risk of haemorrhage in patients receiving diclofenac and anticoagulants concomitantly (see section 4.4). Therefore, to be certain that no change in anticoagulant dosage is required, close monitoring of such patients is required. As with other nonsteroidal anti-inflammatory agents, diclofenac in high dose can reversibly inhibit platelet aggregation.

Other NSAIDs including cyclo-oxygenase-2 selective inhibitors and corticosteroids: Co-administration of diclofenac and other systemic NSAIDs or corticosteroids may increase the risk of gastrointestinal bleeding or ulceration. Avoid concomitant use of two or more NSAIDs (see section 4.4).

Selective serotonin reuptake inhibitors (SSRIs): Concomitant administration of SSRIs may increase the risk of gastrointestinal bleeding (see section 4.4).

Antidiabetics: Clinical studies have shown that diclofenac can be given together with oral antidiabetic agents without influencing their clinical effect. However, there have been isolated reports of hypoglycaemic and hyperglycaemic effects necessitating changes in the dosage of the antidiabetic agents during treatment with diclofenac. For this reason, monitoring of the blood glucose level is recommended as a precautionary measure during concomitant therapy.

Methotrexate: Diclofenac can inhibit the tubular renal clearance of methotrexate hereby increasing methotrexate levels. Caution is recommended when NSAIDs, including diclofenac, are administered less than 24 hours before treatment with methotrexate, since blood concentrations of methotrexate may rise and the toxicity of this substance be increased.

Cases of serious toxicity have been reported when methotrexate and NSAIDs including diclofenac are given within 24 hours of each other. This interaction is mediated through accumulation of methotrexate resulting from impairment of renal excretion in the presence of the NSAID.

Ciclosporin: Diclofenac, like other NSAIDs, may increase the nephrotoxicity of ciclosporin due to the effect on renal prostaglandins. Therefore, it should be given at doses lower than those that would be used in patients not receiving ciclosporin.

Tacrolimus: Possible increased risk of nephrotoxicity when NSAIDs are given with tacrolimus. This might be mediated through renal antiprostaglandin effects of both NSAID and calcineurin inhibitor.

Quinolone antimicrobials: Convulsions may occur due to an interaction between quinolones and NSAIDs. This may occur in patients with or without a previous history of epilepsy or convulsions. Therefore, caution should be exercised when considering the use of a quinolone in patients who are already receiving an NSAID.

Phenytoin: When using phenytoin concomitantly with diclofenac, monitoring of phenytoin plasma concentrations is recommended due to an expected increase in exposure to phenytoin.

Colestipol and cholestyramine: These agents can induce a delay or decrease in absorption of diclofenac. Therefore, it is recommended to administer diclofenac at least one hour before or 4 to 6 hours after administration of colestipol/cholestyramine.

Cardiac glycosides: Concomitant use of cardiac glycosides and NSAIDs in patients may exacerbate cardiac failure, reduce GFR and increase plasma glycoside levels.

Mifepristone: NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone.

Potent CYP2C9 inhibitors: Caution is recommended when co-prescribing diclofenac with potent CYP2C9 inhibitors (such as sulfinpyrazone voriconazole), which could result in a significant increase in peak plasma concentration and exposure to diclofenac due to inhibition of diclofenac metabolism.

4.6 Pregnancy and Lactation

Pregnancy:

Inhibition of prostaglandin synthesis may adversely affect the pregnancy and/or the embryo/foetal development. Data from epidemiological studies suggest an increased risk of miscarriage and of cardiac malformation and gastroschisis after use of a prostaglandin synthesis inhibitor in early pregnancy. The absolute risk for cardiovascular malformation was increased from less than 1%, up to approximately 1.5 %.

The risk is believed to increase with dose and duration of therapy. In animals, administration of a prostaglandin synthesis inhibitor has been shown to result in increased pre- and post-implantation loss and embryo-foetal lethality.

In addition, increased incidences of various malformations, including cardiovascular, have been reported in animals given a prostaglandin synthesis inhibitor during the organogenetic period.

From the 20th week of pregnancy onward, diclofenac use may cause oligohydramnios resulting from foetal renal dysfunction. This may occur shortly after treatment initiation and is usually reversible upon discontinuation. In addition, there have been reports of ductus arteriosus constriction following treatment in the second trimester, most of which resolved after treatment cessation. Therefore, during the first and second trimester of pregnancy, diclofenac should not be given unless clearly necessary. If diclofenac is used by a woman attempting to conceive, or during the first and second trimester of pregnancy, the dose should be kept as low and duration of treatment as short as possible. Antenatal monitoring for oligohydramnios should be considered after exposure to diclofenac for several days from gestational week 20 onward. Diclofenac Sodium 25mg Tablets should be discontinued if oligohydramnios or ductus arteriosus constriction are found. During the third trimester of pregnancy, all prostaglandin synthesis inhibitors may expose the foetus to:

- cardiopulmonary toxicity (premature constriction/closure of the ductus arteriosus and pulmonary hypertension);
- renal dysfunction (see above); the mother and the neonate, at the end of pregnancy, to:
- possible prolongation of bleeding time, an anti-aggregating effect which may occur even at very low doses.
- inhibition of uterine contractions resulting in delayed or prolonged labour.

Consequently, diclofenac sodium tablets are contraindicated during the third trimester of pregnancy (see sections 4.3 and 5.3).

Breast-feeding:

Like other NSAIDs, diclofenac passes into the breast milk in small amounts. Therefore, diclofenac should not be administered during breast feeding in order to avoid undesirable effects in the infant (see section 5.2)

Female Fertility

As with other NSAIDs, the use of diclofenac may impair female fertility and is not recommended in women attempting to conceive. In women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of diclofenac should be considered (see also section 4.4, regarding female fertility).

4.7 Effects on ability to drive and use machine*:

Patients who experience visual disturbances, dizziness, vertigo, somnolence, central nervous system disturbances, drowsiness or fatigue while taking NSAIDs should refrain from driving or operate machinery

4.8 Undesirable Effects*:

Adverse reactions (Table 1) are ranked under heading of frequency, the most frequent first, using the following convention: very common: ($>1/10$); common ($\geq 1/100$, $<1/10$); uncommon ($\geq 1/1,000$, $<1/100$); rare ($\geq 1/10,000$, $<1/1,000$); very rare ($<1/10,000$); Not known: cannot be estimated from the available data.

The following undesirable effects include those reported with either short-term or long-term use.

Table 1

Blood and lymphatic system disorders	
Very rare	Thrombocytopenia, leucopenia, anaemia (including haemolytic and aplastic anaemia), agranulocytosis.
Immune system disorders	
Rare	Hypersensitivity, anaphylactic and anaphylactoid reactions (including hypotension and shock).
Very rare	Angioneurotic oedema (including face oedema).
Psychiatric disorders	
Very rare	Disorientation, depression, insomnia, nightmare, irritability, psychotic disorder.

<u>Nervous system disorders</u>		
<u>Common</u>	<u>Headache, dizziness.</u>	
<u>Rare</u>	<u>Somnolence, tiredness.</u>	
<u>Very rare</u>	<u>Paraesthesia, memory impairment, convulsion, anxiety,</u>	
<u>Unknown</u>	<u>trem aseptic meningitis, taste disturbances,</u>	
	<u>cerebrovascular accid</u>	
	<u>Confusion, hallucinations, disturbances of sensation,</u>	
	<u>malaise.</u>	
<u>Eye disorders</u>		
<u>Very rare</u>	<u>Visual disturbance, vision blurred, diplopia.</u>	
<u>Unknown</u>	<u>Optic neuritis.</u>	
<u>Ear and labyrinth disorders</u>		
<u>Common</u>	<u>Vertigo.</u>	
<u>Very rare</u>	<u>Tinnitus, hearing impaired.</u>	
<u>Cardiac disorders</u>		
<u>Uncommon*</u>	<u>Myocardial infarction, cardiac failure, palpitations, chest</u>	
	<u>pain</u>	
<u>Unknown</u>	<u>Kounis syndrome</u>	
<u>Vascular disorders</u>		
<u>Very rare</u>	<u>Hypertension, hypotension, vasculitis.</u>	
<u>Respiratory, thoracic and mediastinal disorders</u>		
<u>Rare</u>	<u>Asthma (including</u>	<u>dyspnoea).</u>
<u>Very rare</u>	<u>Pneumonitis.</u>	
<u>Gastrointestinal disorders</u>		

**The*

Common	Nausea, vomiting, diarrhoea, dyspepsia, abdominal pain, flatulence, anorexia.
Rare	Gastritis, gastrointestinal haemorrhage, haematemesis, diar haemorrhagic, melaena, gastrointestinal ulcer with or without bleeding or perforation (sometimes fatal particularly in the elderly).
Very rare	Colitis (including haemorrhagic colitis and exacerbation of ulcerative colitis or Crohn's disease), constipation, stomatitis (including ulcerative stomatitis), glossitis, oesophageal diso diaphragm-like intestinal strictures, pancreatitis.
Unknown	Ischaemic colitis
Hepatobiliary disorders	
Common	Transaminases increased.
Rare	Hepatitis, jaundice, liver disorder.
Very rare	Fulminant hepatitis, hepatic necrosis, hepatic failure.
Skin and subcutaneous tissue disorders	
Common	Rash.
Rare	Urticaria.
Very rare	Bullous eruptions, eczema, erythema, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis (Lyell syndrome), dermatitis exfoliative, loss of hair, photosensitivit reaction, purpura , allergic purpura, pruritus.
Unknown	Fixed drug eruption, Generalised bullous fixed drug eruption
Renal and urinary disorders	
Very rare	Acute renal failure, haematuria, proteinuria, nephrotic syndro interstitial nephritis, renal papillary necrosis.
Reproductive system and breast disorders	
Very rare	Impotence
General disorders and administration site conditions	
Rare	Oedema

frequency reflects data from long-term treatment with a high dose (150mg/ day).

Clinical trial and epidemiological data consistently point towards an increased risk of arterial thrombotic events (for example myocardial infarction or stroke) associated with the use of diclofenac, particularly at high dose (150 mg daily) and in long term treatment (see sections 4.3 and 4.4).

Reporting of suspected adverse reactions: Healthcare professionals are requested to

report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Reporting System (PvERS) <https://pv.pharmacyboardkenya.org>.

4.9 Overdose:

Symptoms

There is no typical clinical picture resulting from diclofenac over dosage. Over dosage can cause symptoms such as headache, nausea, vomiting, epigastric pain, gastrointestinal haemorrhage, diarrhoea, dizziness, disorientation, excitation, coma, drowsiness, tinnitus, fainting or convulsions. In the case of significant poisoning acute renal failure and liver damage are possible.

Therapeutic measures

Management of acute poisoning with NSAIDs, including diclofenac, essentially consists of supportive measures and symptomatic treatment. Supportive measures and symptomatic treatment should be given for complications such as hypotension, renal failure, convulsions, gastrointestinal disorder, and respiratory depression. Special measures such as forced diuresis, dialysis or haemo-perfusion are probably of no help in eliminating NSAIDs, including diclofenac, due to the high protein binding and extensive metabolism.

Activated charcoal may be considered after ingestion within one hour of a potentially toxic overdose, and gastric decontamination (e.g. vomiting, gastric lavage) after ingestion of a potentially life threatening overdose.

5. Pharmacological properties:

5.1 Pharmacodynamic Properties:

Pharmacodynamic properties Pharmacotherapeutic group

Non-steroidal anti-inflammatory drugs (NSAIDs).

Mechanism of action

Diclofenac Sodium is a non-steroidal agent with marked analgesic/antiinflammatory and antipyretic properties. It is an inhibitor of prostaglandin synthetase, (cyclo-oxygenase).

Diclofenac sodium *in vitro* does not suppress proteoglycan biosynthesis in cartilage at concentrations equivalent to the concentrations reached in human beings.

There is limited clinical trial experience of the use of diclofenac in JRA/JIA paediatric patients. In a randomised, double-blind, 2-week, parallel group study in children aged 3-15 years with JRA/JIA, the efficacy and safety of daily 2-3 mg/kg BW diclofenac was compared with acetylsalicylic acid (ASS, 50-100 mg/kg BW/d) and placebo - 15 patients in each group. In the global evaluation, 11 of 15 diclofenac patients, 6 of 12 aspirin and 4 of 15 placebo patients showed improvement with the difference being statistically significant ($p < 0.05$). The number of tender joints decreased with diclofenac and ASS but increased with placebo. In a second randomised, double-blind, 6-week, parallel group study in children aged 4-15 years with JRA/JIA, the efficacy of diclofenac (daily dose 2-3 mg/kg BW, n=22) was comparable with that of indomethacin (daily dose 2-3 mg/kg BW, n=23).

5.2 Pharmacokinetics Properties*:

Absorption

Absorption is complete but onset is delayed until passage through the stomach, which may be affected by food which delays stomach emptying. The mean peak plasma diclofenac concentration reached at about 2 hours (50mg dose produces 1511 ± 466 ng/ml).

Bioavailability:

About half of the administered diclofenac is metabolised during its first passage through the liver ("first-pass" effect), the area under the concentrations curve (AUC) following oral administration is about half that following an equivalent parenteral dose.

Pharmacokinetic behaviour does not change on repeated administration. Accumulation does not occur, provided the recommended dosage intervals are observed. The plasma concentrations attained in children given equivalent doses (mg/kg, b.w.) are similar to those obtained in adults.

Distribution

The active substance is 99.7% protein bound, mainly to albumin (99.4%).

Diclofenac enters the synovial fluid, where maximum concentrations are measured 2-4 hours after the peak plasma values have been attained. The apparent half-life for elimination from the synovial fluid is 3-6 hours. Two hours after reaching the peak plasma values, concentrations of the active substance are already higher in the synovial fluid than they are in the plasma and remain higher for up to 12 hours.

Diclofenac was detected in a low concentration (100 ng/mL) in breast milk in one nursing mother. The estimated amount ingested by an infant consuming breast milk is equivalent to a 0.03 mg/kg/day dose (see section 4.6).

Metabolism

Biotransformation of diclofenac takes place partly by glucuronidation of the intact molecule, but mainly by single and multiple hydroxylation and methoxylation, resulting in several phenolic metabolites, most of which are converted to glucuronide conjugates. Two phenolic metabolites are biologically active, but to a much lesser extent than diclofenac.

Elimination

The total systemic clearance of diclofenac in plasma is 263 ± 56 mL/min (mean value $\pm$ SD). The terminal half-life in plasma is 1-2 hours. Four of the metabolites, including the two active ones, also have short plasma half-lives of 1-3 hours.

About 60% of the administered dose is excreted in the urine in the form of the glucuronide conjugate of the intact molecule and as metabolites, most of which are also converted to glucuronide conjugates. Less than 1% is excreted as unchanged substance. The rest of the dose is eliminated as metabolites through the bile in the faeces.

Characteristics in patients

Elderly: No relevant age-dependent differences in the drug's absorption, metabolism, or excretion have been observed, other than the finding that in five elderly patients, a 15 minute iv infusion resulted in 50% higher plasma concentrations than expected with young healthy subjects.

Patients with renal impairment: In patients suffering from renal impairment, no accumulation of the unchanged active substance can be inferred from the single-dose kinetics when applying the usual dosage schedule. At a creatinine clearance of less than 10 mL/min, the calculated steady-state plasma levels of the hydroxy metabolites are about 4 times higher than in normal subjects. However, the metabolites are ultimately cleared through the bile.

Patients with hepatic impairment: In patients with chronic hepatitis or nondecompensated cirrhosis, the kinetics and metabolism of diclofenac are the same as in patients without liver disease.

5.3 Preclinical safety data

Not stated.

6. Pharmaceutical particulars:**6.1 List of Excipients:**

Sorbitol 70% solution
Citric acid
Methyl paraben
Propyl paraben
Sodium saccharine
Propylene glycol
Polysorbate 80
Colloidal silicon dioxide
Hydroxy ethyl cellulose
Xanthan gum
Pineapple flavour
Mixed fruit flavour

6.2 Incompatibilities:

None reported

6.3 Shelf life:

36 months

6.4 Special Precautions for storage:

Do not store above 30°C. Store in the original package.

6.5 Nature and contents of container:

60ml bottle

6.6 Special precautions for disposal:

Not applicable

7. Marketing Authorization Holder:

Krishna Chemists Ltd
3rd Floor Metrix Hardware, Lusaka Road, Industrial Area.

Manufacturing Site Address:

Relax Biotech Pvt. Limited
862/1, Gidc, Makarpura, Vadodara - 390010, Gujarat, India

8. Marketing Authorization Numbers:

H2022/CTD7843/14999

9. Date of first registration /renewal of the registration:

2022 July 14

10. Date of revision of text:

2022 July 14