

Summary of Product Characteristics

MYFORTIC[®] (mycophenolic acid as mycophenolate sodium) 180 mg and 360 mg gastro-resistant tablets Tradename
MYFORTIC 180 mg and 360 mg gastro-resistant tablets.

1 Qualitative and quantitative

composition Active substance

Each gastro-resistant tablet contains 180 mg or 360 mg mycophenolic acid (MPA) equivalent to 192.4 mg and 384.8 mg mycophenolate sodium.

Excipients

Maize starch; povidone (K-30); crospovidone; lactose; colloidal silicon dioxide; magnesium stearate
For full list of excipients, see section 6.1

2 Pharmaceutical form

Gastro-resistant tablets

3 Clinical particulars

3.1 Therapeutic indications

Myfortic is indicated in combination with ciclosporin for microemulsion and corticosteroids for the prophylaxis of acute transplant rejection in patients receiving allogeneic renal transplants.

3.2 Posology and method of administration

The recommended dose is 720 mg (four 180 mg or two 360 mg Myfortic gastro-resistant tablets) twice daily (1,440 mg daily dose). In patients receiving 2 g mycophenolate mofetil (MMF), treatment can be replaced by 720 mg twice daily (1,440 mg daily dose) of Myfortic.

General target population

Treatment with Myfortic should be initiated and maintained by appropriately qualified transplant specialists.

Myfortic should be initiated in *de novo* patients within 48 hours following transplantation. Myfortic can be taken with or without food.

Special populations

Renal impairment

No dose adjustments are needed in patients experiencing delayed post-operative renal graft function (see section 11 Clinical Pharmacology). Patients with severe chronic renal impairment (glomerular filtration rate $<25 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$) should be

carefully monitored. Hepatic impairment

No dose adjustments are needed for renal transplant patients with severe hepatic parenchymal disease.

Pediatric patients (below 18 years)

Safety and efficacy in pediatric patients have not been established. Limited pharmacokinetic data are available for pediatric renal transplant patients (see section 11 Clinical Pharmacology).

Geriatric patients (65 years of age or above)

No dose adjustment is required in this patient population.

Treatment during rejection episodes

Renal transplant rejection does not affect mycophenolic acid pharmacokinetics; dosage reduction or interruption of Myfortic is not required.

Method of administration

Myfortic tablets should not be crushed in order to maintain the integrity of the enteric coating (see sections 11 Clinical pharmacology and 14 Pharmaceutical information).

3.2 Contraindications

Myfortic is contraindicated in patients with hypersensitivity to mycophenolate sodium, mycophenolic acid or mycophenolate mofetil or to any of the excipients (see section 2 Description and composition).

3.3 Special warnings and precautions for use

Patients with rare hereditary deficiency of hypoxanthine-guanine phosphoribosyl-transferase (HGPRT)

Myfortic is an IMPDH (inosine monophosphate dehydrogenase) inhibitor. On theoretical grounds, it should therefore be avoided in patients with a rare hereditary deficiency of hypoxanthine-guanine phosphoribosyl-transferase (HGPRT) such as Lesch-Nyhan and Kelley-Seegmiller syndrome.

Pregnancy, lactation, females and males of reproductive potential

Use of Myfortic during pregnancy is associated with an increased risk of pregnancy loss, including spontaneous abortion and/or congenital malformations. Myfortic therapy should not be initiated in females of reproductive potential until a negative pregnancy test has been obtained. For information on use in pregnancy and contraceptive requirements (see section 9 Pregnancy, lactation, females and males of reproductive potential).

Myfortic should not be used during breast-feeding (see section 9 Pregnancy, lactation, females and males of reproductive potential).

Malignancies

Patients receiving immunosuppressive regimens involving combinations of drugs, including Myfortic, are at increased risk

of developing lymphomas and other malignancies, particularly of the skin (see section 7 Adverse drug reactions). The risk appears to be related to the intensity and duration of immunosuppression rather than to the use of any specific agent. As general advice to minimize the risk of skin cancer, exposure to sunlight and UV light should be limited by wearing protective clothing and using a high protection factor sunscreen.

Infections

Patients receiving Myfortic should be instructed to immediately report any evidence of infection, unexpected bruising, bleeding or any other manifestation of bone marrow depression.

Oversuppression of the immune system increases susceptibility to infection including opportunistic infections, fatal infections and sepsis (see section 7 Adverse drug reactions).

Reactivation of hepatitis B (HBV) or hepatitis C (HCV) have been reported in patients treated with immunosuppressants, including the mycophenolic acid (MPA) derivatives Myfortic and MMF. Monitoring infected patients for clinical and laboratory signs of active HBV or HCV infection is recommended.

Cases of progressive multifocal leukoencephalopathy (PML), sometimes fatal, have been reported in patients treated with MPA derivatives which include mycophenolate mofetil and mycophenolate sodium (see section 7 Adverse drug reactions). The reported cases generally had risk factors for PML, including immunosuppressant therapies and impairment of immune functions. In immunosuppressed patients, physicians should consider PML in the differential diagnosis in patients reporting neurological symptoms and consultation with a neurologist should be considered as clinically indicated. Polyomavirus associated nephropathy (PVAN), especially due to BK virus infection, should be included in the differential diagnosis in immunosuppressed patients with deteriorating renal function (see section 7 Adverse drug reactions). Consideration should be given to reducing the total immunosuppression in patients who develop PML or PVAN. In transplant patients, however, reduced immunosuppression may place the graft at risk.

Blood dyscrasias

Patients receiving Myfortic should be monitored for blood dyscrasias (e.g. neutropenia or anemia - see section 7 Adverse drug reactions), which may be related to MPA itself, comedication, viral infections, or some combination of these causes. Patients taking Myfortic should have complete blood cell counts weekly during the first month, twice monthly for the second and third months of treatment, then monthly throughout the first year. If blood dyscrasias occur (e.g. neutropenia with absolute neutrophil count $<1.5 \times 10^3$ / micro L or anemia) it may be appropriate to interrupt or discontinue Myfortic.

Cases of pure red cell aplasia (PRCA) have been reported in patients treated with MPA derivatives in combination with other immunosuppressants (see section 7 Adverse drug reactions). The

mechanism for MPA derivatives induced PRCA is unknown; the relative contribution of other immunosuppressants and their combinations in an immunosuppressive regimen is also unknown. However, MPA derivatives may cause blood dyscrasias (see above). In some cases PRCA was found to be reversible with dose reduction or cessation of therapy with MPA derivatives. In transplant patients, however, reduced immunosuppression may place the graft at risk. Changes to Myfortic therapy should only be undertaken under appropriate supervision in transplant recipients in order to minimize the risk of graft rejection.

Vaccinations

Patients should be advised that vaccinations may be less effective during treatment with MPA and the use of the live attenuated vaccines should be avoided (see section 8 Interactions).

Influenza vaccination may be of value. Prescribers should refer to national guidelines for influenza vaccination.

Gastrointestinal disorders

As MPA derivatives have been associated with an increased incidence of digestive system adverse events, including infrequent cases of gastrointestinal tract ulceration, hemorrhage and perforation, Myfortic should be administered with caution in patients with active serious digestive system disease.

Combination with other agents

Myfortic has been administered in combination with the following agents in clinical trials: antithymocyte globulin, basiliximab, ciclosporin for microemulsion and corticosteroids. The efficacy and safety of the use of Myfortic with other immunosuppressants have not been studied.

3.4 Interactions with other pharmaceutical products and other forms of interactions

Observed interactions resulting in a concomitant use not recommended

Azathioprine: It is recommended that Myfortic should not be co-administered with azathioprine because such co-administration has not been studied (see section 6 Warnings and precautions).

Live vaccines: Live vaccines should not be given to patients with an impaired immune response. The antibody response to other vaccines may be diminished (see section 6 Warnings and precautions).

Observed interactions to be considered

Aciclovir: Higher plasma concentrations of both MPAG (mycophenolic acid glucuronide) and aciclovir may occur in the presence of renal impairment. Therefore, the potential exists for these two drugs to compete for tubular secretion, resulting in a further increase in the concentration of both MPAG and aciclovir. In this situation patients should be carefully monitored.

Gastroprotective agents

Antacids with magnesium and aluminium hydroxides

The absorption of mycophenolate sodium was decreased when administered with antacids. Co-administration of Myfortic and antacids containing magnesium and aluminium hydroxide results in a 37% decrease in MPA systemic exposure and a 25% decrease in MPA maximal concentration. Caution should be used when co-administering antacids (containing magnesium and aluminium hydroxide) with Myfortic.

Proton pump inhibitors

In healthy volunteers, co-administration of 1000 mg MMF and 40 mg pantoprazole twice daily led to a 27% decrease in MPA AUC and to a 57% decrease in MPA C_{max}. However, in the same study, no changes in the pharmacokinetics of MPA were observed following co-administration of Myfortic and pantoprazole.

Ganciclovir: MPA and MPAG pharmacokinetics are unaffected by the addition of ganciclovir. The clearance of ganciclovir is unchanged in the setting of therapeutic MPA exposure. However, in patients with renal impairment in which Myfortic and ganciclovir are coadministered the dose recommendations for ganciclovir should be observed and patients monitored carefully.

Tacrolimus: In a calcineurin cross-over study in stable renal transplant patients, steady state Myfortic pharmacokinetics were measured during both Neoral[®] and tacrolimus treatments. Mean MPA AUC was 19% higher and C_{max} about 20% lower. Conversely mean MPAG AUC and C_{max} were about 30% lower on tacrolimus treatment compared to Neoral[®] treatment.

Ciclosporin A: When studied in stable renal transplant patients, ciclosporin A pharmacokinetics were unaffected by steady state dosing of Myfortic.

Anticipated interactions to be considered

Cholestyramine and drugs that interfere with enterohepatic circulation: Due to its capacity to block the enteric circulation of drugs, cholestyramine may decrease the systemic exposure of MPA. Caution should be used when co-administering cholestyramine or drugs that interfere with enterohepatic circulation due to the potential to reduce the efficacy of Myfortic.

Oral contraceptives: Oral contraceptives undergo oxidative metabolism while Myfortic is metabolized by glucuronidation. A clinically significant effect of oral contraceptives on Myfortic pharmacokinetics is not anticipated. However, as the long term effect of Myfortic dosing on the pharmacokinetics of oral contraceptives is not known, it is possible that the efficacy of oral contraceptives may be adversely affected (see section 9 Pregnancy, lactation, females and males of reproductive potential).

3.5 Pregnancy, lactation, females and males of reproductive potential

Pregnancy

Risk Summary

Use of Myfortic during pregnancy is associated with an increased risk of spontaneous abortion and congenital malformations. Although there are no adequate and well controlled studies in pregnant women conducted with Myfortic, based on data from the US National Transplant Pregnancy Registry (NTPR), use of mycophenolate mofetil in combination with other immunosuppressants during pregnancy was associated with an increased rate of 22 % (four cases in 18 liveborn with exposure) of congenital malformations, compared to the rate of 4 to 5% for malformations seen among transplant patients in the NTPR. Congenital malformations that have been reported with mycophenolate mofetil include outer ear and other facial abnormalities including cleft lip and palate, congenital diaphragmatic hernia, anomalies of the distal limbs, heart, esophagus and kidney. Use of mycophenolate mofetil during pregnancy was also reported to be associated with increased risk of spontaneous abortion. Since MMF is converted to MPA following oral or IV administration, the above risks must be taken into account for Myfortic as well. The teratogenic potential of MPA was observed in animal studies (see “Animal data”).

Myfortic should be used in pregnant women only if the potential benefit outweighs the potential risk to the fetus. Patients should be instructed to consult their physician immediately should pregnancy occur.

Animal data

In a teratology study in rats, administration of mycophenolate sodium during organogenesis resulted in malformations including anophthalmia, exencephaly and umbilical hernia, at an oral dose as low as 1 mg/kg/day. The systemic exposure at this dose represents 0.05 times the clinical exposure at the MRHD (maximum recommended human dose) of 1440 mg/day Myfortic.

In a pre- and postnatal development study in rats oral administration of mycophenolic acid (as sodium salt) during gestation and lactation caused developmental delays (abnormal pupillary reflex in females and preputial separation in males) at the highest dose of 3 mg/kg, which is below MRHD based on body surface area.

Lactation

Risk Summary

It is not known whether MPA is transferred into human milk. There are no data on the effects of Myfortic on the breastfed child or on milk production. As many drugs are transferred into human milk, and due to the potential for serious adverse reactions in breast-fed newborns/infants, a decision should be made whether to abstain from breast-feeding during treatment and for 6 weeks after stopping the therapy or to abstain from using the medicinal product, taking into account the importance of the drug to the mother (see section 6 Warnings and precautions).

Females and males of reproductive potential

Pregnancy testing

Myfortic therapy should not be initiated until a negative pregnancy test has been obtained.

Contraception

Females

Females of reproductive potential must use effective contraception (methods that result in less than 1 % pregnancy rates) before beginning Myfortic therapy, during therapy and for six weeks after their last Myfortic dose (see section 8 Interactions).

Males

Male patients are recommended to use condoms during treatment, and for a total of 13 weeks after their last Myfortic dose. Accordingly, male patients of reproductive potential should be made aware of and discuss with a qualified health-care professional the potential risks of fathering a child or donating semen. In addition, female partners of the male patients are recommended to use effective contraception (methods that result in less than 1 % pregnancy rates) during treatment and for a total of 13 weeks after the last Myfortic dose.

Infertility

There is no data on the effect of Myfortic on human fertility. Mycophenolate sodium had no effect on male and female rat's fertility at oral doses up to 40 mg/kg/day and 20 mg/kg/day respectively, equivalent to 9 and 4.5 (calculated) times the clinical exposure at the MRHD of 1440 mg Myfortic per day (see section 13 Non-clinical safety data).

3.6 Effects on ability to drive and use machines

Not applicable

3.7 Undesirable effects

Summary of the safety

profile

The following undesirable effects cover adverse drug reactions from two controlled clinical trials. The trials evaluated the safety of Myfortic and mycophenolate mofetil in 423 de novo and in 322 maintenance renal transplant patients (randomized 1:1); the incidence of adverse events was similar between treatments in each population.

The very common ($\geq 10\%$) adverse drug reactions associated with Myfortic in combination with ciclosporin for microemulsion and corticosteroids include leukopenia and diarrhoea.

Malignancies

Patient receiving immunosuppressive regimens involving combinations of drugs, including MPA, are at increased risk of developing lymphomas and other malignancies, particularly of the skin

(see section 6 Warnings and precautions). Overall rates of malignancies observed in Myfortic clinical trials are as follows: lymphoproliferative disease or lymphoma developed in 2 de novo patients (0.9%) and in 2 maintenance patients (1.3%) receiving Myfortic for up to 1 year; non-melanoma skin carcinomas occurred in 0.9% of de novo and 1.8% of maintenance patients receiving Myfortic for up to 1 year; other types of malignancy occurred in 0.5% of de novo and 0.6% of maintenance patients.

Opportunistic infections

All transplant patients are at increased risk of opportunistic infections; the risk increased with total immunosuppressive load (see section 6 Warnings and precautions). The most common opportunistic infections in de novo renal transplant patients receiving Myfortic with other immunosuppressants in controlled clinical trials of renal transplant patients followed for 1 year were CMV (cytomegalovirus), candidiasis and herpes simplex. The overall rate of CMV infections (serology, viremia or disease) observed in Myfortic clinical trials was reported in 21.6% of de novo and in 1.9% of maintenance renal transplant patients.

Tabulated summary of adverse drug reactions from clinical trials

Adverse drug reactions (Table 4-1) are ranked by frequency, with the most frequent first, using the following convention: very common ($\geq 1/10$); common ($\geq 1/100$, $< 1/10$); uncommon ($\geq 1/1,000$, $< 1/100$); rare ($\geq 1/10,000$, $< 1/1,000$) very rare ($< 1/10,000$), including isolated reports. Within each frequency grouping, adverse reactions are ranked in order of decreasing seriousness.

Table 7-1 below contains adverse drug reactions possibly or probably related to Myfortic reported in the two phase III randomized, double blind, controlled, multi-center trials: 1 in de novo kidney transplant patients and 1 in maintenance kidney transplant patients, in which Myfortic was administered at a dose of 1,440 mg /day for 12 months together with ciclosporin microemulsion and corticosteroids. It is compiled according to MedDRA system organ class.

Table 4-1 Adverse drug reactions possibly or probably related to Myfortic reported in the two phase III pivotal trials

Very common	Viral, bacterial and fungal infections
Common	Upper respiratory tract infections, pneumonia
Uncommon	Wound infection, sepsis*, osteomyelitis*
Neoplasms benign and malignant	
Uncommon	Skin papilloma*, basal cell carcinoma*, Kaposi's sarcoma*, Lymphoproliferative disorder, squamous cell carcinoma*
Blood and lymphatic system disorders	
Very common	Leukopenia
Common	Anaemia, thrombocytopenia

Uncommon	Lymphocele*, lymphopenia*, neutropenia*, lymphadenopathy*
Metabolism and nutrition disorders	
Very common	Hypocalcaemia, hypokalaemia, hyperuricaemia
Common	Hyperkalaemia, hypomagnesaemia
Uncommon	Anorexia, hyperlipidaemia, diabetes mellitus*, hypercholesterolaemia*, hypophosphataemia
Psychiatric disorders	
Common	Anxiety
Uncommon	Delusional perception*
Nervous system disorders	
Common	Dizziness, headache
Uncommon	Tremor, insomnia*
Eye disorders	
Uncommon	Conjunctivitis*, blurred vision *
Cardiac disorders	
Uncommon	Tachycardia, pulmonary oedema*
Vascular disorders	
Very common	Hypertension
Common	Aggravated hypertension, hypotension
Respiratory, thoracic and mediastinal disorders	
Common	Cough, dyspnoea, dyspnoea exertional
Uncommon	Interstitial lung disease including fatal pulmonary fibrosis, pulmonary congestion*, wheezing*
Gastrointestinal disorders	
Very common	Diarrhoea
Common	Abdominal distension, abdominal pain, constipation, dyspepsia, flatulence, gastritis, loose stools, nausea, vomiting
Uncommon	Abdominal tenderness, pancreatitis, eructation, halitosis*, ileus*, oesophagitis*, peptic ulcer*, subileus*, gastrointestinal haemorrhage, dry mouth*, lip ulceration*, parotid duct obstruction*, gastro-oesophageal reflux disease*, gingival hyperplasia*, peritonitis*
Hepatobiliary disorders	
Common	Abnormal hepatic function tests
Skin and subcutaneous tissue disorders	
Uncommon	Alopecia, contusion*, acne
Musculoskeletal, connective tissue disorders	
Common	Arthralgia, asthenia, myalgia
Uncommon	Back pain*, muscle cramps
Renal and urinary disorders	
Common	Increased blood creatinine

Uncommon	Haematuria*, renal tubular necrosis*, urethral stricture
General disorders and administration site conditions	
Common	Fatigue, peripheral oedema, pyrexia
Uncommon	Influenza like illness, lower limb oedema *, pain, rigors*, weakness*

* Event reported in a single patient (out of 372) only.

Note: Renal transplant patients were treated with 1,440 mg Myfortic daily for up to one year. A similar profile was seen in the de novo and maintenance transplant population although the incidence tended to be lower in the maintenance patients.

Listing of adverse drug reactions from post-marketing experience

The following adverse drug reactions have been derived from post-marketing experience with Myfortic via spontaneous case reports and literature cases. As these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency which is therefore categorized as not known. Adverse drug reactions are listed according to MedDRA system organ class. Within each system organ class, ADRs are presented in order of decreasing seriousness.

Immune system disorders: Hypersensitivity reactions (including anaphylaxis).

Skin and subcutaneous tissue disorders: Rash has been identified as an adverse drug reaction from post-approval clinical trials, post marketing surveillance and spontaneous reports.

General disorders and administration site conditions: de novo purine synthesis inhibitors-associated acute inflammatory syndrome

The following adverse reactions are attributed to MPA derivatives as a class effect:

Infections and infestations: Serious, sometimes life-threatening infections, including meningitis, infectious endocarditis, tuberculosis, and atypical mycobacterial infection. Polyomavirus associated nephropathy (PVAN), especially due to BK virus infection. Cases of progressive multifocal leukoencephalopathy (PML), sometimes fatal, have been reported (see section 6 Warnings and precautions).

Blood and lymphatic system disorders: Agranulocytosis, neutropenia, pancytopenia. Cases of pure red cell aplasia (PRCA) have been reported in patients treated with MPA derivatives in combination with other immunosuppressants (see section 6 Warnings and precautions).

Gastrointestinal disorders: Colitis, oesophagitis (including CMV-colitis and - oesophagitis), CMV gastritis, pancreatitis, intestinal perforation, gastrointestinal haemorrhage, gastric ulcers, duodenal ulcers, ileus.

Geriatric population (65 years of age or older)

Geriatric patients may generally be at increased risk of adverse drug reactions due to immunosuppression. Geriatric patients receiving Myfortic as part of a combination immunosuppressive regimen, did not show an increased risk of adverse reactions, compared to younger individuals in the Myfortic clinical trials.

Overdosage
There have been anecdotal reports of deliberate or accidental overdoses with Myfortic, whereas not all patients experienced related adverse events.

In overdose cases in which adverse events were reported, the events fall within the known safety profile of the class. Accordingly an overdose of Myfortic could possibly result in oversuppression of the immune system and may increase the susceptibility to infection including opportunistic infections, fatal infections and sepsis. If blood dyscrasias occur (e.g. neutropenia with absolute neutrophil count $<1.5 \times 10^3$ / micro L or anemia) it may be appropriate to interrupt or discontinue Myfortic (see sections 6 Warnings and precautions and 7 Adverse drug reactions).

Although dialysis may be used to remove the inactive metabolite MPAG, it would not be expected to remove clinically significant amounts of the active moiety MPA. This is in large part due to the very high plasma protein binding of MPA, 97%. By interfering with the enterohepatic circulation of MPA, bile acid sequestrants, such as cholestyramine, may reduce systemic MPA exposure.

5 Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic properties

ATC

Pharmacotherapeutic group: immunosuppressant (ATC code L04AA06).

Mechanism of action (MOA)

MPA inhibits the proliferation of T- and B lymphocytes more potently than other cells because in contrast to other cell types that can utilize purine salvage pathways the lymphocyte proliferation is critically dependent on de novo synthesis. Thus, the mode of action is complementary to calcineurin inhibitors, which interfere with cytokine transcription and resting T-lymphocytes.

Pharmacodynamics (PD)

Not applicable.

5.2 Pharmacokinetic properties

Absorption

Following oral administration, mycophenolate sodium is extensively absorbed. Consistent with its enteric coated design, the time to

maximal MPA concentration was approximately 1.5 to 2 hours. *In vitro* studies demonstrated that the enteric coated Myfortic formulation prevents the release of MPA under acidic conditions as in the stomach.

In stable renal transplant patients on ciclosporin for microemulsion based immunosuppression, the gastrointestinal absorption of MPA was 93% and absolute bioavailability was 72%. Myfortic pharmacokinetics are dose proportional and linear over the studied dose range of 180 to 2,160 mg. Compared to the fasting state, administration of 720 mg Myfortic with a high fat meal (55 g fat, 1,000 calories) had no effect on the systemic exposure of MPA (AUC) which is the most relevant PK parameter linked to efficacy. However there was a 33% decrease in the maximal concentration of MPA (C_{max}).

Distribution

The volume of distribution of MPA at steady state is 50 liters. Both mycophenolic acid and mycophenolic acid glucuronide are highly protein bound, 97% and 82%, respectively. The free MPA concentration may increase under conditions of decreased protein binding sites (uremia, hepatic failure, hypoalbuminemia, concomitant use of drugs with high protein binding). This may put patients at increased risk of MPA-related adverse effects.

Biotransformation/metabolism

The half-life of MPA is 11.7 hours and the clearance is 8.6 L/hr. MPA is metabolized principally by glucuronyl transferase to form the phenolic glucuronide of MPA, mycophenolic acid glucuronide (MPAG). MPAG is the predominant metabolite of MPA and does not manifest biologic activity. In stable renal transplant patients on ciclosporin for microemulsion based immunosuppression, approximately 28% of the oral Myfortic dose is converted to MPAG by presystemic metabolism. The half-life of MPAG is longer than that of MPA, approximately 15.7 hours and its clearance is 0.45 L/hr.

Elimination

Although negligible amounts of MPA are present in the urine (<1.0%), the majority of MPA is eliminated in the urine as MPAG. MPAG secreted in the bile is available for deconjugation by gut flora. The MPA resulting from this deconjugation may then be reabsorbed. Approximately 6 to 8 hours after Myfortic dosing a second peak of MPA concentration can be measured, consistent with reabsorption of the deconjugated MPA.

Pharmacokinetics in renal transplant patients on ciclosporin for microemulsion based immunosuppression

Table 5-1 below shows mean pharmacokinetic parameters for MPA following Myfortic administration. Single dose Myfortic pharmacokinetics predicts multiple dose and chronic dosing Myfortic pharmacokinetics. In the early post-transplant period, mean MPA

AUC and mean MPA C_{max} was approximately one-half of that measured six months post-transplant.

Table 5-1 Mean (SD) pharmacokinetic parameters for MPA following oral administration of Myfortic to renal transplant patients on Ciclosporin for microemulsion based Immunosuppression

Adult single dose n = 24	Dose (oral)	T _{max} (hrs)	C _{max} (microgram /mL)	AUC 0- (microgram*hr /mL)
	720 mg	2	26.1 (12.0)	66.5 (22.6)
Adult Multiple dose x 6 days BID n=12	Dose (oral)	T _{max} (hrs)	C _{max} (microgram /mL)	AUC 0-12 (microgram*hr/ mL)
	720 mg	2	37.0 (13.3)	67.9 (20.3)
Adult Multiple dose x 28 days BID n = 36	Dose (oral)	T _{max} (hrs)	C _{max} (microgram /mL)	AUC 0-12 (microgram*hr/ mL)
	720 mg	2.5	31.2(18.1)	71.2(26.3)
Adult Chronic, multiple dosing BID (Study ERLB 301) n=48	Dose	T _{max} (hrs)	C _{max} (microgram /mL)	AUC 0-12 (microgram*hr/ mL)
14 days post-transplant	720 mg	2	13.9 (8.6)	29.1 (10.4)
3 months post-transplant	720 mg	2	24.6 (13.2)	50.7 (17.3)
6 months post-transplant	720 mg	2	23.0 (10.1)	55.7 (14.6)
Pediatric single dose n=10	Dose	T _{max} (hrs)	C _{max} (microgram /mL)	AUC 0- (microgram*hr/ mL)
	450 mg/m ²	2-2.5	31.9 (18.2)	76.2 (25.2)

Special populations

Geriatric population (65 years of age or above)

Based on preliminary data MPA exposure does not appear to vary to a clinically significant degree by age.

Pediatric population (below 18 years)

Safety and efficacy in children have not been established. Limited pharmacokinetics data are available on the use of Myfortic in children. In the table above mean (SD) MPA pharmacokinetics are

shown for stable pediatric renal transplant patients on ciclosporin microemulsion based immunosuppression. Increased variability of MPA C_{max} and AUC were noted in these pediatric patients compared to adult renal transplant patients. Mean MPA AUC at this dose was higher than typically measured in adults receiving 720 mg Myfortic. The mean apparent MPA clearance was approximately 7.7 L/hr. A Myfortic dose of 200 to 300 mg/m² would be expected to result in a MPA AUC of 30 to 50 micrograms hr/mL.

Gender

There are no clinically significant gender differences in Myfortic pharmacokinetics.

Race/ethnicity

Following a single dose administration of 720 mg Myfortic to 18 healthy Japanese and Caucasian subjects, the exposure (AUC_{inf}) for MPA and MPAG were 15 and 22% lower in Japanese subjects compared to Caucasians. The peak MPAG concentrations (C_{max}) were similar between the two populations, however, Japanese subjects had 9.6% higher C_{max} for MPA. These results do not suggest any clinically relevant differences.

Renal impairment

MPA pharmacokinetic appeared to be unchanged over the range of normal to absent renal function. In contrast, MPAG exposure increased with decreased renal function; MPAG exposure being approximately 8 fold higher in the setting of anuria. Clearance of either MPA or MPAG was unaffected by hemodialysis. Free MPA may also significantly increase in the setting of renal failure. This may be due to decreased MPA plasma protein binding in the presence of high blood urea concentration.

Hepatic impairment

In volunteers with alcoholic cirrhosis, hepatic MPA glucuronidation processes were relatively unaffected by hepatic parenchymal disease. Effects of hepatic disease on this process probably depend on the particular disease. However, hepatic disease with predominantly biliary damage, such as primary biliary cirrhosis, may show a different effect.

5.2 Pre-clinical safety data

Safety pharmacology and repeat dose toxicity

The hematopoietic and lymphoid system were the primary organs affected in toxicology studies conducted with mycophenolate sodium in rats and mice. Aplastic, regenerative anemia was identified as the dose-limiting toxicity in MPA-exposed rodents. Evaluation of myelograms showed a marked decrease in erythroid cells (polychromatic erythroblasts and normoblasts) and a dose-dependent spleen enlargement and increase in extramedullary hematopoiesis.

These effects occurred at systemic exposure levels which are equivalent to or less than the clinical exposure at the recommended dose of 1440 mg/day Myfortic in renal transplant patients.

The non-clinical toxicity profile of mycophenolate sodium appears to be consistent with adverse events observed in MPA-exposed humans, which now provide safety data of more relevance to the patient population (see section 7 Adverse drug reactions).

Single oral doses of MPA are moderately well tolerated in rats (LD50 of 350 to 700 mg/kg), well tolerated in mice or monkeys (LD50 of more than 1,000 mg/kg), and extremely well tolerated in rabbits (LD50 of more than 6,000 mg/kg).

Reproductive toxicity

For information on reproductive toxicity, see section 9 Pregnancy, lactation, females and males of reproductive potential.

Carcinogenicity and mutagenicity

In a 104-week oral carcinogenicity study in rats, mycophenolate sodium at daily doses up to 9 mg/kg was not tumorigenic. The highest dose tested resulted in approximately 0.6 to 1.2 times the systemic exposure observed in renal transplant patients at the recommended dose of 1440 mg/day. Similar results were observed in a parallel study in rats performed with mycophenolate mofetil. In a 26-week oral carcinogenicity assay in a P53[±] (heterozygous) transgenic mouse model, mycophenolate sodium at daily doses up to 200 mg/kg was not tumorigenic. The highest dose tested was 200 mg/kg, resulting in approximately 5 times the systemic exposure observed in renal transplant patients (1440 mg/day).

The genotoxic potential of mycophenolate sodium was determined in five assays. MPA was genotoxic in the mouse lymphoma/thymidine kinase assay, the micronucleus test in V79 Chinese hamster cells and the *in vivo* mouse micronucleus assay. Mycophenolate sodium was not genotoxic in the bacterial mutation assay or the chromosomal aberration assay in human lymphocytes. The lowest dose showing genotoxic effects in a mouse bone marrow micronucleus resulted in approximately 3 times the systemic exposure (AUC or C_{max}) observed in renal transplant patients at the tested clinical dose of 1440 mg of Myfortic per day. It is probable that the mutagenic activity observed was due to a shift in the relative abundance of the nucleotides in the cellular pool used for DNA synthesis.

Fertility

Mycophenolate sodium had no effect on male rats fertility at oral doses up to 40 mg/kg/day. The systemic exposure at this dose represents approximately 9 times the clinical exposure at the tested clinical MRHD of 1440 mg Myfortic per day. No effects on female fertility were seen up to a dose of 20 mg/kg/day, a dose at which maternal toxicity and embryotoxicity were already observed.

6.2 List of excipients

The gastro resistant tablet coating of 180 mg Myfortic consists of hypromellose phthalate/ hydroxypropylmethylcellulose phthalate; titanium dioxide; iron oxide yellow; indigotin.

The gastro resistant tablet coating of 360 mg Myfortic consists of hypromellose phthalate/ hydroxypropylmethylcellulose phthalate; titanium dioxide; iron oxide yellow; iron oxide red.

6.3 Incompatibilities

Not applicable

6.4 Shelf life

36 months

6.5 Special precautions for storage

Do not store above 30 °C. Myfortic 180 mg and 360 mg gastro-resistant tablets should be protected from moisture.

Store in the original package and container.

Myfortic must be kept out of the reach and sight of children.

6.6 Nature and contents of container

50, 100, 120 or 250 film coated tablets in double-sided aluminium (PA/AL/PVC) blister foil. Not all pack sizes may be marketed.

6.7 Special precautions for disposal and other handling

Myfortic tablets should not be crushed in order to remain the integrity of the enteric coating

Any unused product or waste material should be disposed of in accordance with local requirements.

7 Marketing authorization holder

NVS Kenya Limited

7 Market authorization Number

MYFORTIC 180MG TABLET:

15957

MYFORTIC 360MG TABLET

:16124

8 Date of first registration/ renewal of the registration

26/02/2026

9 Date of revision of the text

26/02/2026