

Summary of Product Characteristics for Pharmaceutical Products

1. Name of the medicinal product

NIDAMET 400 MG TABLET

2. Qualitative and quantitative composition

Each tablet contains 400mg of Metronidazole.

Excipients with potential clinical effect:

Each tablet contains Lactose monohydrate

For a full list of excipients, see section 6.1

3. Pharmaceutical form

Yellow, circular, flat, beveled edged tablets, scored on one side and plain on the reverse side.

4. Clinical particulars

4.1 Therapeutic indications

Metronidazole is indicated in the prophylaxis and treatment of infections in which anaerobic bacteria have been identified or are suspected to be the cause.

Metronidazole is active against a wide range of pathogenic micro-organisms notably species of Bacteroides, Fusobacteria, Clostridia, Eubacteria, anaerobic cocci and Gardnerella vaginalis.

It is also active against Trichomonas, Entamoeba histolytica, Giardia lamblia and Balantidium coli.

Metronidazole is indicated in adults and children for the following indications:

The prevention of post-operative infections due to anaerobic bacteria, particularly species of Bacteroides and anaerobic streptococci.

The treatment of septicaemia, bacteraemia, peritonitis, brain abscess, necrotising pneumonia, osteomyelitis, puerperal sepsis, pelvic abscess, pelvic cellulitis, and post-operative wound infections from which pathogenic anaerobes have been isolated.

- Urogenital trichomoniasis in the female (trichomonal vaginitis) and in the male.
- Bacterial vaginosis (also known as non-specific vaginitis, anaerobic vaginosis or Gardnerella vaginitis).
- All forms of amoebiasis (intestinal and extra-intestinal disease and that of symptomless cyst passers).
- Giardiasis.
- Acute ulcerative gingivitis.
- Anaerobically-infected leg ulcers and pressure sores.
- Acute dental infections (e.g. acute pericoronitis and acute apical infections). Considerations should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Oral route of administration. Metronidazole tablets should be swallowed with water (not chewed). It is recommended that the tablets be taken during or after a meal.

Prophylaxis against anaerobic infection: Chiefly in the context of abdominal (especially colorectal) and gynaecological surgery.

Adults

400 mg 8 hourly during 24 hours immediately preceding operation followed by postoperative intravenous or rectal administration until the patient is able to take tablets.

Children

Children < 12 years: 20-30mg/kg as a single dose given 1-2 hours before surgery Newborns with a gestation age < 40 weeks: 10mg/kg body weight as a single dose before operation

Anaerobic infections: The duration of a course of Metronidazole treatment is about 7 days but it will depend upon the seriousness of the patient's condition as assessed clinically and bacteriologically.

Treatment of established anaerobic infection:

Adults

800 mg followed by 400 mg 8 hourly.

Children

Children > 8 weeks to 12 years of age: The usual daily dose is 20-30mg/kg/day as a single dose or divided into 7.5mg/kg every 8 hours.

The daily dose may be increased to 40mg/kg, depending on the severity of the infection. Duration of treatment is usually 7 days.

Children < 8 weeks of age: 15mg/kg as a single dose daily or divided into 7.5mg/kg every 12 hours. In newborns with a gestation age < 40 weeks, accumulation of metronidazole can occur during the first week of life, therefore the concentrations of metronidazole in serum should preferable be monitored after a few days therapy.

Protozoal and other infections:

<i>Dosage is given in terms of metronidazole or metronidazole equivalent</i>					
	Duration of dosage in days	Adults and children over 10years	Children		
			7 to 10 years	3 to 7 years	1 to 3 years
<i>Urogenital trichomoniasis</i> Where re-infection is likely, in adults the consort should receive a similar course of treatment concurrently	7 Or	2000mg as a single dose or 200 mg three times daily	40mg/kg orally as a single dose or 15-30 mg/kg/day divided in 2-3 doses; not to exceed 2000mg/dose		
		Or			
	5- 7	400mg twice daily			
<i>Bacterial vaginosis</i>	5-7 or	400 mg twice daily			
	1	2000mg as a single dose			

Amoebiasis (a) Invasive intestinal disease in susceptiblesubjects	5	800 mg threetimes daily	400 mg threetimes daily	200 mg fourtimes daily	200 mg threetimes daily
(b) Intestinal disease inless susceptible subjects and chronic amoebic hepatitis	5-10	400 mg threetimes daily	200 mg threetimes daily	100 mg fourtimes daily	100 mg threetimes daily
(c) Amoebic liver abscess also other forms of extra-intestinalamoebiasis	5	400 mg threetimes daily	200 mg threetimes daily	100 mg fourtimes daily	100 mg threetimes daily
(d) Symptomless cystpassers	5-10	400-800 mg threetimes daily	200-400 mg threetimes daily	100-200 mg fourtimes daily	100-200 mg three times daily
	Alternatively, doses may be expressed by body weight 35 to 50mg/kg daily in 3 divided doses for 5 to 10 days, not to exceed 2400mg/day				
Giardiasis	3	2000mg oncedaily Or	1000mg once daily	600-800 mg oncedaily	500 mg oncedaily
	5	400mg threetimes daily or			
	7-10	500mg twice daily			
	Alternatively, as expressed in mg per kg of body weight: 15-40mg/kg/day divided in 2-3 doses.				
Dosage is given in terms of metronidazole or metronidazole equivalent					
	Duration of dosage in days	Adults and children over 10years	Children		
			7 to 10 years	3 to 7 years	1 to 3 years
Acute ulcerative gingivitis	3	200 mg threetimes daily	100 mg threetimes daily	100 mg twicedaily	50 mg threetimes daily
Acute dental infections	3-7	200 mg threetimes daily			
Leg ulcers andpressure sores	7	400 mg threetimes daily			
Children and infants weighing less than 10 kg should receive proportionally smaller dosages. Elderly: Flagyl is well tolerated by the elderly but a pharmacokinetic study suggests cautious use of high dosageregimens in this age group.					

Eradication of *Helicobacter pylori* in paediatric patients

As a part of a combination therapy, 20mg/kg/day not to exceed 500mg twice daily for 7-14 days. Official guidelines should be consulted before initiating therapy.

4.3 Contraindications

Known hypersensitivity to nitroimidazoles, metronidazole or any of the excipients.

4.4 Special warnings and precautions for use

Regular clinical and laboratory monitoring (especially leucocyte count) are advised if administration of Metronidazole for more than 10 days is considered to be necessary and patients should be monitored for adverse reactions, such as peripheral or central neuropathy (such as paraesthesia, ataxia, dizziness, convulsive seizures)

Metronidazole should be used with caution in patients with active or chronic severe peripheral and central nervous system disease due to the risk of neurological aggravation.

Cases of severe hepatotoxicity/acute hepatic failure, including cases with a fatal outcome with very rapid onset after treatment initiation in patients with cockayne syndrome have been reported with products containing metronidazole for systemic use. In this population, metronidazole should therefore be used after careful benefit-risk assessment and only if no alternative treatment is available. Liver function tests must be performed just prior to the start of therapy, throughout and after end of treatment until liver function is within normal ranges, or until the baseline values are reached. If the liver function tests become markedly elevated during treatment, the drug should be discontinued.

Patients with cockayne syndrome should be advised to immediately report any symptoms of potential liver injury to their physician and stop taking metronidazole.

Cases of severe bullous skin reactions such as Stevens Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) or acute generalised exanthematous pustulosis (AGEP) have been reported with metronidazole. If symptoms or signs of SJS, TEN or AGEP are present, Metronidazole treatment must be immediately discontinued.

There is a possibility that after *Trichomonas vaginalis* has been eliminated a gonococcal infection might persist.

The elimination half-life of metronidazole remains unchanged in the presence of renal failure. The dosage of metronidazole therefore needs no reduction. Such patients however retain the metabolites of metronidazole. The clinical significance of this is not known at present. In patients undergoing haemodialysis, metronidazole and metabolites are efficiently removed during an eight-hour period of dialysis. Metronidazole should therefore be re-administered immediately after haemodialysis. No routine adjustment in the dosage of Metronidazole need be made in patients with renal failure undergoing intermittent peritoneal dialysis (IDP) or continuous ambulatory peritoneal dialysis (CAPD).

Metronidazole is mainly metabolized by hepatic oxidation. Substantial impairment of metronidazole clearance may occur in the presence of advanced hepatic insufficiency. Significant cumulation may occur in patients with hepatic encephalopathy and the resulting high plasma concentrations of metronidazole may contribute to the symptoms of the encephalopathy. Metronidazole should therefore, be administered with

caution to patients with hepatic encephalopathy. The daily dosage should be reduced to one third and may be administered once daily. Patients should be warned that metronidazole may darken urine. Due to inadequate evidence on the mutagenicity risk in humans (see section 5.3), the use of Metronidazole for longer treatment than usually required should be carefully considered.

4.5 Interaction with other medicinal products and other forms of interaction

Patients should be advised not to take alcohol during metronidazole therapy and for at least 48 hours afterwards because of the possibility of a disulfiram-like (antabuse effect) reaction. Psychotic reactions have been reported in patients who were using metronidazole and disulfiram concurrently.

Some potentiation of anticoagulant therapy has been reported when metronidazole has been used with the warfarin type oral anticoagulants. Dosage of the latter may require reducing. Prothrombin times should be monitored. There is no interaction with heparin.

Lithium retention accompanied by evidence of possible renal damage has been reported in patients treated simultaneously with lithium and metronidazole. Lithium treatment should be tapered or withdrawn before administering metronidazole. Plasma concentrations of lithium, creatinine and electrolytes should be monitored in patients under treatment with lithium while they receive metronidazole.

Patients receiving phenobarbital or phenytoin, metabolise metronidazole at a much greater rate than normally, reducing the half-life to approximately 3 hours.

Metronidazole reduces the clearance of 5 fluorouracil and can therefore result in increased toxicity of 5 fluorouracil.

Patients receiving ciclosporin are at risk of elevated ciclosporin serum levels. Serum ciclosporin and serum creatinine should be closely monitored when coadministration is necessary.

Plasma levels of busulfan may be increased by metronidazole which may lead to severe busulfan toxicity.

4.6 Pregnancy and lactation

There is inadequate evidence of the safety of metronidazole in pregnancy but it has been in wide use for many years without apparent ill consequence. Nevertheless metronidazole, like other medicines, should not be given during pregnancy or during lactation unless the physician considers it essential; in these circumstances the short, high-dosage regimens are not recommended.

4.7 Effects on ability to drive and use machines

Patients should be warned about the potential for drowsiness, dizziness, confusion, hallucinations, convulsions or transient visual disorders, and advised not to drive or operate machinery if these symptoms occur.

4.8 Undesirable effects

The frequency of adverse events listed below is defined using the following convention:

very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

Serious adverse reactions occur rarely with standard recommended regimens. Clinicians who contemplate continuous therapy for the relief of chronic conditions, for periods longer than those recommended, are advised to consider the possible therapeutic benefit against the risk of peripheral neuropathy.

Blood and lymphatic system disorders:

Very rare: agranulocytosis, neutropenia, thrombocytopenia, pancytopenia Not known: leucopenia.

Immune system disorders:

Rare: anaphylaxis

Not known: angiodema, urticaria, fever. Metabolism and nutrition disorders:

Not known: anorexia.

Psychiatric disorders:

Very rare: psychotic disorders, including confusion and hallucinations.

Not known: depressed mood

Nervous system disorders: Very rare:

- encephalopathy (eg. confusion, fever, headache, hallucinations, paralysis, light sensitivity, disturbances in sight and movement, stiff neck) and subacute cerebellar syndrome (eg. ataxia, dysathria, gait impairment, nystagmus and tremor) which may resolve on discontinuation of the drug.
- drowsiness, dizziness, convulsions, headaches Not known:
- during intensive and/or prolonged metronidazole therapy, peripheral sensory neuropathy or transient epileptiform seizures have been reported. In most cases neuropathy disappeared after treatment was stopped or when dosage was reduced.
- aseptic meningitis

Eye disorders:

Very rare: vision disorders such as diplopia and myopia, which, in most cases, is transient. Not Known: optic neuropathy/neuritis

Ear and labyrinth disorders

Not known: hearing impaired/hearing loss (including sensorineural), tinnitus Gastrointestinal disorders:

Not known: taste disorders, oral mucositis, furred tongue, nausea, vomiting, gastro-intestinal disturbances such as epigastric pain and diarrhoea.

Hepatobiliary disorders:

Very rare:

- increase in liver enzymes (AST, ALT, alkaline phosphatase), cholestatic or mixed hepatitis and hepatocellular liver injury, jaundice and pancreatitis which is reversible on drug withdrawal.
- cases of liver failure requiring liver transplant have been reported in patients treated with metronidazole in combination with other antibiotic drugs.

Skin and subcutaneous tissue disorders:

Very rare: skin rashes, pustular eruptions, acute generalised exanthematous pustulosis, pruritis, flushing

Not known: erythema multiforme, Stevens-Johnson syndrome or toxic epidermal necrolysis, fixed drug eruption

Musculoskeletal, connective tissue and bone disorders:

Very rare: myalgia, arthralgia.

Renal and urinary disorders:

Very rare: darkening of urine (due to metronidazole metabolite).

Reporting of suspected adverse reactions:

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Pharmacy and Poisons Board- Pharmacovigilance Electronic Reporting System (PvERS);

<https://pv.pharmacyboardkenya.org> .

4.9 Overdose

Single oral doses of metronidazole, up to 12g have been reported in suicide attempts and accidental overdoses. Symptoms were limited to vomiting, ataxia and slight disorientation. There is no specific antidote for metronidazole overdosage. In cases of suspected massive overdose, symptomatic and supportive treatment should be instituted.

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antibacterials for systemic use, ATC code: J01X D01

Metronidazole has antiprotozoal and antibacterial actions and is effective against *Trichomonas vaginalis* and other protozoa including *Entamoeba histolytica* and *Giardia lamblia* and against anaerobic bacteria.

5.2 Pharmacokinetic properties

Metronidazole is rapidly and almost completely absorbed on administration of Metronidazole tablets; peak plasma concentrations occur after 20 min to 3 hours.

The half-life of metronidazole is 8.5 ± 2.9 hours. Metronidazole can be used in chronic renal failure; it is rapidly removed from the plasma by dialysis. Metronidazole is excreted in milk but the intake of a suckling infant of a mother receiving normal dosage would be considerably less than the therapeutic dosage for infants.

5.3 Preclinical safety data

Metronidazole has been shown to be carcinogenic in the mouse and in the rat following chronic oral administration however similar studies in the hamster have given negative results. Epidemiological studies have provided no clear evidence of an increased carcinogenic risk in humans. Metronidazole has been shown to be mutagenic in bacteria in vitro. In studies conducted in mammalian cells in vitro as well as in rodent or humans in vivo, there was inadequate evidence of a mutagenic effect of metronidazole, with some studies reporting mutagenic effects, while other studies were negative.

6. Pharmaceutical particulars

6.1 List of excipients

Lactose Monohydrate 200 mesh

Maize starch dried

Pregelatinized starch

Potassium sorbate

Tartrazine yellow colour

Aerosil

Magnesium stearate

6.2 Incompatibilities

Not applicable

6.3 Shelf-life

3 years

6.4 Special precautions for storage

Store below 30°C in the original packaging (protect from light).

6.5 Nature and contents of container

Metronidazole tablets are available in aluminum/plastic blisters of 14 tablets and HDPE bottles of 100 tablets.

6.6 Special precautions for disposal and other handling

No special requirements

7. Marketing authorization holder

Viva Healthcare Limited

P.O. BOX 56591-00200

Nairobi.

8. Marketing authorization number(s)

H2022/CTD8956/19889

9. Date of first authorization/renewal of the authorization

2023 March 28

10. Date of revision of the text

2023 March 28