

SUMMARY OF PRODUCT CHARACTERISTICS

1. Name of the medicinal product

Sodium Chloride Intravenous Infusion BP 0.9% w/v (NS)

2. Qualitative and quantitative composition

Each ml contains 0.9g Sodium Chloride. For the full list of excipients: see section 6.1

3. Pharmaceutical form

Solution for infusion.

A clear, colourless solution free from visible particles.

4. Clinical particulars

4.1. Therapeutic indications

Sodium Chloride Solution for Infusion is indicated for:

- Treatment of isotonic extracellular dehydration
- Treatment of sodium depletion
- Vehicle or diluent or reconstitution of compatible drugs for parenteral administration.

4.2. Posology and method of administration

Posology

Adults, older people and children:

Doses may be expressed in terms of mEq or mmol of sodium, mass of sodium, or mass of sodium salt (1 g NaCl = 394 mg, 17.1 mEq or 17.1 mmol of Na and Cl).

Fluid balance, serum electrolytes and acid-base balance should be monitored before and during administration, with particular attention to serum sodium in patients with increased non-osmotic vasopressin release (syndrome of inappropriate antidiuretic hormone secretion, SIADH) and in patients comedicated with vasopressin agonist drugs, due to the risk of hospital hyponatraemia (see sections 4.4, 4.5 and 4.8). Monitoring of serum sodium is particularly important for hypotonic fluids.

Sodium Chloride 0.9% intravenous infusion has a tonicity of 308 mOsm/l (approx.)

The infusion rate and volume depend on age, weight, clinical condition (e.g. burns, surgery, head-injury, infections), and concomitant therapy should be determined by the consulting physician experienced in intravenous fluid therapy (see sections 4.4. and 4.8).

Recommended dosage

The recommended dosage for treatment of isotonic extracellular dehydration and sodium depletion is:

- For adults: 500 ml to 3 litres/24h
- For babies and children: 20 to 100 ml per 24h and per kg of body weight, depending of the age and the total body mass.

The recommended dosage when used as a vehicle or diluent ranges from 50 to 250 ml per dose of medicinal product to be administered.

When Sodium Chloride 0.9 % is used as a diluent for injectable preparations of other drugs, the dosage and the infusion rate will also be dictated by the nature and the dose regimen of the prescribed drug.

Method of administration

The solution is for administration by intravenous infusion through a sterile and non-pyrogenic administration set, using aseptic technique. The equipment should be primed with the solution in order to prevent air entering the system.

The product should be inspected visually for particulate matter and discoloration prior to administration. Do not administer unless solution is clear, free from visible particles and the seal is intact.

Do not remove unit from overwrap until ready for use. The inner bag maintains the sterility of the solution. Administer immediately following the insertion of infusion set.

Do not connect flexible plastic containers in series in order to avoid air embolism due to possible residual air contained in the primary container. Pressurizing intravenous solutions contained in flexible plastic containers to increase flow rates can result in air embolism if the residual air in the container is not fully evacuated prior to administration. Use of a vented intravenous administration set with the vent in the open position could result in air embolism. Vented intravenous administration sets with the vent in the open position should not be used with flexible plastic containers.

Additives may be introduced before infusion or during infusion through the injection site.

For information on incompatibilities and preparation of the product (with additives), please see sections 6.2 and 6.6

4.3. Contraindications

The solution is contra-indicated in patient presenting hypernatraemia or hyperchloraemia. The contra-indications related to the added medicinal product should be considered.

4.4 Special warnings and precautions for use Fluid balance/renal function

Use in patients with (severe) renal impairment

Sodium Chloride 0.9% should be administered with particular caution to patients with or at risk of severe renal impairment. In such patients, administration of Sodium Chloride 0.9% may result in sodium retention. (See “Use in patients at risk for sodium retention, fluid overload and oedema” below; for additional considerations.)

Risk of fluid and/or solute overload and electrolyte disturbances

Depending on the volume and rate of infusion, intravenous administration of Sodium Chloride 0.9% can cause:

- Fluid and/or solute overload resulting in overhydration/hypervolemia and, for example, congested states, including central and peripheral oedema.
- Clinically relevant electrolyte disturbances and acid-base imbalance.

In general, the risk of dilutional states (retention of water relative to sodium) is inversely proportional to the electrolyte concentrations of Sodium Chloride 0.9% and its additions. Conversely, the risk of solute overload causing congested states (retention of solute relative to water) is directly proportional to the electrolyte concentrations of Sodium Chloride 0.9% and its additions.

Special clinical monitoring is required at the beginning of any intravenous infusion. Clinical evaluation and periodic laboratory determinations may be necessary to monitor changes in fluid balance, electrolyte concentrations, and acid-base balance during prolonged parenteral therapy or whenever the condition of the patient or the rate of administration warrants such evaluation.

High volume infusion must be used under specific monitoring in patients with cardiac or pulmonary failure and in patients with nonosmotic vasopressin release (including SIADH), due to the risk of hospital-acquired hyponatraemia (see below).

Hyponatraemia

Patients with non-osmotic vasopressin release (e.g. in acute illness, pain, post-operative stress, infections, burns, and CNS diseases), patients with heart-, liver- and kidney diseases and patients exposed to vasopressin agonists (see section 4.5) are at particular risk of acute hyponatraemia upon infusion of hypotonic fluids.

Acute hyponatraemia can lead to acute hyponatraemic encephalopathy (cerebral oedema) characterized by headache, nausea, seizures, lethargy and vomiting. Patients with cerebral oedema are at particular risk of severe, irreversible and lifethreatening brain injury.

Children, women in the fertile age and patients with reduced cerebral compliance (e.g. meningitis, intracranial bleeding, cerebral contusion and brain oedema) are at particular risk of the severe and lifethreatening brain swelling caused by acute hyponatraemia.

Use in patients at risk for sodium retention, fluid overload and oedema

Sodium Chloride 0.9% should be used with particular caution, if at all, in patients with or at risk for:

- Hyponatraemia. Rapidly correcting hyponatraemia once adaptation has occurred may lead to cerebral oedema, potentially resulting in seizures, permanent brain damage, or death.

Hyperchloraemia

- Metabolic acidosis, which may be worsened by prolonged use of this product, especially in patients with renal impairment.

- Hypervolaemia such as congestive heart failure and pulmonary oedema may be precipitated, particularly in patients with cardiovascular disease.

- Iatrogenic hyperchloraemic metabolic acidosis (e.g., during intravenous volume resuscitation)

- Conditions that may cause sodium retention, fluid overload and oedema (central and peripheral), such as patients with primary hyperaldosteronism, secondary hyperaldosteronism, associated with, for example,

- hypertension,

- congestive heart failure,

- liver disease (including cirrhosis),

- renal disease (including renal artery stenosis, nephrosclerosis) or preeclampsia. Medications that may increase the risk of sodium and fluid retention, such as corticosteroids

Infusion reactions
Symptoms of unknown aetiology which can appear to be hypersensitivity reactions have been reported very rarely in association with infusion of Sodium Chloride 0.9 %. These have been characterized as hypotension, pyrexia, tremor, chills, urticaria, rash and pruritus. Stop the infusion immediately if signs or symptoms of these reactions develop. Appropriate therapeutic countermeasures should be instituted as clinically indicated.

Specific patient groups

The consulting physician should be experienced in this product's use and safety in these special populations that are especially sensitive to rapid changes in serum sodium levels.

Rapid correction of hyponatraemia and hypernatraemia is potentially dangerous (risk of serious neurologic complications). See section "Hyponatraemia/hypernatraemia" above Paediatric population

Plasma electrolyte concentrations should be closely monitored in the paediatric population as this population may have impaired ability to regulate fluids and electrolytes. Repeated infusions of sodium chloride should therefore only be given after determination of the serum sodium level.

Geriatric population

When selecting the type of infusion solution and the volume/rate of infusion for a geriatric patient, consider that geriatric patients are generally more likely to have cardiac, renal, hepatic, and other diseases or concomitant drug therapy.

For information on preparation of the product and additives, please see section 6.6

4.5. Interaction with Other Medicinal Products and Other Forms Of Interaction

Drugs leading to an increased vasopressin effect

The below listed drugs increase the vasopressin effect, leading to reduced renal electrolyte free water excretion and may increase the risk of hospital acquired hyponatraemia following inappropriately balanced treatment with i.v. fluids (see sections 4.2, 4.4 and 4.8).

- Drugs stimulating vasopressin release include: Chlorpropamide, clofibrate, carbamazepine, vincristine, selective serotonin reuptake inhibitors, 3,4-methylenedioxy-N-methamphetamine, ifosfamide, antipsychotics, narcotics
- Drugs potentiating vasopressin action include: Chlorpropamide, NSAIDs, cyclophosphamide
- Vasopressin analogues include: Desmopressin, oxytocin, terlipressin

Other medicinal products increasing the risk of hyponatraemia also include diuretics in general and antiepileptics such as oxcarbazepine.

Caution is advised in patients treated with lithium. Renal sodium and lithium clearance may be increased during administration of Sodium Chloride 0.9%. Administration of Sodium Chloride 0.9% may result in decreased lithium levels.

Corticoids/Steroids and carbenoxolone, are associated with the retention of sodium and water (with oedema and hypertension). See Section

4.4. Special warnings and precautions for use.

4.6. Fertility, Pregnancy and Lactation

There are no adequate data from the use of Sodium Chloride 0.9% in pregnant or lactating women. The physician should carefully consider the potential risks and benefits for each specific patient before administering Sodium Chloride 0.9%.

Sodium Chloride 0.9% should be administered with special caution for pregnant women during labour particularly as to serum-sodium if administered in combination with oxytocin (see section 4.4, 4.5 and 4.8).

Caution is advised with patients with pre-eclampsia (See Section 4.4. Special warnings and precautions for use).

When a medicinal product is added, the nature of the drug and its use during pregnancy and lactation has to be considered separately.

4.7. Effects on ability to drive and use machines

No studies have been conducted on the influence of Sodium Chloride 0.9% on the ability to operate an automobile or other heavy machinery.

4.8. Undesirable effects

Injudicious intravenous saline therapy (e.g. post-operatively and in patients with impaired cardiac or renal function) may cause hypernatraemia. Osmotically induced water shifts decrease intracellular volume, resulting in dehydration of internal organs, especially the brain, which may lead to thrombosis and haemorrhage.

General adverse effects of sodium chloride excess in the body include nausea, vomiting, diarrhoea, abdominal cramps, thirst, reduced salivary and lachrymal secretions, sweating, fever, hypotension, tachycardia, renal failure, peripheral and pulmonary oedema, respiratory arrest, headache, dizziness, restlessness, irritability, weakness, muscular twitching and rigidity, convulsions, coma and death. Excess chloride in the body may cause a loss of bicarbonate, with an acidifying effect. With judicious use of intravenous saline therapy, these side effects can be avoided.

If administered sub-cutaneously, any addition to the isotonic solution could render it hypertonic and cause pain at the site of injection.

Administration of large doses may give rise to sodium accumulation, oedema, and hyperchloraemic acidosis.

The following adverse reactions have been reported in post-marketing experience. The frequency of the adverse drug reactions listed in this section cannot be estimated from the available data.

System Organ Class (SOC)	Adverse reactions (Preferred Term)	Frequency
Nervous system disorders	Tremor Acute hyponatraemic encephalopathy*	Not known
Metabolism and nutrition disorders	Hospital acquired hyponatraemia*	Not known
Vascular disorders	Hypotension	Not known
Skin and subcutaneous tissue disorders	Urticaria Rash Pruritus	Not known
General disorders and administration site conditions:	Infusion site reactions, such as • Infusion site erythema, • Vein irritation, Injection site streaking, burning sensation, • Local pain or reaction, • Infusion site urticaria • Infection at the site of injection, • Venous thrombosis or phlebitis extending from the site of	Not known

	injection, extravasation and hypervolemia • Pyrexia • Chills	
--	--	--

- Hypernatraemia (eg. when administered to patients with nephrogenic diabetes insipidus or high nasogastric output)
- Hyperchloraemic metabolic acidosis
- Hyponatraemia, which may be symptomatic. Hyponatraemia may occur when normal free water excretion is impaired. (eg SIADH or postoperative) General adverse effects of sodium excess are described in section 4.9 Overdose.

Additives

When Sodium Chloride 0.9% is used as a diluent for injectable preparations of other drugs, the nature of additives will determine the likelihood of any other undesirable effect.

If an adverse event occurs the patient should be evaluated and appropriate counter measures be started, if needed the infusion should be stopped. The remaining part of the solution should be kept for investigation if deemed necessary.

Reporting of suspected adverse reactions

Reporting of suspected adverse reactions: Healthcare professionals are requested to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org/>

4.9. Overdose

Injudicious intravenous saline therapy (e.g. post-operatively or in patients with impaired cardiac or renal function) may cause hypernatraemia. Osmotically induced water shift decreases intracellular volume, resulting in dehydration of internal organs, especially the brain, which may lead to thrombosis and haemorrhage.

Generally, adverse effects of sodium chloride excess in the body include: nausea, vomiting, diarrhoea, abdominal cramps, thirst, reduced salivary and lachrymal secretions, sweating, fever, hypotension, tachycardia, renal failure, peripheral and pulmonary oedema, respiratory arrest, headache, dizziness, restlessness, irritability, weakness, muscular twitching and rigidity, convulsions, coma and death. Excess chloride in the body may cause a loss of bicarbonate, with an acidifying effect.

With judicious use of intravenous saline therapy, these effects can be avoided. Diuretics may be used to treat oedema resulting from isotonic expansion, and appropriate replacement therapy should be employed to avoid fluid and electrolyte imbalance.

Treatment of hypervolaemic hypernatraemia requires removal of sodium in excess of water and can be achieved by replacing diuretic-induced sodium and water losses with only water. The basic aim of therapy is to restore the volume and composition of the body fluids to normal.

Excess chloride in the body may cause a loss of bicarbonate with an acidifying effect.

When Sodium Chloride 0.9% is used as a diluent for injectable preparations of other drugs, the signs and symptoms of over infusion will be related to the nature of the additives being used. In the event of accidental over infusion, treatment should be discontinued and the patient should be observed for the appropriate signs and symptoms related to the drug administered. The relevant and supportive measures should be provided as necessary

5. Pharmacological properties

5.1. Pharmacodynamic properties

Pharmacotherapeutic Group – Electrolyte Solutions, sodium chloride

ATC Code: B05XA03

The principal determinant of the effective osmolality of the extracellular fluids (and also of the intracellular fluids, since they remain in osmotic equilibrium with the extracellular fluids) is the extracellular fluid sodium concentration. The reason for this is because sodium is the most abundant positive ion of the extracellular fluid. Negative ion concentrations of the body fluids are adjusted to equal those of the positive ions by renal acid-base control mechanisms. In addition chloride is the major anion in extracellular fluid and is involved in maintaining the acid- base balance. Solutions of sodium chloride resemble extracellular fluid. Furthermore, glucose and urea, the most abundant of the non-ionic osmolar solutes in extracellular fluids, normally only represent about 3% of the total osmolality. Therefore, in effect, the extracellular fluid sodium ion concentration controls over 90% of the effective osmotic pressure of the extracellular fluid. Sodium chloride remains the most important single salt for prophylaxis or replacement therapy of deficits of extracellular fluid. Volume contraction, whether isotonic, hypotonic or hypertonic, may seriously impair the circulation (cardiac output falls and microcirculation is compromised) and prompt infusion of isotonic sodium chloride solution is indicated. Even with moderately severe hyponatraemia or hypernatraemia, the disorder may be corrected with isotonic saline solution, provided there is normal renal function to allow physiological adjustments to be made by the kidneys, resulting in the excretion of urine at a concentration appropriate to the underlying situation.

5.2. Pharmacokinetic properties

Sodium is predominantly excreted by the kidney, but there is extensive renal reabsorption. Small amounts of sodium are lost in the faeces and sweat.

The homeostatic mechanisms involved in maintaining constant ion concentrations are well described in standard text books of physiology and biochemistry and are not, therefore, included in this report.

5.3. Preclinical safety data

The safety of sodium chloride in animals is not relevant in view of its presence as a normal component in animal and human plasma

6. Pharmaceutical particulars

6.1. List of excipients

Water For Injections BP.

6.2. Incompatibilities

The addition of sodium chloride to mannitol 20 or 25% may cause precipitation of the mannitol. As with all parenteral solutions compatibility of the additives with the solution must be assessed before addition. In the absence of compatibility studies, this solution must not be mixed with other medicinal products. Those additives known to be incompatible should not be used

6.3. Shelf life

36 months.

Use immediately after first opening.

6.4. Special precautions for storage

Do not store above 30°C and do not freeze.

Store below 30°C.

6.5. Nature and contents of container

Pack sizes: 100mL, 250mL, and 500 mL.

The bottles are made from Low Density Polyethylene plastic; the bottles are then overwrapped with a protective plastic pouch.

6.6. Special precautions for disposal and other handling

For I.V. Injection

Use as directed by the physician. Keep out of reach of children.

Do not use unless solution is clear and the container is undamaged. Any contents of the product remaining after use should be discarded.

If used as solution additive, the following precautions should be followed. Before adding a drug, verify it is soluble and stable in water at the pH range of the Sodium Chloride 0.9% Intravenous Infusion solution. Additives may be introduced before infusion or during infusion through the injection site.

It is the responsibility of the physician to judge the incompatibility of an additive medication with the Sodium Chloride 0.9% Intravenous Infusion

solution by checking for eventual color change and/or eventual precipitate, insoluble complexes or crystals apparition. The Instructions for Use of the medication to be added must be consulted.

When additive is used, verify isotonicity prior to parenteral administration. Thorough and careful aseptic mixing of any additive is mandatory. Solutions containing additives should be used immediately and not stored.

Adding other medication or using an incorrect administration technique might cause the appearance of fever reactions due to the possible introduction of pyrogens. In case of adverse reaction, infusion must be stopped immediately.

7. MARKETING AUTHORISATION HOLDER

GOODMED HEALTHCARE LIMITED

P.O BOX 76337, NAIROBI, KENYA

8. MARKETING AUTHORISATION NUMBER(S)

H2026/CTD12388/26233

9. DATE OF FIRST AUTHORISATION / RENEWAL OF THE AUTHORISATION

Date of first authorisation: 18.01.2026

10. DATE OF REVISION OF THE TEXT

18.01.2026