

## Summary of Product Characteristics (SmPC)

### 1 NAME OF THE MEDICINAL PRODUCT

Panadol ActiFast 500mg Soluble Tablets

### 2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 500 mg of Paracetamol

Excipients with known effect: each tablet contains 427mg sodium and 50mg sorbitol. For the full

list of excipients see Section 6.1.

### 3 PHARMACEUTICAL FORM

Soluble Tablet.

Round white tablets with bevelled edges.

### 4 CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

As a mild analgesic and antipyretic. The tablets are recommended for use in the short-term management of the symptoms of headache, toothache, menstrual pain, musculoskeletal disorders and for symptoms of common colds and 'flu including sore throat, headache, muscle ache and fever. Panadol may also be used in the symptomatic relief of mild to moderate pain associated with osteoarthritis, as diagnosed by a doctor.

#### 4.2 Posology and method of administration

For oral administration only.

##### **Adults (including the elderly) and children aged 16 years and over:**

One or two tablets dissolved in water up to 4 times daily as required.

##### **Children aged 10-15 years:**

One tablet dissolved in water up to 4 times daily as required.

Children should not be given paracetamol for more than 3 days without consulting a doctor.

Not recommended for children under the age of 10 years. Minimum dosing interval: 4 hours. Not more than 4 doses should be taken in any 24 hour period. Do not exceed the stated dose. Should not be used with other paracetamol containing products.

The lowest dose necessary to achieve efficacy should be used.

##### Renal Impairment

It is recommended, when giving paracetamol to patients with renal impairment, to reduce the dose and to increase the minimum interval between each administration to at least 6 hours unless directed otherwise by a physician. See Table below.

Adults:

Glomerular filtration rate	Dose
10-50 ml/min	500mg every 6 hours
< 10 ml/min	500mg every 8 hours

### Hepatic Impairment

In patients with hepatic impairment or Gilbert's Syndrome, the dose should be reduced or the dosing interval prolonged. The daily dose should not exceed 2g/day unless directed by a physician.

### The Elderly

Experience has indicated that normal adult dosage is usually appropriate. However, in frail, immobile, elderly subjects or in elderly patients with renal or hepatic impairment, a reduction in the amount or frequency of dosing may be appropriate.

The maximum daily dose should not exceed 60mg/kg/day (up to a maximum of 2g per day) in the following situations, unless directed by a physician:

- Weight less than 50kg
- Chronic alcoholism
- Dehydration
- Chronic malnutrition

### **4.3 Contraindications**

Known hypersensitivity to paracetamol or any of the other ingredients.

### **4.4 Special warnings and precautions for use**

Contains paracetamol. Do not use with any other paracetamol-containing products. The concomitant use with other products containing paracetamol may lead to an overdose. Paracetamol overdose may cause liver failure which can lead to liver transplant or death.

Underlying liver disease increases the risk of paracetamol related liver damage. Patients who have been diagnosed with liver or kidney impairment must seek medical advice before taking this medication.

Cases of hepatic dysfunction/failure have been reported in patients with depleted glutathione levels, such as those who are severely malnourished, anorexic, have a low body mass index or are chronic heavy users of alcohol.

Cases of high anion gap metabolic acidosis (HAGMA) due to pyroglutamic acidosis have been reported in patients with severe illness such as severe renal impairment and sepsis, or in patients with malnutrition and other sources of glutathione deficiency (e.g. chronic alcoholism) who were treated with paracetamol at therapeutic dose for a prolonged period or a combination of paracetamol and flucloxacillin. If HAGMA due to pyroglutamic acidosis is suspected, prompt discontinuation of paracetamol and close monitoring is recommended. The measurement of urinary 5-oxoproline may be useful to identify pyroglutamic acidosis as underlying cause of HAGMA in patients with multiple risk factors.

Do not exceed the stated dose.

If symptoms persist, consult your doctor. Prolonged use except under medical supervision may be harmful. This product should only be used when clearly necessary. Keep out of the sight and reach of children.

### **4.5 Interaction with other medicinal products and other forms of interaction**

The speed of absorption of paracetamol may be increased by metoclopramide or domperidone

and absorption reduced by cholestyramine. The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular daily use of paracetamol with increased risk of bleeding; occasional doses have no significant effect. Caution should be taken when paracetamol is used concomitantly with flucloxacillin as concurrent intake has been associated with high anion gap metabolic acidosis due to pyroglutamic acidosis, especially in patients with risks factors (see section 4.4)

#### **4.6 Fertility, pregnancy**

##### **and lactation Pregnancy**

A large amount of data on pregnant women indicate neither malformative, nor fetoneonatal toxicity. Epidemiological studies on neurodevelopment in children exposed to paracetamol in utero show inconclusive results. If clinically needed, paracetamol can be used during pregnancy however it should be used at the lowest effective dose for the shortest possible time and at the lowest possible frequency.

##### **Lactation**

Paracetamol is excreted in breast milk. However, the level of paracetamol present is not considered to be harmful. Available published data do not contraindicate breastfeeding.

#### **4.7 Effects on ability to drive and use machines**

None stated.

#### **4.8 Undesirable effects**

Events reported from extensive post-marketing experience at therapeutic/labelled dose and considered attributable are tabulated below by System Organ Class and frequency.

Frequencies are defined as: very common ( $\geq 1/10$ ), common ( $\geq 1/100$ ,  $< 1/10$ ), uncommon ( $\geq 1/1,000$ ,  $< 1/100$ ), rare ( $\geq 1/10,000$ ,  $< 1/1,000$ ), very rare ( $< 1/10,000$ ), not known (cannot be estimated from available data).

Adverse event frequencies have been estimated from spontaneous reports received through post marketing data.

<b>Body System</b>	<b>Undesirable Effect</b>	<b>Frequency</b>
<b>Paracetamol</b>		
Blood and lymphatic system disorders	Thrombocytopenia	Very rare
Metabolism and nutrition disorders	High anion gap metabolic acidosis*	Not known
Immune System disorders	Anaphylaxis, Cutaneous hypersensitivity reactions, including, among others, skin rashes, angioedema, Stevens Johnson syndrome and Toxic Epidermal Necrolysis.  Very rare cases of serious skin reactions	Very rare
Respiratory, thoracic and mediastinal disorders	Bronchospasm in patients sensitive to aspirin and other NSAIDs	Very rare
Hepatobiliary disorders	Hepatic dysfunction	Very rare

\* Cases of high anion gap metabolic acidosis due to pyroglutamic acidosis have been observed in patients with risk factors using paracetamol (see section 4.4). Pyroglutamic acidosis may occur as a consequence of low glutathione levels in these patients.

#### Reporting of suspected adverse reactions

Reporting of suspected adverse reactions: Healthcare professionals are requested to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS) <https://pv.pharmacyboardkenya.org>

#### **4.9 Overdose Symptoms and signs**

Paracetamol overdose may cause liver failure which can lead to liver transplant or death. Acute pancreatitis has been observed, usually with hepatic dysfunction and liver toxicity.

There is a risk of poisoning with paracetamol particularly in elderly subjects, young children, patients with liver disease, cases of chronic alcoholism and in patients with chronic malnutrition. Overdosing may be fatal in these cases.

Symptoms generally appear within the first 24 hours and may comprise: nausea, vomiting, anorexia, pallor, and abdominal pain, or patients may be asymptomatic.

Overdose of paracetamol in a single administration in adults or in children can cause liver cell necrosis likely to induce complete and irreversible necrosis, resulting in hepatocellular insufficiency, metabolic acidosis and encephalopathy which may lead to coma and death. Simultaneously, increased levels of hepatic transaminases (AST, ALT), lactate dehydrogenase and bilirubin are observed together with increased prothrombin levels that may appear 12 to 48 hours after administration.

Liver damage is likely in adults who have taken more than the recommended amounts of paracetamol. It is considered that excess quantities of toxic metabolite (usually adequately detoxified by glutathione when normal doses of paracetamol are ingested), become irreversibly bound to liver tissue.

Some patients may be at increased risk of liver damage from

paracetamol toxicity. Risk Factors include: If the patient;

- Is on long-term treatment with carbamazepine, phenobarbitone, phenytoin, primidone, rifampicin, St John's Wort or other drugs that induce liver enzymes.
- Regularly consumes ethanol in excess of recommended amounts
- Is likely to be glutathione depleted e.g. eating disorders, cystic fibrosis, HIV infection, starvation, cachexia

### Emergency Procedure:

Immediate transfer to hospital.

Blood sampling to determine initial paracetamol plasma concentration. In the case of a single acute overdose, paracetamol plasma concentration should be measured 4 hours post ingestion.

Administration of activated charcoal should be considered if >150mg/kg paracetamol has been taken within 1 hour.

The antidote N-acetylcysteine, should be administered as soon as possible in accordance with

National treatment guidelines. Symptomatic treatment should be implemented.

### **Sodium bicarbonate**

High doses of sodium bicarbonate may be expected to induce gastrointestinal symptoms including belching and nausea. In addition, high doses of sodium bicarbonate may cause hypernatraemia; electrolytes should be monitored and patients managed accordingly.

## **5 PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Paracetamol has analgesic and antipyretic actions.

### **5.2 Pharmacokinetic properties**

Paracetamol is rapidly and almost completely absorbed from the gastro-intestinal tract. Concentration in plasma reaches a peak in 30-60 minutes. Plasma half-life is 1-4 hours. Paracetamol is relatively uniformly distributed throughout most body fluids. Plasma protein binding is variable. Excretion is almost exclusively renal, in the form of conjugated metabolites.

### **5.3 Preclinical safety data**

Conventional studies using the currently accepted standards for the evaluation of toxicity to reproduction and development are not available.

## **6 PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

Saccharin sodium  
 Citric acid (anhydrous)  
 Sorbitol powder E420  
 Sodium hydrogen carbonate  
 Sodium laurilsulfate  
 Anhydrous sodium carbonate  
 Povidone

Dimeticone

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

4 years.

## **6.4 Special precautions for storage**

Do not store above 25°C. Keep in the original package in order to protect from moisture.

## **6.5 Nature and contents of container**

Panadol Soluble tablets are packed into PFPF laminate sachets. The strips are then enclosed in cardboard cartons to give pack sizes of 12, 24 or 60 tablets. Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal and other handling**

No special requirements.

## **7 MARKETING AUTHORISATION HOLDER**

Haleon Ireland  
Limited 12  
Riverwalk  
Citywest Business  
Campus Dublin 24  
Ireland

## **8 MARKETING AUTHORISATION NUMBER**

H97/456

## **9 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 29<sup>th</sup> June 2007

Date of last renewal: 29<sup>th</sup> June 2012

## **10 DATE OF REVISION OF THE TEXT**

28/02/202