

Summary of Product Characteristics

1 Name of the medicinal product

Tasemic 500mg/5ml Solution for Injection

2 Qualitative and quantitative composition

Each ampoule contains 500mg of tranexamic acid per 5ml (100mg/ml). For a full list of excipients, see section 6.1

3 Pharmaceutical form

Solution for injection
Clear, colourless solution

4 Clinical particulars

4.1 Therapeutic indications

Prevention and treatment of haemorrhages due to general or local fibrinolysis in adults and children from one year.

Specific indications include:

- Haemorrhage caused by general or local fibrinolysis such as:
- Menorrhagia and metrorrhagia,
- Gastrointestinal bleeding,
- Haemorrhagic urinary disorders, further to prostate surgery or surgical procedures affecting the urinary tract,
- Ear Nose Throat surgery (adenoidectomy, tonsillectomy, dental extractions),
- Gynaecological surgery or disorders of obstetric origin,
- Thoracic and abdominal surgery and other major surgical intervention such as cardiovascular surgery,- Management of haemorrhage due to the administration of a fibrinolytic agent.

4.2 Posology and method of administration

Posology

Adults

Unless otherwise prescribed, the following doses are recommended:

1. Standard treatment of local fibrinolysis:

500mg (1 ampoule of 5ml) to 1000mg (2 ampoules of 5ml) tranexamic acid by slow intravenous injection (= 1ml/minute) two to three times daily

2. Standard treatment of general fibrinolysis:

1000mg (2 ampoules of 5ml) tranexamic acid by slow intravenous injection (= 1 ml/minute) every 6 to 8 hours, equivalent to 15 mg/kg body weight.

Renal impairment

In renal insufficiency leading to a risk of accumulation, the use of tranexamic acid is contra-indicated in patient with severe renal impairment (see section 4.3). For patient with mild to moderate renal impairment, the dosage of tranexamic acid should be reduced according to the serum creatinine level:

Serum creatinine	Dose IV	Administration
------------------	---------	----------------

µmol/l	mg/10 ml		
120 to 249	1.35 to 2.82	10 mg/kg BW	Every 12 hours
250 to 500	2.82 to 5.65	10 mg/kg BW	Every 24 hours
> 500	> 5.65	5 mg/kg BW	Every 24 hours

Hepatic impairment

No dose adjustment is required in patient with hepatic impairment.

Paediatric Population:

In children from 1 year, for current approved indications as described in section 4.1, the dosage is in the region of 20 mg/kg/day. However, data on efficacy, posology and safety for these indications are limited. The efficacy, posology and safety of tranexamic acid in children undergoing cardiac surgery have not been fully established. Currently available data are limited and are described in section 5.1.

Elderly:

No reduction in dosage is necessary unless there is evidence of renal failure.

Method of administration

The administration is strictly limited to slow intravenous injection.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1;
- Acute venous or arterial thrombosis (see section 4.4);
- Fibrinolytic conditions following consumption coagulopathy except in those with predominant activation of the fibrinolytic system with acute severe bleeding (see section 4.4);
- Severe renal impairment (risk of accumulation);
- History of convulsions;
- Intrathecal and intraventricular injection, intracerebral application (risk of cerebral oedema and convulsions).

4.4 Special warnings and precautions for use

The indications and method of administration indicated above should be followed strictly:

- Intravenous injections should be given very slowly
- Tranexamic acid should not be administered by the intramuscular route.

Convulsions

Cases of convulsions have been reported in association with tranexamic acid treatment. In coronary artery bypass graft (CABG) surgery, most of these cases were reported following intravenous (i.v.) injection of tranexamic acid in high doses. With the use of the recommended lower doses of Tranexamic Acid,

the incidence of post-operative seizures was the same as that in untreated patients.

Visual disturbances

Attention should be paid to possible visual disturbances including visual impairment, vision blurred, impaired colour vision and if necessary the treatment should be discontinued. With continuous long-term use of Tranexamic Acid solution for injection, regular ophthalmologic examinations (eye examinations including visual acuity, colour vision, fundus, visual field etc.) are indicated. With pathological ophthalmic changes, particularly with diseases of the retina, the physician must decide after consulting a specialist on the necessity for the long-term use of Tranexamic Acid solution for injection in each individual case.

Haematuria

In case of haematuria from the upper urinary tract, there is a risk for urethral obstruction.

Thromboembolic events

Before use of Tranexamic Acid, risk factors of thromboembolic disease should be considered. In patients with a history of thromboembolic diseases or in those with increased incidence of thromboembolic events in their family history (patients with a high risk of thrombophilia), Tranexamic acid solution for injection should only be administered if there is a strong medical indication after consulting a physician experienced in hemostaseology and under strict medical supervision (see section 4.3).

Tranexamic acid should be administered with care in patients receiving oral contraceptives because of the increased risk of thrombosis (See section 4.5.).

Disseminated intravascular coagulation

Patients with disseminated intravascular coagulation (DIC) should in most cases not be treated with tranexamic acid (see section 4.3). If tranexamic acid is given it must be restricted to those in whom there is predominant activation of the fibrinolytic system with acute severe bleeding. Characteristically, the haematological profile approximates to the following: reduced euglobulin clot lysis time; prolonged prothrombin time; reduced plasma levels of fibrinogen, factors V and VIII, plasminogen fibrinolysin and alpha-2 macroglobulin; normal plasma levels of P and P complex; i.e. factors II (prothrombin), VIII and X; increased plasma levels of fibrinogen degradation products; a normal platelet count. The foregoing presumes that the underlying disease state does not of itself modify the various elements in this profile. In such acute cases a single dose of 1g tranexamic acid is frequently sufficient to control bleeding. Administration of Tranexamic acid in DIC should be considered only when appropriate haematological laboratory facilities and expertise are available.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed. Simultaneous treatment with anticoagulants must take place under the strict supervision of a physician experienced in this field. Medicinal products that act on haemostasis should be given with caution to patients treated with tranexamic acid. There is a theoretical risk of increased thrombus-formation potential, such as with oestrogens. Alternatively, the antifibrinolytic action of the drug may be antagonised with thrombolytic drugs.

4.6 Fertility, pregnancy and lactation

Women of childbearing potential have to use effective contraception during treatment.

Pregnancy

There are insufficient clinical data on the use of tranexamic acid in pregnant women. As a result, although studies in animals do not indicate teratogenic effects, as a precaution for use, tranexamic acid is not recommended during the first trimester of pregnancy. Limited clinical data on the use of tranexamic acid in different clinical haemorrhagic settings during the second and third trimesters did not identify deleterious effect for the foetus. Tranexamic acid should be used throughout pregnancy only if the expected benefit justifies the potential risk.

Breastfeeding

Tranexamic acid is excreted in human milk. Therefore, breastfeeding is not recommended.

Fertility

There are no clinical data on the effects of tranexamic acid on human fertility.

4.7 Effects on ability to drive and use machines

No studies have been performed on the ability to drive and use machines.

4.8 Undesirable effects

The ADRs reported from clinical studies and post-marketing experience are listed below according to system organ class.

Tabulated list of adverse reactions

Adverse reactions reported are presented in table below. Adverse reactions are listed according to MedDRA primary system organ class. Within each system organ class, adverse reactions are ranked by frequency.

Term	Frequency of occurrence
Very common	($\geq 1/10$)
Common	($\geq 1/100$ to $< 1/10$)
Uncommon	($\geq 1/1\ 000$ to $< 1/100$)
Rare	($\geq 1/10\ 000$ to $< 1/1\ 000$)

Very rare	(<1/10 000)
Not known	(cannot be estimated from the available data)

System Organ Class	Frequency	Undesirable Effects
Skin and subcutaneous tissues disorders	Uncommon	Dermatitis allergic
Gastrointestinal disorders	Common	Diarrhoea, Vomiting, Nausea
Nervous system disorders	Not known	Dizziness, convulsions particularly in case of misuse (refer to sections 4.3 and 4.4)
Eye disorders	Not known	Visual disturbances including impaired colour vision
Vascular disorders	Not known	Malaise with hypotension, with or without loss of consciousness (generally following a too fast intravenous injection, exceptionally after oral administration) Arterial or venous thrombosis at any sites
Immune system disorders	Not known	Hypersensitivity reactions including anaphylaxis

4.9 Overdose

No cases of overdose have been reported.

Signs and symptoms may include nausea, vomiting, dizziness, headache, hypotension, and convulsions. It has been shown that convulsions tend to occur at higher frequency with increasing dose. Management of overdose should be supportive.

Maintain a high fluid intake to promote renal excretion. Anticoagulant treatment should be considered. There is a risk of thrombosis in predisposed individuals.

5 Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antihemorrhagics,

Antifibrinolytics, Aminoacids ATC code: B02A A02

Tranexamic acid exerts an anti haemorrhagic activity by inhibiting the fibrinolytic properties of plasmin.

A complex involving tranexamic acid, plasminogen is constituted; the tranexamic acid being linked to plasminogen when transformed into plasmin.

The activity of the tranexamic acid-plasmin complex on the activity on fibrin is lower than the activity of free plasmin alone.

In vitro studies showed that high tranexamic dosages decreased the activity of complement.

Paediatric population

In children over one year old: Literature review identified 12 efficacy studies in paediatric cardiac surgery which have included 1073 children, 631 having received tranexamic acid. Most of them were controlled versus placebo. Studied population was heterogenic in terms of age, surgery types, dosing schedules. Study results with tranexamic acid suggest reduced blood loss and reduced blood product requirements in paediatric cardiac surgery under cardiopulmonary bypass when there is a high risk of haemorrhage, especially in cyanotic patients or patients undergoing repeat surgery.

The most adapted dosing schedule appeared to be:

- first bolus of 10 mg/kg after induction of anaesthesia and prior to skin incision,
- continuous infusion of 10 mg/kg/h or injection into the CPB pump prime at a dose adapted on the CPB procedure, either according to patient weight with a 10 mg/kg dose, either according to CPB pump prime volume,
- last injection of 10 mg/kg at the end of CPB.

While studied in very few patients, the limited data suggest that continuous infusion is preferable, since it would maintain therapeutic plasma concentration throughout surgery.

No specific dose-effect study or PK study has been conducted in children.

5.2 Pharmacokinetic properties

Absorption

Peak plasma concentrations of tranexamic acid are obtained rapidly after a short intravenous infusion after which plasma concentrations decline in a multi-exponential manner.

Distribution

The plasma protein binding of tranexamic acid is about 3% at therapeutic plasma levels and seems to be fully accounted for by its binding to plasminogen. Tranexamic acid does not bind to serum albumin. The initial volume of distribution is about 9 to 12 liters.

Tranexamic acid passes through the placenta. Following administration of an intravenous injection of 10 mg/kg to 12 pregnant women, the concentration of tranexamic acid in serum ranged 10-53 µg/mL while that in cord blood ranged 4-31 µg/mL. Tranexamic acid diffuses rapidly into joint fluid and the synovial membrane. Following administration of an intravenous injection of 10 mg/kg to 17 patients undergoing knee surgery, concentrations in the joint fluids were similar to those seen in corresponding serum samples. The concentration of tranexamic acid in a number of other tissues is a fraction of that observed in the blood (breast milk, one hundredth; cerebrospinal fluid, one tenth; aqueous humor, one tenth). Tranexamic acid has been detected in semen where it inhibits fibrinolytic activity but does not influence sperm migration.

Elimination

It is excreted mainly in the urine as unchanged drug. Urinary excretion via glomerular filtration is the main route of elimination. Renal clearance is equal to plasma clearance (110 to 116 mL/min). Excretion of tranexamic acid is about 90% within the first 24 hours after intravenous administration of 10 mg/kg body weight. Half-life of tranexamic acid is approximately 3 hours.

Special populations

Plasma concentrations increase in patients with renal failure.

No specific PK study has been conducted in children.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and toxicity to reproduction. Epileptogenic activity has been observed in animals with intrathecal use of tranexamic acid.

6 Pharmaceutical particulars

6.1 List of excipients

Water for Injections

6.2 Incompatibilities

Tranexamic Acid Solution for Injection should not be added to blood for transfusion, or to injections containing penicillin.

6.3 Shelf life

2 years

6.4 Special precautions for storage

Store below 30°C. Keep the ampoules in the original packaging to protect from light. Do not freeze.

6.5 Nature and contents of container

Type I glass 5ml ampoules packed in outer cardboard carton. Pack of 10 ampoules.

6.6 Special precautions for disposal and other handling

Discard if solution is coloured. For single use only, any remaining solution should be discarded.

7 Marketing authorisation holder

Tasa Pharma Limited
Unit C1-C3
Kay complex
Nairobi
Kenya

8. Marketing Authorization Number

H2022/CTD8986/20508

9. Date of first authorization/renewal of the authorization

21/07/2021

10. Date of revision of the text

21/07/2021