

SUMMARY PRODUCT CHARACTERISTICS

TRANEXAMIC ACID TABLETS BP 500 MG



1. NAME OF THE MEDICINAL PRODUCT

1.1 Name of the Medicinal Product

UBAMIC

TRANEXAMIC ACID TABLETS BP 500 MG

1.2 Strength

500 mg

1.3 Pharmaceutical Form

Tablet, Oral dosage form

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film coated tablet contains:

Tranexamic acid BP 500 mg

Excipients q.s.

Colour: Titanium Dioxide BP

3. PHARMACEUTICAL FORM

Tablet

White to off White colored oblong shaped, film coated tablet breakline on one side & plain on another side.

4. CLINICAL PARTICULARS

4.1. Therapeutic indications

Tranexamic Acid is indicated for short term use for haemorrhage or risk of haemorrhage in those with increased fibrinolysis or fibrinogenolysis. Local fibrinolysis as occurs in the following conditions:

- 1.a) Prostatectomy and bladder surgery
 - b) Menorrhagia
 - c) Epistaxis
 - d) Conisation of the cervix
 - e) Traumatic hyphaema
2. Management of dental extraction in haemophiliacs.
 3. Hereditary angioneurotic oedema.

4.2. Posology and method of administration

Posology

Adults:

Local Fibrinolysis: The recommended standard dose is 15-25mg/kg bodyweight (i.e. 2-3 tablets) two to three times daily. For the indications listed below the following doses may be used:

1a. Prostatectomy: Prophylaxis and treatment of haemorrhage in high risk patients should commence pre- or post-operatively with an injectable form; thereafter 2 tablets three to four times daily until macroscopic haematuria is no longer present.

1b. Menorrhagia: Recommended dosage is 2 tablets 3 times daily as long as needed for up to 4 days. If very heavy menstrual bleeding, dosage may be increased. A total dose of 4g daily (8 tablets) should not be exceeded. Treatment with tranexamic acid should not be initiated until menstrual bleeding has started.

1c. Epistaxis: When repeated bleeding is anticipated oral therapy (2 tablets three times daily) should be administered for 7 days.

1d. Cervix Conisation: 3 tablets three times daily

1e. Traumatic Hyphaema: 2-3 tablets 3 times daily. The dose is based on 25mg/kg three times a day.

2. Haemophilia: In the management of dental extractions 2-3 tablets every eight hours. The dose is based on 25mg/kg.

3. Hereditary angioneurotic oedema: Some patients are aware of the onset of illness; suitable treatment for these patients is intermittently 2-3 tablets two to three times daily for some days. Other patients are treated continuously at this dosage.

Pediatric population:

This should be calculated according to bodyweight at 25mg/kg per dose at the adult dosing frequencies. However, data on efficacy, posology and safety for these indications are limited.

Elderly:

No reduction in dosage is necessary unless there is evidence of renal failure (see guidelines below).

Renal insufficiency

By extrapolation from clearance data relating to the intravenous dosage form, the following reduction in the oral dosage is recommended for patients with mild to moderate renal insufficiency:

SUMMARY PRODUCT CHARACTERISTICS**TRANEXAMIC ACID TABLETS BP 500 MG**

Serum Creatinine($\mu\text{mol/l}$)	Oral Dose	Dose Frequency
120-249	15 mg/kg body weight	twice daily
250-500	15 mg/kg body weight	daily

Method of administration

Oral

4.3. Contra-indications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1

Severe renal failure because of risk of accumulation.

Active thromboembolic disease.

History of venous or arterial thrombosis

Fibrinolytic conditions following consumption coagulopathy

History of convulsions

4.4. Special warnings and special precautions for use

In case of haematuria of renal origin (especially in haemophilia), there is a risk of mechanical anuria due to formation of a ureteral clot.

In the long-term treatment of patients with hereditary angioneurotic oedema, regular eye examinations (e.g. visual acuity, slit lamp, intraocular pressure, visual fields) and liver function tests should be performed.

Patients with irregular menstrual bleeding should not use Tranexamic Acid until the cause of irregular bleeding has been established. If menstrual bleeding is not adequately reduced by Tranexamic Acid, an alternative treatment should be considered.

Tranexamic acid should be administered with care in patients receiving oral contraceptives because of the increased risk of thrombosis.

Patients with a previous thromboembolic event and a family history of thromboembolic disease (patients with thrombophilia) should use Tranexamic Acid only if there is a strong medical indication and under strict medical supervision.

The blood levels are increased in patients with renal insufficiency. Therefore a dose reduction is recommended (see section 4.2).

SUMMARY PRODUCT CHARACTERISTICS

TRANEXAMIC ACID TABLETS BP 500 MG



The use of tranexamic acid in cases of increased fibrinolysis due to disseminated intravascular coagulation is not recommended.

Patients who experience visual disturbance should be withdrawn from treatment.

Clinical experience with Tranexamic Acid in menorrhagic children under 15 years of age is not available.

4.5. Interactions with other Drug products and other forms of interaction

Tranexamic Acid will counteract the thrombolytic effect of fibrinolytic preparations.

4.6. Pregnancy and lactation

Pregnancy

Although there is no evidence from animal studies of a teratogenic effect, the usual caution with use of drugs in pregnancy should be observed.

Tranexamic acid crosses the placenta.

Breast-feeding

Tranexamic acid passes into breast milk to a concentration of approximately one hundredth of the concentration in the maternal blood. An antifibrinolytic effect in the infant is unlikely.

4.7. Effects on ability to drive and use machines

Tranexamic Acid has no or negligible influence on the ability to drive and use machines.

4.8. ADVERSE REACTIONS

Adverse effects have been ranked under headings of frequency using the following convention:

Very common ($\geq 1/10$)
Common ($\geq 1/100, < 1/10$)
Uncommon ($\geq 1/1000, < 1/100$)
Rare ($\geq 1/10,000, < 1/1,000$)
Very rare ($< 1/10,000$) including isolated reports
Not known (cannot be estimated from the available data).

The following undesirable effects have been reported

Immune system disorders

Very rare: Hypersensitivity reactions including anaphylaxis

Gastrointestinal disorders

SUMMARY PRODUCT CHARACTERISTICS

TRANEXAMIC ACID TABLETS BP 500 MG



Very rare: Digestive effects such as nausea, vomiting and diarrhoea may occur but disappear when the dosage is reduced

Skin and subcutaneous tissue disorders

Rare: Allergic skin reactions.

Vascular disorders

Rare: thromboembolic events.

Very rare: Arterial or venous thrombosis at any sites

Eye disorders

Rare: impaired colour vision and other visual disturbances, retinal/artery occlusion

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the Yellow Card Scheme (www.mhra.gov.uk/yellowcard).

4.9 Overdose

Symptoms may be nausea, vomiting, orthostatic symptoms and/or hypotension. Initiate vomiting, then stomach lavage, and charcoal therapy. Maintain a high fluid intake to promote renal excretion. There is a risk of thrombosis in predisposed individuals. Anticoagulant treatment should be considered.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antifibrinolytic agent. ATC code: B02AA02

Tranexamic acid is an antifibrinolytic compound which is a potent competitive inhibitor of the activation of plasminogen to plasmin. At much higher concentrations it is a non competitive inhibitor of plasmin. The inhibitory effect of tranexamic acid in plasminogen activation by urokinase has been reported to be 6-100 times and by streptokinase 6-40 times greater than that of aminocaproic acid. The antifibrinolytic activity of tranexamic acid is approximately ten times greater than that of aminocaproic acid.

5.2 Pharmacokinetic Properties

Absorption

Peak plasma Tranexamic acid concentration is obtained immediately after intravenous administration (500mg). Then concentration decreases until the 6th hour. Elimination half-life is about 3 hours.

Distribution

SUMMARY PRODUCT CHARACTERISTICS

TRANEXAMIC ACID TABLETS BP 500 MG



Tranexamic acid administered parenterally is distributed in a two compartment model. Tranexamic acid is delivered in the cell compartment and the cerebrospinal fluid with delay. The distribution volume is about 33% of the body mass.

Tranexamic acid crossed the placenta, and may reach one hundredth of the serum peak concentration in the milk of lactating women.

Elimination

Tranexamic acid is excreted in urine as unchanged compound. 90% of the administered dose is excreted by the kidney in the twelve first hours after administration (glomerular excretion without tubular reabsorption).

Following oral administration, 1.13% and 39% of the administered dose were recovered after 3 and 24 hours respectively.

Plasma concentrations are increased in patients with renal insufficiency.

5.3. Preclinical safety data

There are no preclinical data of relevance to the prescriber which are additional to that already included in other sections of the Summary of Product Characteristics.

6. PHARMACEUTICAL PARTICULARS

6.1. List of Excipients

Maize Starch, Lactose, Microcrystalline cellulose, Sodium benzoate, Purified Talc, Magnesium Stearate, Cross Carmellose Sodium, Colloidal Silicon Dioxide, Isopropyl alcohol, Methylene Dichloride, Titanium Dioxide BP.

6.2. Incompatibilities

None

6.3. Shelf life

36 Months.

6.4 Special precautions for storage

Store below above 30 °C.

Protect from light & moisture.

6.5. Nature and contents of container

Aluminum - Aluminum Blister Pack

10 Tablet are blister packed with Aluminum - Aluminum foil; such 3 blister are packed in one carton pack.

Pack size: 3 x 10 Tablets (i.e. 30 Tablets) in one carton box along with packing leaflet.

SUMMARY PRODUCT CHARACTERISTICS	
TRANEXAMIC ACID TABLETS BP 500 MG	

**6.6. Special precautions for disposal and other handling**

There are no special storage precautions. Any unused product or waste material should be disposed of in accordance with local requirements.

7.0 Registrant

Eastleigh Pharmaceuticals Co. Ltd
P.O Box 167-00610 Nairobi, Kenya

8.0 Manufacturer

MARS REMEDIES PVT LTD
Address: 635, GIDC Estate, Waghodia-391760, Vadodara, GUJARAT INDIA

9.0 Date of Publication or Revision

Last revised on Dec-2020

10. DOSIMETRY (IF APPLICABLE)

Not Applicable

11. Instructions for Preparation of Radiopharmaceuticals (If Applicable):

Not Applicable