

1. Name of the medicinal product

Virux 400 Tablet

2. Qualitative and quantitative composition

Each Tablet contains Acyclovir USP 400 mg.

3. Pharmaceutical form

Tablet

A white to off white round shaped tablet having both side plain. Free from any visible defects.

4. Clinical particulars

4.1 Therapeutic indications

Acyclovir Tablets are indicated for the treatment of herpes simplex virus infections of the skin and mucous membranes including initial and recurrent genital herpes (excluding neonatal HSV and severe HSV infections in immunocompromised children).

Acyclovir Tablets are indicated for the suppression (prevention of recurrences) of recurrent herpes simplex infections in immunocompetent patients.

Acyclovir Tablets are indicated for the prophylaxis of herpes simplex infections in immunocompromised patients.

Acyclovir Tablets are indicated for the treatment of varicella (chickenpox) and herpes zoster (shingles) infections

4.2 Posology and method of administration

1. Dosage in adults

Treatment of herpes simplex infections: 200 mg Acyclovir should be taken five times daily at approximately four hourly intervals omitting the night time dose. Treatment should continue for 5 days, but in severe initial infections this may have to be extended.

In severely immunocompromised patients (e.g. after marrow transplant) or in patients with impaired absorption from the gut the dose can be doubled to 400 mg Acyclovir, or alternatively, intravenous dosing could be considered.

Dosing should begin as early as possible after the start of an infection; for recurrent episodes this should preferably be during the prodromal period or when lesions first appear.

Suppression of herpes simplex infections in immunocompetent patients: 200 mg Acyclovir should be taken four times daily at approximately six-hourly intervals.

Many patients may be conveniently managed on a regimen of 400 mg Acyclovir twice daily at approximately twelve-hourly intervals.

Dosage titration down to 200 mg Acyclovir taken thrice daily at approximately eight-hourly intervals or even twice daily at approximately twelve-hourly intervals may prove effective.

Some patients may experience break-through infection on total daily doses of 800 mg Acyclovir.

Therapy should be interrupted periodically at intervals of six to twelve months, in order to observe possible changes in the natural history of the disease.

Prophylaxis of herpes simplex infections in immunocompromised patients: 200 mg Acyclovir should be taken four times daily at approximately six-hourly intervals.

In severely immunocompromised patients (e.g. after marrow transplant) or in patients with impaired absorption from the gut, the dose can be doubled to 400 mg Acyclovir, or alternatively, intravenous dosing could be considered.

The duration of prophylactic administration is determined by the duration of the period at risk.

Treatment of varicella and herpes zoster infections: 800 mg Acyclovir should be taken five times daily at approximately four-hourly intervals, omitting the night time dose. Treatment should continue for seven days.

In severely immunocompromised patients (e.g. after marrow transplant) or in patients with impaired absorption from the gut, consideration should be given to intravenous dosing.

Dosing should begin as early as possible after the start of an infection: Treatment of herpes zoster yields better results if initiated as soon as possible after the onset of the rash. Treatment of chickenpox in immunocompetent patients should begin within 24 hours after onset of the rash.

2. Paediatric population

Treatment of herpes simplex infections, and prophylaxis of herpes simplex infections in the immunocompromised: Children aged two years and over should be given adult dosages and children below the age of two years should be given half the adult dose.

For treatment on neonatal herpes virus infections, intravenous Acyclovir is recommended.

Treatment of varicella infection

6 years and over: 800 mg Acyclovir four
times daily 2 - 5 years: 400 mg Acyclovir
four times daily Under 2 years: 200 mg
Acyclovir four times daily Treatment
should continue for five days.

Dosing may be more accurately calculated as 20 mg/kg bodyweight (not to exceed 800 mg) Acyclovir four times daily.

No specific data are available on the suppression of herpes simplex infections or the treatment of herpes zoster infections in immunocompetent children.

3. Dosage in the elderly

The possibility of renal impairment in the elderly must be considered and the dosage should be adjusted accordingly (see Dosage in renal impairment below). Adequate hydration of elderly patients taking high oral doses of Acyclovir should be maintained.

4. Dosage in renal impairment

Caution is advised when administering Acyclovir to patients with impaired renal function. Adequate hydration should be maintained.

In the management of herpes simplex infections in patients with impaired renal function, the recommended oral doses will not lead to accumulation of Acyclovir above levels that have been established safe by intravenous infusion. However, for patients with severe renal impairment (creatinine clearance less than 10 ml/minute) an adjustment of dosage to 200 mg Acyclovir twice daily at approximately twelve-hourly intervals is recommended.

In the treatment of herpes zoster infections it is recommended to adjust the dosage to 800 mg Acyclovir twice daily at approximately twelve hourly intervals for patients with severe renal impairment (creatinine clearance less than 10 ml/minute), and to 800 mg Acyclovir three times daily at intervals of approximately eight hours for patients with moderate renal impairment (creatinine clearance in the range 10 – 25 ml/minute).

Method of administration:

Oral. Acyclovir tablets may be dispersed in a minimum of 50 ml of water or swallowed whole with a little water. Ensure that patients on high doses of Acyclovir are adequately hydrated.

4.3 Contraindications

Acyclovir is contraindicated in patients known to be hypersensitive to Acyclovir.

4.4 Special warnings and precautions for use

Use in patients with renal impairment and in elderly patients:

Acyclovir is eliminated by renal clearance; therefore, the dose must be adjusted in patients with renal impairment.

Elderly patients are likely to have reduced renal function and therefore the need for dose adjustment must be considered in this group of patients. Both elderly

patients and patients with renal impairment are at increased risk of developing neurological side effects and should be closely monitored for evidence of these effects. In the reported cases, these reactions were generally reversible on discontinuation of treatment.

Prolonged or repeated courses of acyclovir in severely immune-compromised individuals may result in the selection of virus strains with reduced sensitivity, which may not respond to continued acyclovir treatment.

Hydration status: Care should be taken to maintain adequate hydration in patients receiving high oral doses of acyclovir.

The risk of renal impairment is increased by use with other nephrotoxic drugs.

The data currently available from clinical studies is not sufficient to conclude that treatment with aciclovir reduces the incidence of chickenpox-associated complications in immunocompetent patients.

Paediatric population:

Oral aciclovir should be used in paediatric population mainly for the treatment of non-severe skin and mucosa HSV infections. For the treatment of neonatal HSV and severe HSV infections in immunocompromised children IV aciclovir should be used.

Interaction with other medicinal products and other forms of interaction

Acyclovir is eliminated primarily unchanged in the urine via active renal tubular secretion. Any drugs administered concurrently that compete with this mechanism may increase acyclovir plasma concentrations.

Probenecid and cimetidine increase the AUC of acyclovir by this mechanism, and reduce acyclovir renal clearance. Similarly increases in plasma AUCs of acyclovir and of the inactive metabolite of mycophenolate mofetil, an immunosuppressant agent used in transplant patients have been shown when the drugs are co administered. However, no dosage adjustment is necessary because of the wide therapeutic index of acyclovir.

An experimental study on five male subjects indicates that concomitant therapy with acyclovir increases AUC of totally administered theophylline with approximately 50%. It is recommended to measure plasma concentrations during concomitant therapy with acyclovir.

4.6 Fertility, pregnancy and lactation

Pregnancy category B. Acyclovir should not be used during pregnancy unless the potential benefit justifies the potential risk to the fetus.

Caution should be exercised when it is administered to a nursing mother.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

Undesirable effects

Nausea, vomiting, diarrhoea, abdominal pains, fatigue, fever, pruritus, rashes, increases in blood urea and creatinine, Hepatitis, jaundice, reversible rises in bilirubin and liver related enzyme, angioedema.

Reporting of suspected adverse reactions: Healthcare professionals are requested to report any suspected adverse reactions via national Pharmacovigilance Electronic Reporting Systems.

4.8 Overdose

Symptoms and signs

Acyclovir is only partly absorbed in the gastrointestinal tract. Patients have ingested overdoses of up to 20g acyclovir on a single occasion, usually without toxic effects. Accidental, repeated overdoses of oral acyclovir over several days have been associated with gastrointestinal effects (such as nausea and vomiting) and neurological effects (headache and confusion).

Overdosage of intravenous acyclovir has resulted in elevations of serum creatinine, blood urea nitrogen and subsequent renal failure. Neurological effects including confusion, hallucinations, agitation, seizures and coma have been described in association with intravenous overdosage.

Management

Patients should be observed closely for signs of toxicity. Haemodialysis significantly enhances the removal of acyclovir from the blood and may, therefore, be considered a management option in the event of symptomatic overdose.

Reporting of suspected adverse reactions: Healthcare professionals are asked to report any suspected adverse reactions via pharmacy and poisons board, Pharmacovigilance Electronic Reporting System (PvERS)
<https://pv.pharmacyboardkenya.org>

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Direct acting antivirals, Nucleosides and nucleotides excl. reverse transcriptase inhibitors.

ATC code: J05AB01

Mechanism of action

Acyclovir is a synthetic purine nucleoside analogue with *in vitro* and *in vivo* inhibitory activity against human herpes viruses, including herpes simplex virus (HSV) types I and II and varicella zoster virus (VZV).

The inhibitory activity of acyclovir for HSV I, HSV II and VZV is highly selective. The enzyme thymidine kinase (TK) of normal, uninfected cells does not use acyclovir effectively as a substrate, hence toxicity of mammalian host cells is low; however, TK encoded by HSV and VZV converts acyclovir to acyclovir monophosphate, a nucleoside analogue which is further converted to the diphosphate and finally to the triphosphate by cellular enzymes. Acyclovir triphosphate interferes with the viral DNA polymerase and inhibits viral DNA replication with resultant chain termination following its incorporation into the viral DNA.

Prolonged or repeated courses of acyclovir in severely immuno-compromised individuals may result in the selection of virus strains with reduced sensitivity, which may not respond to continued acyclovir treatment. Most of the clinical isolates with reduced sensitivity have been relatively deficient in viral TK, however, strains with altered viral TK or viral DNA polymerase have also been reported. *In vitro* exposure of HSV isolates to acyclovir can also lead to the emergence of less sensitive strains. The relationship between the *in vitro*-determined sensitivity of HSV isolates and clinical response to Acyclovir therapy is not clear.

5.2 Pharmacokinetic properties

Acyclovir is only partially absorbed from the gut. Mean steady state peak plasma concentrations (C^{ss}_{max}) following doses of 200 mg administered four-hourly were 3.1 microMol (0.7 micrograms/ml) and equivalent trough plasma levels (C^{ss}_{min}) were 1.8 microMol (0.4 micrograms/ml). Corresponding C^{ss}_{max} levels following doses of 400 mg and 800 mg administered four-hourly were 5.3 microMol (1.2 micrograms/ml) and 8 microMol (1.8 micrograms/ml)

respectively and equivalent C^{ss}_{min} levels were 2.7 microMol (0.6 micrograms/ml) and 4 microMol (0.9 micrograms/ml).

In adults the terminal plasma half-life of acyclovir after administration of intravenous aciclovir is about 2.9 hours. Most of the drug is excreted unchanged by the kidney. Renal clearance of

acyclovir is substantially greater than creatinine clearance, indicating that tubular secretion, in addition to glomerular filtration contributes to the renal elimination of the drug. 9-carboxymethoxymethylguanine is the only significant metabolite of acyclovir, and accounts for approximately 10-15% of the administered dose recovered from the urine. When acyclovir is given one hour after 1 gram of probenecid the terminal half-life and the area under the plasma concentration time curve is extended by 18% and 40% respectively.

In adults, mean steady state peak plasma concentrations (C^{ss}_{max}) following a one hour infusion of 2.5 mg/kg, 5 mg/kg and 10 mg/kg were 22.7 microMol (5.1 micrograms/ml), 43.6 microMol (9.8 micrograms/ml) and 92 microMol (20.7 micrograms/ml), respectively. The corresponding trough levels (C^{ss}_{min}) 7 hours later were 2.2 microMol (0.5 micrograms/ml),

3.1 microMol (0.7 micrograms/ml), and 10.2 microMol (2.3 micrograms/ml), respectively.

In children over 1 year of age similar mean peak (C^{ss}_{max}) and trough (C^{ss}_{min}) levels were observed when a dose of 250 mg/m² was substituted for 5 mg/kg and a dose of 500 mg/m² was substituted for 10 mg/kg. In neonates and young infants (0 to 3 months of age) treated with doses of 10 mg/kg administered by infusion over a one-hour period every 8 hours the C^{ss}_{max} was found to be 61.2 microMol (13.8 micrograms/ml) and C^{ss}_{min} to be 10.1 microMol (2.3 micrograms/ml). The terminal plasma half-life in these patients was 3.8 hours. A separate group of neonates treated with 15 mg/kg every 8 hours showed approximate dose proportional increases, with a C_{max} of 83.5 micromolar (18.8 microgram/ml) and C_{min} of 14.1 micromolar (3.2 microgram/ml).

In the elderly, total body clearance falls with increasing age associated with decreases in creatinine clearance although there is little change in the terminal plasma half-life.

In patients with chronic renal failure the mean terminal half-life was found to

be 19.5 hours. The mean acyclovir half-life during haemodialysis was 5.7 hours. Plasma acyclovir levels dropped approximately 60% during dialysis. Cerebrospinal fluid levels are approximately 50% of corresponding plasma levels. Plasma protein binding is relatively low (9 to 33%) and drug interactions involving binding site displacement are not anticipated.

5.3 Preclinical safety data

Mutagenicity:

The results of a wide range of mutagenicity tests *in vitro* and *in vivo* indicate that acyclovir is unlikely to pose a genetic risk to man.

Carcinogenicity:

Acyclovir was not found to be carcinogenic in long term studies in the rat and the mouse.

Teratogenicity:

Systemic administration of acyclovir in internationally accepted standard tests did not produce embryotoxic or teratogenic effects in rats, rabbits or mice.

In a non-standard test in rats, foetal abnormalities were observed, but only following such high subcutaneous doses that maternal toxicity was produced. The clinical relevance of these findings is uncertain.

Fertility:

Largely reversible adverse effects on spermatogenesis in association with overall toxicity in rats and dogs have been reported only at doses of acyclovir greatly in excess of those employed therapeutically. Two generation studies in mice did not reveal any effect of acyclovir on fertility.

6. Pharmaceutical particulars

6.1 List of excipients

Microcrystalline Cellulose (Avicel PH 101)
Microcrystalline Cellulose (Avicel PH 102)
Sodium Starch Glycolate (Primojel)
Povidone (K-30)
Magnesium Stearate

6.2 Incompatibilities

Not applicable

6.3 Shelf life

2 years

6.4 Special precautions for storage

Store below 30°C. Protect from light and moisture. Keep out of the reach of children.

6.5 Nature and contents of container

Box containing 20 tablets in Alu-PVDC blister pack.

6.6 Special precautions for disposal and other handling

Any unused product or waste material should be disposed of in accordance with local requirements.

7. Supplier

Square Pharmaceuticals PLC.

Square Centre

48, Mohakhali C.A, Dhaka – 1212,

Bangladesh

Tel: +880 2 8833047-56, +880 2 8859007 (10 lines)

Fax: +880 2 8828768, +880 2 8834941

E-mail: info@squaregroup.com

8. Market authorization number

H2011/21685/255

9. Date of first registration

17/01/2019

10. Date of revision of the text

19/07/2026

Reference list:

- https://www.accessdata.fda.gov/drugsatfda_docs/label/2005/018828s030,020089s019,0_19909s020lbl.pdf
- <https://www.squarepharma.com.bd/product-details.php?pid=305>