

Summary of Product Characteristics for Pharmaceutical Products

1. Name of the medicinal product:

Zerodol

2. Qualitative and quantitative composition

Each film coated tablet contains:

Aceclofenac BP 100mg

For a full list of excipients, see section 6.1

3. Pharmaceutical form

Tablet

4. Clinical particulars

4.1 Therapeutic indications

Aceclofenac is indicated for the relief of pain and inflammation associated with rheumatoid arthritis, osteoarthritis or ankylosing spondylitis.

4.2 Posology and method of administration

Posology

Aceclofenac tablets should be swallowed whole with a sufficient quantity of liquid. When aceclofenac was administered to fasting and fed healthy volunteers only the rate and not the extent of aceclofenac absorption was affected and as such aceclofenac should be taken preferably with or after food.

The usual dose of aceclofenac is 100 mg given twice daily by mouth.

One tablet in the morning and one in the evening.

Renal impairment

There is no evidence that the dosage of aceclofenac needs to be modified in patients with mild renal impairment, but as with other NSAIDs caution should be exercised.

Hepatic impairment

There is some evidence that the dose of aceclofenac should be reduced in patients with hepatic impairment and it is suggested that an initial daily dose of 100 mg be used.

4.3 Contraindications

Hypersensitivity to aceclofenac or to any of the excipients included in section 6.1. Aceclofenac should not be administered to patients with a history of gastrointestinal bleeding or perforation associated with previous treatment with NSAIDs, peptic ulcer / active or repeated gastrointestinal bleeding (two or more different proven episodes of ulceration or bleeding). It should not be administered in patients with active bleeding or bleeding disorders. This medication should not be administered to patients with serious renal or hepatic impairment. Airtal should not be administered to patients with established congestive cardiac impairment (NYHA classification II-IV), ischaemic cardiopathy, peripheral arterial pathology and/or cerebrovascular illness. Aceclofenac should not be prescribed in the third trimester of pregnancy. Airtal should not be administered during the first and second trimesters of pregnancy unless considered strictly necessary. In this case the dose and duration of treatment should be reduced as far as possible (see section 4.6). Aceclofenac should not be prescribed during lactation (see section 4.6). Airtal should not be administered in patients in whom acetylsalicylic acid or non-steroidal anti-inflammatory drugs trigger asthma attacks, acute rhinitis or urticaria, or in patients with hypersensitivity to these compounds

4.4 Special warnings and precautions for use

Close medical surveillance is imperative in patients with symptoms indicative of gastrointestinal disorders, with a history suggestive of gastrointestinal ulceration, with ulcerative colitis or with Crohn's disease, bleeding diathesis or haematological abnormalities.

Gastrointestinal bleeding or ulcerative perforation, haematemesis and melaena have in general more serious consequences in the elderly.

They can occur at any time during treatment, with or without warning symptoms or a previous history. In the rare instances, where gastrointestinal bleeding or ulceration occurs in patients receiving aceclofenac, the drug should be withdrawn.

Close medical surveillance is also imperative in patients suffering from severe impairment of hepatic function.

As with other NSAIDs, allergic reactions, including anaphylactic/ anaphylactoid reactions, can also occur without earlier exposure to the drug.

PRECAUTIONS

Patients with mild renal or cardiac impairment and the elderly should be kept under surveillance, since the use of NSAIDs may result in deterioration of renal function. The lowest effective dose should be used

and renal function monitored regularly.

The importance of prostaglandins in maintaining renal blood flow should be taken into account in patients with impaired cardiac or renal function, those being treated with diuretics or recovering from major surgery.

Effects on renal function are usually reversible on withdrawal of aceclofenac.

If abnormal liver function tests persist or worsen, clinical signs or symptoms consistent with liver disease develop or if other manifestations occur (eosinophilia, rash), aceclofenac should be discontinued. Hepatitis may occur without prodromal symptoms. Use of aceclofenac in patients with hepatic porphyria may trigger an attack. Aceclofenac may reversibly inhibit platelet aggregation.

Caution should also be exercised in patients with history of coagulation defects and history of liver dysfunction long term treatment.

NSAIDs should be given with care to patients with a history of heart failure or hypertension since oedema has been reported in association with NSAID administration.

Use with caution in patients suffering from or with a history of bronchial asthma since NSAIDs have been known to cause bronchospasm in such patients.

4.5 Interaction with other medicinal products and other forms of

Drug interactions associated with aceclofenac are similar to those observed with other NSAIDs.

Lithium: Aceclofenac, like many NSAIDs, may increase plasma concentration of lithium.

Cardiac glycosides: Through their renal effects, NSAIDs may increase plasma glycoside (including digoxin) levels, exacerbate cardiac failure and reduce the glomerular filtration rate in patients receiving glycosides.

Diuretics: Aceclofenac, like other NSAIDs may inhibit the activity of diuretics. Although it was not shown to affect blood pressure control when co-administered with bendrofluazide, interactions with other diuretics cannot be ruled out. When concomitant administration with potassium sparing diuretics is employed, serum potassium should be monitored. Diuretics can increase the risk of nephrotoxicity of NSAIDs.

Anticoagulants: Like other NSAIDs, aceclofenac may enhance the activity of anticoagulants. Close monitoring of patients on combined anticoagulant and aceclofenac therapy should be undertaken.

Antidiabetic agents: There have been isolated reports of hypoglycaemic

and hyperglycaemic effects. Thus with aceclofenac, consideration should be given to adjustment of the dosage of hypoglycaemic agents.

Methotrexate: Caution should be exercised if NSAIDs and methotrexate are administered within 24 hours of each other, since NSAIDs may increase methotrexate plasma levels, resulting in increased toxicity.

Mifepristone: NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone.

Other NSAIDs and Steroids: Concomitant therapy with aspirin, other NSAIDs and steroids may increase the frequency of adverse reactions, including the risk of GI bleeding.

Cyclosporin: Cyclosporin nephrotoxicity may be increased by the effect of NSAIDs on renal prostaglandins.

Quinolone antimicrobials: Convulsions may occur due to an interaction between quinolones and NSAIDs. This may occur in patients with or without a previous history of epilepsy or convulsions. Therefore, caution should be exercised when considering the use of a quinolone in patients who are already receiving a NSAID.

4.6 Pregnancy and Lactation

The drug is not recommended in pregnant or breast feeding women

4.7 Effects on ability to drive and use machines

Patients suffering from dizziness, vertigo, or other central nervous system disorders whilst taking NSAIDs should refrain from driving or handling dangerous machinery.

4.8 Undesirable effects

Aceclofenac is well tolerated, with most adverse events being minor and reversible.

The most frequent are gastrointestinal disorders, in particular dyspepsia, abdominal pain, nausea and diarrhea, and occasional occurrence of dizziness. Dermatological complaints including pruritus and rash and abnormal hepatic enzyme and serum creatinine levels have also been reported. If serious reactions occur, aceclofenac should be withdrawn.

Although the incidence of GI adverse events with aceclofenac was similar to those of comparator NSAIDs in individual clinical trials, withdrawal rates due to these events were significantly lower with aceclofenac than with ketoprofen and tenoxicam. Costs incurred as a result of adverse event management are lower with aceclofenac than with a range of comparator NSAIDs. Although statistical analyses were not consistently available, faecal bleeding and endoscopy studies in humans have indicated overall less GI bleeding and GI mucosal damage with aceclofenac than with

naproxen or diclofenac.

The following adverse events [described as common (<10% - > 1%), uncommon (<1%->0.1%), rare (<0.1% - >0.01%) and very rare/isolated reports (<0.01%)] were reported in clinical studies and post marketing surveillance

Blood and lymphatic system disorders: Rare: Anaemia. Very rare: Granulocytopenia, thrombocytopenia, neutropenia, haemolytic anaemia

Immune system disorders: Rare: Anaphylactic reaction (including shock), hypersensitivity

Metabolism and nutrition disorders: Very rare: Hyperkalemia

Psychiatric disorders: Very rare: Depression, abnormal dreams, insomnia

Nervous system disorders: Common: Dizziness. Very rare: Paraesthesia, tremor, somnolence, headache, dysgeusia (abnormal taste)

Eye disorders: Rare: Visual disturbance

Ear and labyrinth disorders: Very rare: Vertigo

Cardiac disorders: Very rare: Palpitations

Vascular disorders: Very rare: Flushing, hot flush

Respiratory, thoracic and mediastinal disorders: Rare: dyspnoea.

Very rare: Bronchospasm, stridor

Gastrointestinal disorders: Common: Dyspepsia, abdominal pain, nausea and diarrhea. Uncommon: Flatulence, gastritis, constipation, vomiting, mouth ulceration. Rare: Melaena. Very rare: Stomatitis, haematemesis, gastrointestinal haemorrhage, gastric ulcer, pancreatitis

Hepatobiliary disorders: Very rare: Hepatitis, jaundice

Skin and subcutaneous tissue disorders: Uncommon: Pruritus, rash, dermatitis, urticaria. Rare: Face oedema. Very rare: Purpura, bullous dermatitis

Renal and urinary disorders: Very rare: Renal insufficiency, nephritic syndrome

General disorders and administration site conditions: Very rare: Oedema, fatigue, cramps in legs

Laboratory investigations: Common: Increased hepatic enzyme.

Uncommon: Increased blood urea, increased blood creatinine. Very rare: Increased blood alkaline phosphatase, weight increase

Reporting of suspected adverse reactions

Reporting of suspected adverse reactions: Healthcare professionals are requested to report any suspected adverse reactions via

pharmacy and poisons board, Pharmacovigilance Electronic Reporting System(PvERS) <https://pv.pharmacyboardkenya.org>

4.9 Overdose

Treatment for acute intoxication due to non-steroidal anti-inflammatory drugs consists, essentially of support and symptomatic measures for complications such as hypotension, renal impairment, convulsions, gastrointestinal irritation and respiratory depression. There are no data regarding the consequences of overdose of Airtal in humans. Therapeutic measures to adopt are: in the case of overdose, avoid absorption of the drug as far as possible through gastric lavage and treatment with activated carbon. Specific treatments such as forced diuresis, dialysis or haemoperfusion probably do not contribute to the elimination of non-steroidal anti-inflammatory drugs due to their high rate of protein binding and extensive metabolism

5. Pharmacological properties

5.1 Pharmacodynamic properties

The mode of action of aceclofenac is largely based on the inhibition of prostaglandin synthesis. Aceclofenac is a potent inhibitor of the enzyme cyclo-oxygenase, which is involved in the production of prostaglandins.

Aceclofenac has been shown to exert effects on a variety of mediators of inflammation. The drug inhibits synthesis of the inflammatory cytokines interleukin (IL)-1 and tumour necrosis factor and inhibits prostaglandin E (PGE) production. Effects on cell adhesion molecules from 2 neutrophils have also been noted. *In vitro* data indicate inhibition of cyclo-oxygenase (COX)-1 and 2 by aceclofenac in whole blood assays, with selectivity for COX-2 being evident.

In contrast to some other NSAIDs, aceclofenac has shown stimulatory effects on cartilage matrix synthesis, that may be linked to the ability of the drug to inhibit IL-1 activity. *In vitro* data indicate stimulation by the drug of synthesis of glycosaminoglycan in osteoarthritic cartilage. There is also evidence that aceclofenac stimulates the synthesis of IL-1 receptor antagonist in human articular chondrocytes subjected to inflammatory stimuli and that 4'-hydroxyaceclofenac has chondroprotective properties attributable to suppression of IL-1 mediated pro-matrix metalloproteinase production and proteoglycan release.

In patients with osteoarthritis of the knee, aceclofenac decreases pain, reduces disease severity and improves the functional capacity of the knee. It reduces joint inflammation, pain intensity and the duration of morning stiffness in patients with rheumatoid arthritis. The duration of morning stiffness and pain intensity are reduced and spinal mobility improved, by aceclofenac in patients with ankylosing spondylitis

5.2 Pharmacokinetic properties

Aceclofenac is rapidly and completely absorbed after oral administration. Peak plasma concentrations are reached 1 to 3 hours after an oral dose. The drug is highly protein bound (>99%). The presence of food does not alter the extent of absorption of aceclofenac but the absorption rate is reduced. The plasma concentration of aceclofenac was approximately twice that in synovial fluid after multiple doses of the drug in patients with knee pain and synovial fluid effusion.

Aceclofenac is metabolised to a major metabolite, 4'-hydroxyaceclofenac and to a number of other metabolites including 5-hydroxyaceclofenac, 4'-hydroxydiclofenac, diclofenac and 5-hydroxydiclofenac. These other metabolites account for the fate of approximately 20% of each dose of aceclofenac. Renal excretion is the main route of elimination of aceclofenac with 70 to 80% of an administered dose found in the urine, mainly as the glucuronides of aceclofenac and its metabolites. Of each dose of aceclofenac, 20% is excreted in the faeces. The plasma elimination half life of the drug is approximately 4 hours.

5.3 Preclinical safety data

No additional data available.

6. Pharmaceutical Particulars

6.1 List of Excipients

Microcrystalline Cellulose BP/EP
Croscarmellose Sodium BP/EP
Sodium Starch Glycollate BP/EP
Colloidal Anhydrous Silica BP/EP
Sodium Lauryl Sulphate BP/EP
Ferric Oxide Red USP-NF
Hypromellose BP/EP
Polyoxy-40 Hydrogenated Castor oil USP-NF
Isopropyl alcohol BP/EP
Methylene Chloride EP
Titanium Dioxide BP/EP
Purified Talc BP/EP

Methylene Chloride EP
Dibutyl Phthalate BP/EP
Crospovidone BP/EP
Stearic Acid BP/EP

6.2 Shelf-Life

36 Months

6.3 Special Precautions for storage

Store at a temperature not exceeding 30°C. Protect from light

6.4 Nature and Content of container

Blister strip of 10 tablets

6.5 Special precautions for disposal and other handling

--

7. Marketing Authorization Holder

IPCA LABORATORIES LIMITED
Regd. Off.: 48, Kandivli Ind. Estate,
Mumbai 400 067, India.

Manufacturing Site

Ipca Laboratories Limited,

P.O Sejavta, Ratlam- 457001, Madhya Pradesh

8. Marketing Authorization Number

--

9. Date of first authorization/renewal of the authorization

--

10. Date of revision of the text

--